


1. Answer the following questions about magnesium.


A. An incomplete mass spectrum for magnesium is shown in the diagram.



The percent abundance of magnesium-24 is 79%. The percent abundances of the other two natural isotopes of magnesium, magnesium-25 and magnesium-26, are approximately equal.

- Complete the mass spectrum in part A by drawing thick lines in the appropriate locations to represent the percent abundance of magnesium-25 and magnesium-26.
- Describe the difference in atomic structure that accounts for the difference in mass between magnesium-25 and magnesium-26.

A student prepares a $1.85 \times 10^{-3} M$ solution of $\text{Mg}(\text{NO}_3)_2(aq)$ in beaker 1 and a $2.80 \times 10^{-4} M$ solution of $\text{NaOH}(aq)$ in beaker 2, as shown.




B. The particle diagram shown represents a magnesium ion, Mg^{2+} , in beaker 1. A sodium ion, Na^+ , in beaker 2 has a weaker attraction to water than the Mg^{2+} does. Explain this phenomenon using Coulomb's law and each of the following.


- The relative charge of the ions
- The relative radii of the ions

C. Calculate the pH of the solution in beaker 2.

- D. A student combines 35.00 mL of $1.85 \times 10^{-3} M \text{Mg}(\text{NO}_3)_2(aq)$ with 50.00 mL of $2.80 \times 10^{-4} M \text{NaOH}(aq)$, as shown in the diagram. Calculate $[\text{Mg}^{2+}]$ after the two solutions are combined but before any reaction takes place. (Assume that volumes are additive.)



- E. The dissolution of magnesium hydroxide is represented by the following equation.



- Write the expression for the solubility product constant, K_{sp} .
 - After the two solutions are combined in beaker 3 as described in part D, but before any reaction takes place, $[\text{OH}^-] = 1.65 \times 10^{-4} M$. Using your answer to part D, calculate the value of the reaction quotient, Q .
 - Using the reaction quotient, Q , predict whether a precipitate should form as the mixture in beaker 3 approaches equilibrium. Justify your answer.
- F. In a separate experiment, the student adds $\text{HNO}_3(aq)$ to decrease the pH of a saturated solution containing undissolved $\text{Mg}(\text{OH})_2(s)$. Does the amount of undissolved $\text{Mg}(\text{OH})_2(s)$ increase, decrease, or remain the same as the $\text{HNO}_3(aq)$ is added? Justify your answer.

6. A student is studying the properties of CaSO_4 and PbSO_4 . The student has samples of both compounds, which are white powders.

(a) The student tests the electrical conductivity of each solid and observes that neither solid conducts electricity. Describe the structures of the solids that account for their inability to conduct electricity.


The student places excess $\text{CaSO}_4(s)$ in a beaker containing 100 mL of water and places excess $\text{PbSO}_4(s)$ in another beaker containing 100 mL of water. The student stirs the contents of the beakers and then measures the electrical conductivity of the solution in each beaker. The student observes that the conductivity of the solution in the beaker containing the $\text{CaSO}_4(s)$ is higher than the conductivity of the solution in the beaker containing the $\text{PbSO}_4(s)$.

(b) Which compound is more soluble in water, $\text{CaSO}_4(s)$ or $\text{PbSO}_4(s)$? Justify your answer based on the results of the conductivity test.

GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

The left side of the diagram below shows a particulate representation of the contents of the beaker containing the $\text{CaSO}_4(s)$ from the solution conductivity experiment.




(c) Draw a particulate representation of $\text{PbSO}_4(s)$ and the ions dissolved in the solution in the beaker on the right in the diagram. Draw the particles to look like those shown to the right of the beaker. Draw an appropriate number of dissolved ions relative to the number of dissolved ions in the beaker on the left.


(d) The student attempts to increase the solubility of $\text{CaSO}_4(s)$ by adding 10.0 mL of 2 M $\text{H}_2\text{SO}_4(aq)$ to the beaker, and observes that additional precipitate forms in the beaker. Explain this observation.

GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.



3. To determine the molar mass of an unknown metal, M, a student reacts iodine with an excess of the metal to form the water-soluble compound MI_2 , as represented by the equation above. The reaction proceeds until all of the I_2 is consumed. The $\text{MI}_2(aq)$ solution is quantitatively collected and heated to remove the water, and the product is dried and weighed to constant mass. The experimental steps are represented below, followed by a data table.



Data for Unknown Metal Lab	
Mass of beaker	125.457 g
Mass of beaker + metal M	126.549 g
Mass of beaker + metal M + I_2	127.570 g
Mass of MI_2 , first weighing	1.284 g
Mass of MI_2 , second weighing	1.284 g

- (a) Given that the metal M is in excess, calculate the number of moles of I_2 that reacted.
- (b) Calculate the molar mass of the unknown metal M.

The student hypothesizes that the compound formed in the synthesis reaction is ionic.

- (c) Propose an experimental test the student could perform that could be used to support the hypothesis. Explain how the results of the test would support the hypothesis if the substance was ionic.



The student hypothesizes that Br_2 will react with metal M more vigorously than I_2 did because Br_2 is a liquid at room temperature.

- (d) Explain why I_2 is a solid at room temperature whereas Br_2 is a liquid. Your explanation should clearly reference the types and relative strengths of the intermolecular forces present in each substance.

While cleaning up after the experiment, the student wishes to dispose of the unused solid I_2 in a responsible manner. The student decides to convert the solid I_2 to $\text{I}^-(aq)$ anion. The student has access to three solutions, $\text{H}_2\text{O}_2(aq)$, $\text{Na}_2\text{S}_2\text{O}_3(aq)$, and $\text{Na}_2\text{S}_4\text{O}_6(aq)$, and the standard reduction table shown below.

Half reaction	E° (V)
$\text{S}_4\text{O}_6^{2-}(aq) + 2 e^- \rightarrow 2 \text{S}_2\text{O}_3^{2-}(aq)$	0.08
$\text{I}_2(s) + 2 e^- \rightarrow 2 \text{I}^-(aq)$	0.54
$\text{O}_2(g) + 2 \text{H}^+(aq) + 2 e^- \rightarrow \text{H}_2\text{O}_2(aq)$	0.68

- (e) Which solution should the student add to $\text{I}_2(s)$ to reduce it to $\text{I}^-(aq)$? Circle your answer below. Justify your answer, including a calculation of E° for the overall reaction.



- (f) Write the balanced net-ionic equation for the reaction between I_2 and the solution you selected in part (e).