


1. Answer the following questions about magnesium.


A. An incomplete mass spectrum for magnesium is shown in the diagram.



The percent abundance of magnesium-24 is 79%. The percent abundances of the other two natural isotopes of magnesium, magnesium-25 and magnesium-26, are approximately equal.

- i. Complete the mass spectrum in part A by drawing thick lines in the appropriate locations to represent the percent abundance of magnesium-25 and magnesium-26.
- ii. Describe the difference in atomic structure that accounts for the difference in mass between magnesium-25 and magnesium-26.

A student prepares a $1.85 \times 10^{-3} M$ solution of $\text{Mg}(\text{NO}_3)_2(aq)$ in beaker 1 and a $2.80 \times 10^{-4} M$ solution of $\text{NaOH}(aq)$ in beaker 2, as shown.




B. The particle diagram shown represents a magnesium ion, Mg^{2+} , in beaker 1. A sodium ion, Na^+ , in beaker 2 has a weaker attraction to water than the Mg^{2+} does. Explain this phenomenon using Coulomb's law and each of the following.


- i. The relative charge of the ions
- ii. The relative radii of the ions

C. Calculate the pH of the solution in beaker 2.


- D. A student combines 35.00 mL of $1.85 \times 10^{-3} M \text{Mg}(\text{NO}_3)_2(aq)$ with 50.00 mL of $2.80 \times 10^{-4} M \text{NaOH}(aq)$, as shown in the diagram. Calculate $[\text{Mg}^{2+}]$ after the two solutions are combined but before any reaction takes place. (Assume that volumes are additive.)




- E. The dissolution of magnesium hydroxide is represented by the following equation.




- Write the expression for the solubility product constant, K_{sp} .
 - After the two solutions are combined in beaker 3 as described in part D, but before any reaction takes place, $[\text{OH}^{-}] = 1.65 \times 10^{-4} M$. Using your answer to part D, calculate the value of the reaction quotient, Q .
 - Using the reaction quotient, Q , predict whether a precipitate should form as the mixture in beaker 3 approaches equilibrium. Justify your answer.
- F. In a separate experiment, the student adds $\text{HNO}_3(aq)$ to decrease the pH of a saturated solution containing undissolved $\text{Mg}(\text{OH})_2(s)$. Does the amount of undissolved $\text{Mg}(\text{OH})_2(s)$ increase, decrease, or remain the same as the $\text{HNO}_3(aq)$ is added? Justify your answer.



2. A student investigates the reactions of nitrogen oxides. One of the reactions in the investigation requires an equimolar mixture of $\text{NO}(g)$ and $\text{NO}_2(g)$, which the student produces by using the reaction represented above.
- (a) The particle-level representation of the equimolar mixture of $\text{NO}(g)$ and $\text{NO}_2(g)$ in the flask at the completion of the reaction between $\text{NO}(g)$ and $\text{O}_2(g)$ is shown below in the box on the right. In the box below on the left, draw the particle-level representation of the reactant mixture of $\text{NO}(g)$ and $\text{O}_2(g)$ that would yield the product mixture shown in the box on the right. In your drawing, represent oxygen atoms and nitrogen atoms as indicated below.




The student reads in a reference text that $\text{NO}(g)$ and $\text{NO}_2(g)$ will react as represented by the equation below. Thermodynamic data for the reaction are given in the table below the equation.



ΔH_{298}°	ΔS_{298}°	ΔG_{298}°
$-40.4 \text{ kJ/mol}_{\text{rxn}}$	$-138.5 \text{ J/(K} \cdot \text{mol}_{\text{rxn}})$	$0.87 \text{ kJ/mol}_{\text{rxn}}$

- (b) The student begins with an equimolar mixture of $\text{NO}(g)$ and $\text{NO}_2(g)$ in a rigid reaction vessel and the mixture reaches equilibrium at 298 K.
- Calculate the value of the equilibrium constant, K , for the reaction at 298 K.
 - If both P_{NO} and P_{NO_2} in the vessel are initially 1.0 atm, will $P_{\text{N}_2\text{O}_3}$ at equilibrium be equal to 1.0 atm? Justify your answer.
- (c) The student hypothesizes that increasing the temperature will increase the amount of $\text{N}_2\text{O}_3(g)$ in the equilibrium mixture. Indicate whether you agree or disagree with the hypothesis. Justify your answer.

$\text{N}_2\text{O}_3(g)$ reacts with water to form nitrous acid, $\text{HNO}_2(aq)$, a compound involved in the production of acid rain. The reaction is represented below.




(d) The skeletal structure of the HNO_2 molecule is shown in the box below.

- (i) Complete the Lewis electron-dot diagram of the HNO_2 molecule in the box below, including any lone pairs of electrons.




- (ii) Based on your completed diagram above, identify the hybridization of the nitrogen atom in the HNO_2 molecule.


To produce an aqueous solution of HNO_2 , the student bubbles $\text{N}_2\text{O}_3(g)$ into distilled water. Assume that the reaction goes to completion and that HNO_2 is the only species produced. To determine the concentration of $\text{HNO}_2(aq)$ in the resulting solution, the student titrates a 100. mL sample of the solution with 0.100 M $\text{KOH}(aq)$. The neutralization reaction is represented below.



The following titration curve shows the change in pH of the solution during the titration.



- (e) Use the titration curve and the information above to
- determine the initial concentration of the $\text{HNO}_2(aq)$ solution
 - estimate the value of $\text{p}K_a$ for $\text{HNO}_2(aq)$
- (f) During the titration, after a volume of 15 mL of 0.100 M $\text{KOH}(aq)$ has been added, which species, $\text{HNO}_2(aq)$ or $\text{NO}_2^-(aq)$, is present at a higher concentration in the solution? Justify your answer.



$$K = 7.7 \times 10^7$$

6. The polyatomic ion $\text{C}_{10}\text{H}_{12}\text{N}_2\text{O}_8^{4-}$ is commonly abbreviated as EDTA^{4-} . The ion can form complexes with metal ions in aqueous solutions. A complex of EDTA^{4-} with Ba^{2+} ion forms according to the equation above. A 50.0 mL volume of a solution that has an $\text{EDTA}^{4-}(\text{aq})$ concentration of 0.30 M is mixed with 50.0 mL of 0.20 M $\text{Ba}(\text{NO}_3)_2$ to produce 100.0 mL of solution.
- (a) Considering the value of K for the reaction, determine the concentration of $\text{Ba}(\text{EDTA})^{2-}(\text{aq})$ in the 100.0 mL of solution. Justify your answer.
- (b) The solution is diluted with distilled water to a total volume of 1.00 L. After equilibrium has been reestablished, is the number of moles of $\text{Ba}^{2+}(\text{aq})$ present in the solution greater than, less than, or equal to the number of moles of $\text{Ba}^{2+}(\text{aq})$ present in the original solution before it was diluted? Justify your answer.