

3. The Rocky Mountains, the headwaters for the Colorado River, have experienced a severe drought for many of the past twenty years. States and cities downstream that depend on the river water have had to find solutions to the reduced water flow.

One city located downstream from the Rocky Mountains is Las Vegas, Nevada. In Las Vegas, one solution to reduce the amount of water used for irrigation is to remove landscaping and lawns that use grasses not native to the local area and replace them with desert-tolerant landscaping.

- (a) **Identify** a typical type of plant that would be used in desert-tolerant landscaping.
- (b) **Justify** the removal of landscaping and lawns that use grasses not native to the local area by describing an additional advantage, other than reducing the amount of water needed for irrigation.

Warmer seasonal temperatures in the Rocky Mountains have led to earlier mountain snowmelt and an increased risk of forest fires as soil and vegetation dry out.

- (c) **Describe** how a prescribed burn would reduce the severity and spread of forest fires.
- (d) **Describe** a disadvantage of a prescribed burn in a forested ecosystem.

Lake Powell is a large reservoir on the Colorado River. Lake Powell is formed by a dam with a hydroelectric power plant. At full capacity or full pool, Lake Powell contains 25.16 million acre-feet (maf) of water. Currently, Lake Powell is at 36% full capacity.

- (e) **Calculate** the number of million acre-feet (maf) of water currently in Lake Powell. **Show** your work.

In 2021, after years of drought, Lake Powell held 7.9 million acre-feet (maf) of water. One million acre-feet is equivalent to 3.26×10^{11} gallons.

- (f) The watershed of the Upper Colorado River contributes an average of 9.60 maf of water to Lake Powell annually. The melted mountain snow found in the watershed contributes 50% of the average river flow into Lake Powell. In 2021, the river flow was 36% of the annual average. Assuming that all resources to that flow contributed the same proportions as in prior years, **calculate** the amount of water (in million acre-feet) that was contributed by the melted mountain snow in 2021. **Show** your work.
- (g) The average household in the United States consumes 5.0×10^4 gallons of water in a year. **Calculate** the number of households that could be supported for one year by the average flow of 9.60 maf of water into Lake Powell. **Show** your work.

Begin your response to this question at the top of a new page in the separate Free Response booklet and fill in the appropriate circle at the top of each page to indicate the question number.

STOP

END OF EXAM

ENVIRONMENTAL SCIENCE

SECTION II


Time—1 hour and 10 minutes

3 Questions

Directions: Answer all three questions, which are weighted equally; the suggested time is about 22 minutes for answering each question. Write all your answers in the Free Response booklet. Where calculations are required, clearly show how you arrived at your answer. Where explanation or discussion is required, support your answers with relevant information and/or specific examples. You may plan your answers in this orange booklet, but no credit will be given for anything written in this booklet. **You will only earn credit for what you write in the separate Free Response booklet.**

1. Two restoration methods designed to reclaim the overburden piles removed from open pit coal mines in temperate forests of eastern Europe were studied over several years. The first method was spontaneous restoration, where the soil and rock were dumped in noncompacted, random piles and not replanted with vegetation. The second method was technical restoration, where the land was leveled by heavy machinery, organic material was added, and eight plant species were added. Technical restoration started 5 years after the piles were formed, so no bar is included in the 1–5 year range in the graph.

Number of Plant Species in Spontaneous and Technically Restored Overburden Piles over Time



- (a) Based on the data in the graph, **identify** the average number of species found in technically restored piles 16–25 years after restoration started.
- (b) Based on the data in the graph, **describe** the relationship between the average number of species and the age of the spontaneous restoration piles.

Scientists hypothesized that after 35 years, the number of animal species in the spontaneously restored piles would be higher than the number of animal species in the technically restored piles.

- (c) Using the data in the graph, **make a claim** that supports or refutes the hypothesis that the number of animal species in the spontaneously restored piles would be higher than the number of animal species in the technically restored piles.
- (d) **Describe** a characteristic of a plant species that would have been present in the spontaneous restoration piles from years 1 to 5.
- (e) **Explain** how the return of plant life would have been altered in the spontaneous piles if the mining had taken place in the tundra rather than in temperate forests.

Limiting runoff on disturbed land improves soil and water quality. To model water runoff, students set up three large plastic trays and filled each tray with loam soil to 1 centimeter below the edge. Loam soil is a balance of sand, silt, and clay soil. One tray was seeded with wheatgrass and grown for 30 days, one tray was covered with small wood chips, and one tray was soil only. Five hundred milliliters of water was sprinkled evenly onto the soil surface of each tray. All three trays were placed on a slight slope, and water that ran off the surface was collected. The results are shown in the following table.

Tray	Water Collected (mL)	Water Appearance
Wheatgrass grown in soil; no small wood chips	190	Clear
Small wood chips covered soil; no wheatgrass	300	Transparent; light brown
Soil only; no wheatgrass or small wood chips	360	Cloudy; dark brown


- (f) **Identify** the scientific question for the investigation.
- (g) **Identify** a dependent variable in the experiment.
- (h) **Explain** how the results of the experiment would likely have been altered if the soil had contained a higher percentage of sand, which has the largest particle size of the components of loam.

Runoff from mining waste can also affect water quality.

- (i) **Describe** one ecological impact that could result from the drainage from mine waste into nearby waterways.
- (j) **Describe** one strategy to reduce the drainage from mine waste into waterways downstream from mines.

Begin your response to this question at the top of a new page in the separate Free Response booklet and fill in the appropriate circle at the top of each page to indicate the question number.

3. An Arctic food web includes the following organisms.



Note: Figures not drawn to scale.

(a) Refer to the food web above to complete the following table.

	Organism from Arctic food web
(i) Identify a primary producer	
(ii) Identify a primary consumer	
(iii) Identify a secondary consumer	

(b) Other than showing which organisms are consumed by other organisms, **describe** what is indicated by the direction of the arrows in the diagram.

As the amount of sea ice has decreased, larger expanses of the Arctic Ocean are now completely free of sea ice for several weeks each summer. Ringed seals, the preferred prey of polar bears, come to holes in the sea ice to breathe.

- (c) **Describe** how the change in sea ice habitat is affecting polar bears' ability to hunt and feed.
- (d) **Explain** how melting sea ice leads to a feedback loop that increases Arctic warming.
- (e) Many species, including some whales and birds, will travel thousands of kilometers during annual migrations.
 - (i) **Provide** one reason a species may migrate a long distance.
 - (ii) The North Atlantic right whale migrates between subtropical and polar waters annually. Nearly 50 percent of right whale deaths are due to human activities. **Describe** one commercial activity, other than whaling, that may result in the death of right whales.
 - (iii) **Describe** one strategy that could reasonably be implemented to decrease right whale deaths caused by the commercial activity you described in part (ii).