

There is a natural climate phenomenon that occurs in the equatorial Pacific Ocean because of periodic fluctuations in sea surface conditions and atmospheric circulation. This phenomenon happens every few years, resulting in shifts in temperature and precipitation in many parts of the world.

Figure 1 shows temperature and precipitation patterns associated with one phase of this phenomenon in the equatorial Pacific Ocean. Figure 2 shows the associated temperature and precipitation patterns that occur over North America.

Figure 1. Effects of Changes in Sea Surface Conditions in Equatorial Pacific Ocean





Figure 2. Polar Jet Stream and Climate Conditions in North America



2. Respond to parts A, B, C, D, E, F, G, H, I, and J.

- A. **Identify** the sea surface condition for the eastern equatorial area of the Pacific Ocean illustrated in Figure 1.
- B. Based on the information in Figure 1, **identify** the climate phenomenon associated with the sea surface conditions shown in the equatorial area of the Pacific Ocean.
- C. Based on the information in Figure 2, **describe** a difference in climate patterns between the regions A and B.
- D. Increases in regional precipitation can lead to increased flooding in certain locations. **Describe** one reason why there is an increased risk of flooding in urban areas during extended periods of precipitation.
- E. **Propose** a realistic solution a city could implement to decrease the risk of flooding in urban areas.
- F. **Justify** the solution proposed in part E by providing an additional advantage other than a reduction in the risk of flooding.
- G. Terrestrial biomes are primarily determined by temperature and precipitation patterns. **Describe** one difference between the climate of a temperate seasonal forest and that of a savanna.
- H. Droughts are one environmental problem that can affect the frequency and severity of forest fires. **Identify** the ecological process that occurs following a forest fire that leaves the soil intact.
- I. **Describe** one way burning forests contribute to atmospheric pollution.
- J. **Describe** one sustainable forestry practice that could be used to reduce the occurrence or severity of forest fires.

ENVIRONMENTAL SCIENCE


SECTION II

Time—1 hour and 10 minutes

3 Questions


Directions: Answer all three questions, which are weighted equally; the suggested time is about 22 minutes for answering each question. Write all your answers in the Free Response booklet. Where calculations are required, clearly show how you arrived at your answer. Where explanation or discussion is required, support your answers with relevant information and/or specific examples. You may plan your answers in this orange booklet, but no credit will be given for anything written in this booklet. **You will only earn credit for what you write in the separate Free Response booklet.**

1. The following diagram shows how an area changed over time. The dominant vegetation in each stage is labeled in the diagram. Each roman numeral represents a stage in the ecological process.



- (a) **Identify** the ecological change over time illustrated in the diagram.
- (b) **Identify** the stage in which pioneer species first appear.
- (c) **Describe** how soil forms from stage II to stage III.
- (d) **Identify** a cultural ecosystem service that may be provided by a mature forest.

Researchers are interested in investigating the effect of the removal of trees on the water quality of nearby waterways. To conduct their experiment, the researchers locate two identical forest watersheds, both with a river running through them. Forest A is an intact forest and forest B is a clear-cut forest. Four stream monitoring stations are set up in each forest study area. Each station monitors a variety of water quality indicators, including water temperature, at various locations along the river. The experimental setup is shown in the following diagrams. The arrows shown in the rivers represent the direction of water flow. The triangles in the river show the location of the stream monitoring stations.



- (e) **Identify** a testable hypothesis being investigated by the researchers.
- (f) **Identify** the independent variable in the experiment.
- (g) **Describe** the purpose of forest A in the experiment.
- (h) **Explain** why the water temperature might change because of clear-cutting trees in forest B.

After the experiment was completed, land developers constructed a golf course in the clear-cut area of forest B. To prepare the land, developers removed any remaining stumps and newly grown trees and began planting and fertilizing new grass. The researchers are interested in the effect the golf course may have on the water quality of the river. To monitor the river in forest B, the researchers repeated the experiment once the golf course had been constructed.

- (i) **Explain** why an indicator of water quality other than temperature could be altered by the golf course in forest B.
- (j) **Explain** how the aquatic organisms living in the lake illustrated in the diagram of forest B are likely to be affected by the new golf course.

Begin your response to this question at the top of a new page in the separate Free Response booklet and fill in the appropriate circle at the top of each page to indicate the question number.

ENVIRONMENTAL SCIENCE
SECTION II
Time—1 hour and 10 minutes
3 Questions

Directions: Answer all three questions, which are weighted equally; the suggested time is about 22 minutes for answering each question. Write all your answers in the Free Response booklet. Where calculations are required, clearly show how you arrived at your answer. Where explanation or discussion is required, support your answers with relevant information and/or specific examples. You may plan your answers in this orange booklet, but no credit will be given for anything written in this booklet. **You will only earn credit for what you write in the separate Free Response booklet.**

- 1. Researchers interested in sustainability developed three new varieties of genetically modified green beans designed to produce higher yields in arid regions. Four plots (A–D) were set up on a floodplain of a river to grow beans. Plots A, B, and C were each planted with seeds of different types of genetically modified strains of green beans. Plot D was planted with seeds of unmodified green beans. Each plot was given equal amounts of fertilizer, which contains phosphorus and nitrogen, and water and was irrigated using spray irrigation for one hour per day. Throughout the growing season, researchers measured the amount of green beans harvested within each plot. The plots with genetically modified beans had higher crop yields than the plot with unmodified beans.
 - (a) **Identify** the control group in this experiment.
 - (b) **Identify** the scientific question for the investigation.
 - (c) Researchers repeated the experiment by modifying the length of time for the spray irrigation to 20 minutes per day. **Explain** how the results of the experiment could be altered with this modification.
- Researchers also monitored the amount of sediment and fertilizer washing into the river from each plot in the original study. The soil temperature was also measured in each plot at the same depth and at the same time each day approximately 0.05 meters below the surface of the soil.

Genetically Modified Green Bean Experimental Results

Plot	Sediment Runoff (mg/L)	Phosphorus Runoff (mg/L)	Nitrogen Runoff (mg/L)	Soil Temperature (°C)
A (Type 1 GMO beans)	2.4	0.11	1.07	18.6
B (Type 2 GMO beans)	2.1	0.02	0.56	18.3
C (Type 3 GMO beans)	0.9	0.04	0.68	18.3
D (Unmodified beans)	3.5	0.15	1.24	17.6

- (d) Based on the data in the table, **identify** the plot with the lowest soil temperature.
- (e) **Describe** how sediment runoff and fertilizer runoff compare between the unmodified green beans and the genetically modified green beans.

- (f) The Type 2 GMO beans in Plot B were developed to grow more quickly than the unmodified beans in Plot D. Researchers have hypothesized that the Type 2 beans would use fertilizer more completely than the other varieties. Based on the data in the table and the experimental design, **explain** whether the researchers' hypothesis was supported or refuted.

Once the experiment was concluded, the researchers burned the plots to remove the crops that had been planted. After a few years, the researchers returned to the plots and observed a variety of plants, insects, and bird species living there.

- (g) **Describe** the ecological process that occurred on the plots after the crops were burned.

A survey indicated that one of the plots had twice the plant diversity that the other plots had. Over the next five years, the river occasionally flooded the plots, killing off many of the species that inhabited the plots.

- (h) After each flooding event, the plot with twice the plant diversity returned to its prior level of biodiversity more quickly than the other plots did. **Explain** why a community with more plant diversity will recover more quickly from the flooding.

- (i) After the last flooding event, a beetle not previously known in the area appeared in one of the plots with less plant diversity. Over a period of a few months, the new beetle population increased, whereas the existing beetle species in the plot had declining populations. **Explain** why the new beetle species could be better able to successfully populate this plot than the existing beetle species could.

- (j) **Describe** one realistic method to prevent the new beetle from spreading beyond the experimental plot.

Begin your response to this question at the top of a new page in the separate Free Response booklet and fill in the appropriate circle at the top of each page to indicate the question number.

ENVIRONMENTAL SCIENCE

SECTION II


Time—1 hour and 10 minutes

3 Questions

Directions: Answer all three questions, which are weighted equally; the suggested time is about 22 minutes for answering each question. Write all your answers in the Free Response booklet. Where calculations are required, clearly show how you arrived at your answer. Where explanation or discussion is required, support your answers with relevant information and/or specific examples. You may plan your answers in this orange booklet, but no credit will be given for anything written in this booklet. **You will only earn credit for what you write in the separate Free Response booklet.**

- Two restoration methods designed to reclaim the overburden piles removed from open pit coal mines in temperate forests of eastern Europe were studied over several years. The first method was spontaneous restoration, where the soil and rock were dumped in noncompacted, random piles and not replanted with vegetation. The second method was technical restoration, where the land was leveled by heavy machinery, organic material was added, and eight plant species were added. Technical restoration started 5 years after the piles were formed, so no bar is included in the 1–5 year range in the graph.

Number of Plant Species in Spontaneous and Technically Restored Overburden Piles over Time



- Based on the data in the graph, **identify** the average number of species found in technically restored piles 16–25 years after restoration started.
- Based on the data in the graph, **describe** the relationship between the average number of species and the age of the spontaneous restoration piles.

Scientists hypothesized that after 35 years, the number of animal species in the spontaneously restored piles would be higher than the number of animal species in the technically restored piles.

- (c) Using the data in the graph, **make a claim** that supports or refutes the hypothesis that the number of animal species in the spontaneously restored piles would be higher than the number of animal species in the technically restored piles.
- (d) **Describe** a characteristic of a plant species that would have been present in the spontaneous restoration piles from years 1 to 5.
- (e) **Explain** how the return of plant life would have been altered in the spontaneous piles if the mining had taken place in the tundra rather than in temperate forests.

Limiting runoff on disturbed land improves soil and water quality. To model water runoff, students set up three large plastic trays and filled each tray with loam soil to 1 centimeter below the edge. Loam soil is a balance of sand, silt, and clay soil. One tray was seeded with wheatgrass and grown for 30 days, one tray was covered with small wood chips, and one tray was soil only. Five hundred milliliters of water was sprinkled evenly onto the soil surface of each tray. All three trays were placed on a slight slope, and water that ran off the surface was collected. The results are shown in the following table.

Tray	Water Collected (mL)	Water Appearance
Wheatgrass grown in soil; no small wood chips	190	Clear
Small wood chips covered soil; no wheatgrass	300	Transparent; light brown
Soil only; no wheatgrass or small wood chips	360	Cloudy; dark brown

- (f) **Identify** the scientific question for the investigation.
- (g) **Identify** a dependent variable in the experiment.
- (h) **Explain** how the results of the experiment would likely have been altered if the soil had contained a higher percentage of sand, which has the largest particle size of the components of loam.
- Runoff from mining waste can also affect water quality.
- (i) **Describe** one ecological impact that could result from the drainage from mine waste into nearby waterways.
- (j) **Describe** one strategy to reduce the drainage from mine waste into waterways downstream from mines.

Begin your response to this question at the top of a new page in the separate Free Response booklet and fill in the appropriate circle at the top of each page to indicate the question number.

ENVIRONMENTAL SCIENCE

SECTION II

Time—1 hour and 30 minutes


4 Questions

Directions: Answer all four questions, which are weighted equally; the suggested time is about 22 minutes for answering each question. Write all your answers on the pages following the questions in this book. Where calculations are required, clearly show how you arrived at your answer. Where explanation or discussion is required, support your answers with relevant information and/or specific examples.

1. The piping plover is a globally threatened species, with perhaps as few as 2,000 nesting pairs remaining in the Atlantic population of these migratory shorebirds. Plovers were nearly hunted to extinction in the nineteenth century.

Plovers prefer to nest and search for food on open sandy beaches between sparsely vegetated sand dunes. Washovers, where sand is washed up from the beach and buries vegetated areas during big storms, are also important to plovers because they provide moist and undisturbed habitat for plover adults and their chicks. Ecologists with the Elliott Oceanographic Institute mapped the distribution of plover nests during two seasonal surveys of nesting sites over a ten-year period on Assateague Island. This is a barrier island, a long, narrow island running parallel to the Atlantic coast of Virginia.


SPRING PIPING PLOVER NESTING SITES
ON ASSATEAGUE ISLAND, VIRGINIA



- (a) Use the maps provided to answer the following questions.
- Identify** the preferred nesting habitat for piping plovers.
 - Describe** the change in the number of piping plover nests on Assateague Island between 1999 and 2009.
 - Describe** one likely reason for the change in the number of piping plover nests between 1999 and 2009.
- (b) Coastal species are affected by more than just natural events.
- Special beach restrictions can help piping plovers during nesting season. **Describe** one restriction that could reasonably be implemented to help prevent the destruction of plover nests by human actions.
 - In addition to providing habitat for piping plovers, barrier islands (and closely related landforms) are important for other reasons. **Explain** one way that these features help to preserve and protect the environment in coastal regions.
 - Identify** one human action that directly threatens coastal habitats and **describe** one impact on species, other than the piping plover, that use the habitat.

Approximately 40% of the United States population resides in coastal areas, such as areas near Assateague Island, where sea level rise and shoreline erosion is occurring.

- Identify** one economic impact on coastal communities that has resulted from rising sea levels.
- Describe** TWO methods that may be used locally to protect coastal communities from rising sea levels.



3. Plate-tectonic theory states that the Earth's lithosphere is broken into very slowly moving pieces or plates. Plate movements over vast stretches of time have led to the current orientation of our continents and oceans. Individual events along plate boundaries, such as earthquakes and volcanic eruptions, pose periodic threats to human activity and ecosystems. The "Ring of Fire" is a term that describes the location of increased seismic and volcanic activity around the margins of the Pacific Ocean basin. On the map above, each dot represents a volcano or an earthquake.
- Japan, Indonesia, and the Philippines are examples of volcanic island chains that have formed along subduction zones between plates in the western Pacific.
 - Describe** what happens when two tectonic plates collide along a subduction zone.
 - Explain** how subduction leads to volcanic activity.
 - Although the landscape following a volcanic eruption may appear unable to support ecological communities, over time the area can be transformed through succession.
 - What is primary succession?
 - Explain** how primary succession can lead to soil formation on a newly formed volcanic landscape.
 - In addition to volcanic activity, highly destructive tsunamis are generated along Pacific Plate subduction zones.
 - Explain** how a tsunami is generated along a subduction zone.
 - Describe** one negative ecological impact that tsunamis have on coastal environments.
 - Southern California experiences periodic devastating earthquakes along the San Andreas Fault, which is a transform boundary located along the eastern edge of the Pacific Plate.
 - Describe** what happens to the tectonic plates along a transform boundary at the moment when an earthquake occurs.
 - Describe** what happens to the tectonic plates along a transform boundary during the time between earthquakes.