


The following information applies to parts A, B, C, D, and E.

Climate change can affect both terrestrial and marine organisms. Researchers studying the effect of climate change on bird distribution mapped elevation zones at which five bird species lived within a tropical mountain range. Birds are distributed in part according to climate-related adaptations. Researchers modeled where the birds would most likely live in the future based on climate change predictions.

The results of this study are shown in the following diagram. Each unique shape in the diagram represents a different bird species, labeled A through E. Current distribution is shown on the left side of the figure, and predicted future distribution is shown on the right.

**The following information applies to parts F, G, and H.**

Some organisms such as birds can move in response to environmental changes, while other organisms such as corals cannot. Researchers studying the effect of climate change on marine organisms conducted two experiments involving photosynthetic algae that have a symbiotic relationship with corals.

In Experiment 1, the researchers placed photosynthetic algal colonies living in corals in three different saltwater tanks. The temperature of Tank 1 was set to 27°C, the average temperature at which most corals live. Water circulation in each tank mimicked natural ocean currents. After 30 days, the researchers counted the number of bleached corals. The tank setups are shown in the following list.

- Tank 1: 27°C and no sediment
- Tank 2: 32°C and no sediment
- Tank 3: 37°C and no sediment

In Experiment 2, the researchers used the same photosynthetic algae living in corals but modified the conditions of the tanks as shown in the following list. Water circulation kept the sediment in the tanks suspended. After 30 days, the researchers counted the number of bleached corals.

- Tank 1: 27°C and no sediment
- Tank 2: 27°C and moderate sediment
- Tank 3: 27°C and high sediment

1. Respond to parts A, B, C, D, E, F, G, H, I, and J.

Part A

Based on the information in the diagram, **identify** the elevation zone that has the highest species richness for the current distribution.

Part B

Describe a climate condition at high elevations that is different than it is at low elevations.

Part C

A predator at low elevations currently feeds on Species C and D. Based on the information in the diagram, **describe** what could happen to the predator species in the future in response to the predicted changes in the prey community.

Part D

Species C is considered a generalist, while Species E is considered a specialist. **Explain** why Species C and E are predicted to respond differently to climate change.

Part E

Based on the information in the diagram, **explain** why the bird community at low elevations in the future would be less likely to recover from natural disruptions than the current low-elevation bird community.

Part F

Identify the independent variable in Experiment 1.

Part G

Identify a likely scientific question for Experiment 1.

Part H

In Experiment 2 the temperature of the tanks was held constant, while the amount of sediment was varied. **Explain** how this modification could affect the results of this experiment.

Part I


Changes in climate also affect Arctic ecosystems. One impact of climate change is the loss of ice and snow in polar regions. **Describe** a positive feedback loop that results from the loss of ice and snow in polar regions.

Part J

Describe one problem that Arctic species, such as polar bears or ringed seals, face because of warming in polar regions.

Serengeti National Park is located in Tanzania, Africa. The park is located in a valley along a tectonic plate boundary called the East African Rift. Figure 1 shows the East African Rift and other plate boundaries near the east coast of Africa.


Figure 1. Tectonic Plate Movement near East Coast of Africa



- ~ Plate boundaries
- East African Rift
- ➔ Direction of plate movement

Serengeti National Park contains a variety of tropical habitats but is dominated by wideopen lands that include a variety of grass species with scattered trees and shrubs. Most grass growth occurs after the March through May rainy season. Many of the park's herbivores, such as wildebeest and zebra, migrate through the park in an annual cycle following the growth of new grass (Figure 2).

Figure 2. Annual Migration of Wildebeest in Serengeti National Park



2. Respond to parts A, B, C, D, E, F, G, H, I, and J.

- A.** Based on the information in Figure 1, **identify** the type of plate boundary that runs beneath Serengeti National Park.
- B.** Based on the information provided, **identify** the dominant biome within the Serengeti.
- C.** Wildebeest give birth to their young when new grasses are growing. Based on the information in Figure 2, **identify** the location in the Serengeti where wildebeest are most likely to give birth.
- D.** The Serengeti has two primary types of grasses: long and short. The two grass types provide similar levels of nutrition for animals. Wildebeest primarily consume long grasses, while other herbivores such as gazelle consume the short grasses. These two similar species coexist in Serengeti National Park.

Explain why resource partitioning allows for the coexistence of the wildebeest and the gazelle in the Serengeti.

- E.** The Tanzanian government plans to build a highway across the northern portion of the Serengeti, as shown in Figure 2.

Describe one way the planned highway shown in Figure 2 could negatively affect the wildebeest population in the Serengeti.

- F. Propose** a solution to the negative effect created by the planned highway described in part E.
- G.** Many organisms live in grasslands. One weed species, the spotted knapweed, was introduced at one location in North America in the 1890s and is currently found throughout the rangelands of the western United States. The spotted knapweed poses several environmental problems in North America. Farmers often use crop rotation to control populations of pests like the spotted knapweed. A more traditional method of controlling pests involves spraying pesticides.

Describe a characteristic of an invasive species, such as the spotted knapweed, that allows them to outcompete native species.

- H. Justify** the use of crop rotation as an environmental solution by providing an additional advantage, other than controlling pest populations.
- I. Describe** an environmental problem associated with controlling pest populations with large amounts of pesticide.
- J. Propose** a solution to controlling pests, other than pesticides or crop rotation.

3. Habitat destruction and fragmentation can have many different effects on species.

- (a) **Describe** a characteristic of a specialist species that would make the specialist species more likely to be negatively affected by habitat fragmentation than a generalist species.

A rapidly growing suburban municipality purchases nearby forested land and proposes a newly planned housing development, which would involve clear-cutting much of the area for the construction of single-family homes. While evaluating the land, the development committee discovers that development will encroach upon the habitat of a wood thrush population. The wood thrush is a solitary, territorial bird whose preferred habitat consists of large, intact densely forested areas. Wood thrush populations are also threatened by cowbirds. A cowbird lays her egg in an existing nest of a wood thrush. After the cowbird egg hatches, the cowbird pushes the unhatched wood thrush eggs out. The wood thrush parents raise the cowbird hatchling as their own.

- (b) **Identify** the symbiotic relationship between the wood thrush and the cowbird.
- (c) **Describe** one ecological advantage of leaving areas of undeveloped forest in the development plan as compared to clear-cutting the property.
- (d) **Propose** a solution that will minimize the effect of development on the resident population of wood thrush while still meeting the municipality's need for a housing development.
- (e) A male wood thrush needs a minimum of 800 m^2 of territory for reproduction. The municipal development committee has set a biodiversity preservation target of 275 male wood thrush territories. **Calculate** the area that must be set aside to support the goal of 275 male thrush territories. **Show** your work.
- (f) A real estate developer wants to build houses on the property. The plan will support 1,000 lots with a lot size of $1,100 \text{ m}^2$. The developer has proposed setting aside land equal to 10% of the size of each lot it sells. **Calculate** the maximum number of male wood thrush territories that could be created under this proposal. **Show** your work.
- (g) **Calculate** the percentage of each of the 1,000 lots that would need to be set aside in order to support the goal of 275 male wood thrush territories. **Show** your work.

Begin your response to this question at the top of a new page in the separate Free Response booklet and fill in the appropriate circle at the top of each page to indicate the question number.

STOP

END OF EXAM