

**The following information applies to parts A, B, C, D, E, and F.**

Ocelots are small wildcats that were previously found in 25 counties in Texas where shrubland is prevalent. As a result of land use changes, only 74 ocelots remain in 4 counties in Texas today. Rodents such as pocket mice are the main source of food for ocelots, but ocelots also prey on other small animals such as snakes and lizards.

**3. Respond to parts A, B, C, D, E, and F, and all subparts.****Part A**

**Identify** one environmental consequence that could result from having a smaller population of ocelots.

**Part B**

**Calculate** the percent change in the number of Texas counties where ocelots have been found historically as compared to today. **Show** your work.

**Part C**

Researchers expect the ocelot population to grow at a constant rate of 2.8% per year. **Calculate** the number of years it will take for the ocelot population to double based on a constant rate of growth. **Show** your work.

**Part D**

Pocket mice eat plants in the shrubland, while snakes are carnivores. **Explain** why the amount of primary production energy available to ocelots would change if ocelots ate only carnivores such as snakes and not pocket mice.

**Part E**

Ocelots need 720 grams of food per day. Pocket mice weigh an average of 43 grams each. If the ocelot diet consists of 46% pocket mice by weight, **calculate** how many pocket mice an ocelot would need to consume in one day, assuming each mouse weighs the average amount. **Show** your work.

**Part F**


As urbanization has increased in southern Texas, motor vehicle collisions have become the main cause of death in ocelots, which poses an added threat to the remaining ocelots.

- i. **Propose** a realistic solution to reduce the number of ocelots killed by vehicles.
- ii. **Justify** the solution proposed in part F (i) by explaining an additional advantage, other than reducing the number of ocelots killed by vehicles.

**STOP**  
**END OF EXAM**


Serengeti National Park is located in Tanzania, Africa. The park is located in a valley along a tectonic plate boundary called the East African Rift. Figure 1 shows the East African Rift and other plate boundaries near the east coast of Africa.

**Figure 1. Tectonic Plate Movement near East Coast of Africa**



Serengeti National Park contains a variety of tropical habitats but is dominated by wideopen lands that include a variety of grass species with scattered trees and shrubs. Most grass growth occurs after the March through May rainy season. Many of the park's herbivores, such as wildebeest and zebra, migrate through the park in an annual cycle following the growth of new grass (Figure 2).

**Figure 2. Annual Migration of Wildebeest in Serengeti National Park**



**2. Respond to parts A, B, C, D, E, F, G, H, I, and J.**

- A.** Based on the information in Figure 1, **identify** the type of plate boundary that runs beneath Serengeti National Park.
- B.** Based on the information provided, **identify** the dominant biome within the Serengeti.
- C.** Wildebeest give birth to their young when new grasses are growing. Based on the information in Figure 2, **identify** the location in the Serengeti where wildebeest are most likely to give birth.
- D.** The Serengeti has two primary types of grasses: long and short. The two grass types provide similar levels of nutrition for animals. Wildebeest primarily consume long grasses, while other herbivores such as gazelle consume the short grasses. These two similar species coexist in Serengeti National Park.

**Explain** why resource partitioning allows for the coexistence of the wildebeest and the gazelle in the Serengeti.

- E.** The Tanzanian government plans to build a highway across the northern portion of the Serengeti, as shown in Figure 2.

**Describe** one way the planned highway shown in Figure 2 could negatively affect the wildebeest population in the Serengeti.

- F. Propose** a solution to the negative effect created by the planned highway described in part E.
- G.** Many organisms live in grasslands. One weed species, the spotted knapweed, was introduced at one location in North America in the 1890s and is currently found throughout the rangelands of the western United States. The spotted knapweed poses several environmental problems in North America. Farmers often use crop rotation to control populations of pests like the spotted knapweed. A more traditional method of controlling pests involves spraying pesticides.

**Describe** a characteristic of an invasive species, such as the spotted knapweed, that allows them to outcompete native species.

- H. Justify** the use of crop rotation as an environmental solution by providing an additional advantage, other than controlling pest populations.
- I. Describe** an environmental problem associated with controlling pest populations with large amounts of pesticide.
- J. Propose** a solution to controlling pests, other than pesticides or crop rotation.