

ENVIRONMENTAL SCIENCE

SECTION II

Time—1 hour and 10 minutes

3 Questions

Directions: Answer all three questions, which are weighted equally; the suggested time is about 22 minutes for answering each question. Write all your answers in the Free Response booklet. Where calculations are required, clearly show how you arrived at your answer. Where explanation or discussion is required, support your answers with relevant information and/or specific examples. You may plan your answers in this orange booklet, but no credit will be given for anything written in this booklet. **You will only earn credit for what you write in the separate Free Response booklet.**

1. Researchers interested in sustainability developed three new varieties of genetically modified green beans designed to produce higher yields in arid regions. Four plots (A–D) were set up on a floodplain of a river to grow beans. Plots A, B, and C were each planted with seeds of different types of genetically modified strains of green beans. Plot D was planted with seeds of unmodified green beans. Each plot was given equal amounts of fertilizer, which contains phosphorus and nitrogen, and water and was irrigated using spray irrigation for one hour per day. Throughout the growing season, researchers measured the amount of green beans harvested within each plot. The plots with genetically modified beans had higher crop yields than the plot with unmodified beans.

- (a) **Identify** the control group in this experiment.
- (b) **Identify** the scientific question for the investigation.
- (c) Researchers repeated the experiment by modifying the length of time for the spray irrigation to 20 minutes per day. **Explain** how the results of the experiment could be altered with this modification.

Researchers also monitored the amount of sediment and fertilizer washing into the river from each plot in the original study. The soil temperature was also measured in each plot at the same depth and at the same time each day approximately 0.05 meters below the surface of the soil.

Genetically Modified Green Bean Experimental Results

Plot	Sediment Runoff (mg/L)	Phosphorus Runoff (mg/L)	Nitrogen Runoff (mg/L)	Soil Temperature (°C)
A (Type 1 GMO beans)	2.4	0.11	1.07	18.6
B (Type 2 GMO beans)	2.1	0.02	0.56	18.3
C (Type 3 GMO beans)	0.9	0.04	0.68	18.3
D (Unmodified beans)	3.5	0.15	1.24	17.6

- (d) Based on the data in the table, **identify** the plot with the lowest soil temperature.
- (e) **Describe** how sediment runoff and fertilizer runoff compare between the unmodified green beans and the genetically modified green beans.

- (f) The Type 2 GMO beans in Plot B were developed to grow more quickly than the unmodified beans in Plot D. Researchers have hypothesized that the Type 2 beans would use fertilizer more completely than the other varieties. Based on the data in the table and the experimental design, **explain** whether the researchers' hypothesis was supported or refuted.

Once the experiment was concluded, the researchers burned the plots to remove the crops that had been planted. After a few years, the researchers returned to the plots and observed a variety of plants, insects, and bird species living there.

- (g) **Describe** the ecological process that occurred on the plots after the crops were burned.

A survey indicated that one of the plots had twice the plant diversity that the other plots had. Over the next five years, the river occasionally flooded the plots, killing off many of the species that inhabited the plots.

- (h) After each flooding event, the plot with twice the plant diversity returned to its prior level of biodiversity more quickly than the other plots did. **Explain** why a community with more plant diversity will recover more quickly from the flooding.
- (i) After the last flooding event, a beetle not previously known in the area appeared in one of the plots with less plant diversity. Over a period of a few months, the new beetle population increased, whereas the existing beetle species in the plot had declining populations. **Explain** why the new beetle species could be better able to successfully populate this plot than the existing beetle species could.
- (j) **Describe** one realistic method to prevent the new beetle from spreading beyond the experimental plot.

Begin your response to this question at the top of a new page in the separate Free Response booklet and fill in the appropriate circle at the top of each page to indicate the question number.

ENVIRONMENTAL SCIENCE

SECTION II


Time—1 hour and 10 minutes

3 Questions

Directions: Answer all three questions, which are weighted equally; the suggested time is about 22 minutes for answering each question. Write all your answers in the Free Response booklet. Where calculations are required, clearly show how you arrived at your answer. Where explanation or discussion is required, support your answers with relevant information and/or specific examples. You may plan your answers in this orange booklet, but no credit will be given for anything written in this booklet. **You will only earn credit for what you write in the separate Free Response booklet.**

1. Two restoration methods designed to reclaim the overburden piles removed from open pit coal mines in temperate forests of eastern Europe were studied over several years. The first method was spontaneous restoration, where the soil and rock were dumped in noncompacted, random piles and not replanted with vegetation. The second method was technical restoration, where the land was leveled by heavy machinery, organic material was added, and eight plant species were added. Technical restoration started 5 years after the piles were formed, so no bar is included in the 1–5 year range in the graph.

Number of Plant Species in Spontaneous and Technically Restored Overburden Piles over Time



- (a) Based on the data in the graph, **identify** the average number of species found in technically restored piles 16–25 years after restoration started.
- (b) Based on the data in the graph, **describe** the relationship between the average number of species and the age of the spontaneous restoration piles.

Scientists hypothesized that after 35 years, the number of animal species in the spontaneously restored piles would be higher than the number of animal species in the technically restored piles.

- (c) Using the data in the graph, **make a claim** that supports or refutes the hypothesis that the number of animal species in the spontaneously restored piles would be higher than the number of animal species in the technically restored piles.
- (d) **Describe** a characteristic of a plant species that would have been present in the spontaneous restoration piles from years 1 to 5.
- (e) **Explain** how the return of plant life would have been altered in the spontaneous piles if the mining had taken place in the tundra rather than in temperate forests.

Limiting runoff on disturbed land improves soil and water quality. To model water runoff, students set up three large plastic trays and filled each tray with loam soil to 1 centimeter below the edge. Loam soil is a balance of sand, silt, and clay soil. One tray was seeded with wheatgrass and grown for 30 days, one tray was covered with small wood chips, and one tray was soil only. Five hundred milliliters of water was sprinkled evenly onto the soil surface of each tray. All three trays were placed on a slight slope, and water that ran off the surface was collected. The results are shown in the following table.

Tray	Water Collected (mL)	Water Appearance
Wheatgrass grown in soil; no small wood chips	190	Clear
Small wood chips covered soil; no wheatgrass	300	Transparent; light brown
Soil only; no wheatgrass or small wood chips	360	Cloudy; dark brown


- (f) **Identify** the scientific question for the investigation.
- (g) **Identify** a dependent variable in the experiment.
- (h) **Explain** how the results of the experiment would likely have been altered if the soil had contained a higher percentage of sand, which has the largest particle size of the components of loam.

Runoff from mining waste can also affect water quality.

- (i) **Describe** one ecological impact that could result from the drainage from mine waste into nearby waterways.
- (j) **Describe** one strategy to reduce the drainage from mine waste into waterways downstream from mines.

Begin your response to this question at the top of a new page in the separate Free Response booklet and fill in the appropriate circle at the top of each page to indicate the question number.

2. Biologists collected data on the abundance of lizards on several Pacific islands with and without mongoose populations. The surveys were conducted at different locations on the islands by counting the number of lizards observed per hour and included habitats with both undisturbed areas and anthropogenically disturbed areas.



- (a) The graph shows data collected by biologists on lizard numbers on several Pacific islands.
- Based on the data in the graph, **identify** the habitat type with the greatest number of lizards.
 - Based on the data in the graph, **describe** the relationship between the lizard numbers observed on islands with and without mongoose populations.

Globally, islands vary in size, location, and habitat.

- (b) Based on the theory of island biogeography, **describe** the characteristics of an island with the greatest species diversity.
- (c) Island ecosystems are often threatened by invasive species, such as the mongoose populations studied in the survey.
- Describe** one environmental problem caused by invasive animal species.
 - Invasive species are often pests. **Make a claim** that proposes one way to control pest species.
 - Justify** the solution proposed in (c)(ii) by providing an additional benefit of the solution.
- (d) Islands provide humans with many ecosystem services. However, humans may alter ecosystems when they obtain these services. Large-scale commercial sugarcane farming on the Pacific islands has led to significant ecological damage.
- Identify** a provisioning ecosystem service provided by primary forests.
 - Based on the data in the graph, **explain** how replacing primary forests with sugarcane fields may have affected the number of lizards counted on the mongoose-free Pacific islands.
 - The long-term use of monocultures in commercial sugarcane farming is one cause of the ecological damage. **Describe** one problem associated with monocultures.

One solution to the ecological damage of monocultures is to use a form of crop rotation using legumes.

- (e) **Describe** an advantage of crop rotation using legumes on soil fertility.

3. North Carolina is one of the fastest-growing states by population in the United States, with most growth concentrated in major cities like Charlotte. Increased population leads to increased demands for resources such as power and water.

The majority of electrical energy in North Carolina is produced during the combustion of coal.

- (a) **Identify** one negative human health effect linked to the exposure to pollutants resulting from the combustion of coal.
- (b) Coal ash, the powdery, solid substance that remains after coal combustion, presents a variety of potential environmental issues. Disposal of toxic coal ash is a growing concern in the state. Coal ash can be mixed with water and disposed of in large, unlined pits or as dry ash in landfills.
- (i) **Describe** an environmental problem associated with coal ash waste disposal.
- (ii) A proposed solution is to dispose of North Carolina's coal ash in clay-lined pits. **Justify** this solution by providing one advantage of using clay soil.
- (c) The population of Charlotte, North Carolina, has grown over the last few decades. The table shows the increase in the population size of Charlotte from 2013 to 2019.

Population of Charlotte, North Carolina, 2013–2019

Year	Population
2013	757,278
2014	774,807
2015	792,137
2016	808,834
2017	826,060
2018	841,611
2019	857,425

- (i) **Calculate** the percent change in Charlotte's population from 2013 to 2019. **Show** your work.
- (ii) Based on Charlotte's 2019 growth rate of 1.88%, **calculate** the year when the population of Charlotte will double, assuming the growth rate stays the same. **Show** your work.
- (iii) The average Charlotte resident uses 90 gallons of water per day. **Calculate** the gallons of water used by the population of Charlotte in the year 2018. **Show** your work.
- (d) Drought periods are becoming more frequent in North Carolina, causing water resources to become more scarce. **Describe** one realistic action that citizens could take to reduce domestic outdoor water use.

Begin your response to this question at the top of a new page in the separate Free Response booklet and fill in the appropriate circle at the top of each page to indicate the question number.

STOP

END OF EXAM

4. Soil is a complex mixture of living organisms and organic material, along with minerals and other abiotic components. Soils help sustain life and support ecosystem functions.

- (a) **Describe** how TWO climate factors affect the rate of soil formation.
- (b) As soils form, distinct layers known as horizons develop over time. One of these is the A horizon.
 - (i) **Identify** one specific biotic component of the A horizon.
 - (ii) **Identify** one abiotic component of the A horizon.

Resources such as soil and water can be degraded by human activities.

- (c) Nitrate levels exceeding the United States Environmental Protection Agency's primary drinking water standard have been found in the groundwater of areas with intensive agriculture.
 - (i) **Identify** one agricultural practice that can lead to elevated nitrate levels in groundwater.
 - (ii) **Describe** how the practice you identified in (c)(i) leads to elevated nitrate levels in groundwater.
- (d) Acid deposition has affected soil quality in many parts of the northeastern United States.
 - (i) **Explain** one way acid deposition onto soil can affect plant health.
 - (ii) **Describe** one method for remediating soil affected by acid deposition.
- (e) Climate change is causing far-reaching ecosystem changes, including soil degradation in many of the world's biomes. **Describe** TWO ways that climate change can degrade soil.

STOP

END OF EXAM