

There is a natural climate phenomenon that occurs in the equatorial Pacific Ocean because of periodic fluctuations in sea surface conditions and atmospheric circulation. This phenomenon happens every few years, resulting in shifts in temperature and precipitation in many parts of the world.

Figure 1 shows temperature and precipitation patterns associated with one phase of this phenomenon in the equatorial Pacific Ocean. Figure 2 shows the associated temperature and precipitation patterns that occur over North America.

Figure 1. Effects of Changes in Sea Surface Conditions in Equatorial Pacific Ocean





Figure 2. Polar Jet Stream and Climate Conditions in North America



2. Respond to parts A, B, C, D, E, F, G, H, I, and J.

- A. **Identify** the sea surface condition for the eastern equatorial area of the Pacific Ocean illustrated in Figure 1.
- B. Based on the information in Figure 1, **identify** the climate phenomenon associated with the sea surface conditions shown in the equatorial area of the Pacific Ocean.
- C. Based on the information in Figure 2, **describe** a difference in climate patterns between the regions A and B.
- D. Increases in regional precipitation can lead to increased flooding in certain locations. **Describe** one reason why there is an increased risk of flooding in urban areas during extended periods of precipitation.
- E. **Propose** a realistic solution a city could implement to decrease the risk of flooding in urban areas.
- F. **Justify** the solution proposed in part E by providing an additional advantage other than a reduction in the risk of flooding.
- G. Terrestrial biomes are primarily determined by temperature and precipitation patterns. **Describe** one difference between the climate of a temperate seasonal forest and that of a savanna.
- H. Droughts are one environmental problem that can affect the frequency and severity of forest fires. **Identify** the ecological process that occurs following a forest fire that leaves the soil intact.
- I. **Describe** one way burning forests contribute to atmospheric pollution.
- J. **Describe** one sustainable forestry practice that could be used to reduce the occurrence or severity of forest fires.

3. The Rocky Mountains, the headwaters for the Colorado River, have experienced a severe drought for many of the past twenty years. States and cities downstream that depend on the river water have had to find solutions to the reduced water flow.

One city located downstream from the Rocky Mountains is Las Vegas, Nevada. In Las Vegas, one solution to reduce the amount of water used for irrigation is to remove landscaping and lawns that use grasses not native to the local area and replace them with desert-tolerant landscaping.

- (a) **Identify** a typical type of plant that would be used in desert-tolerant landscaping.
- (b) **Justify** the removal of landscaping and lawns that use grasses not native to the local area by describing an additional advantage, other than reducing the amount of water needed for irrigation.

Warmer seasonal temperatures in the Rocky Mountains have led to earlier mountain snowmelt and an increased risk of forest fires as soil and vegetation dry out.

- (c) **Describe** how a prescribed burn would reduce the severity and spread of forest fires.
- (d) **Describe** a disadvantage of a prescribed burn in a forested ecosystem.

Lake Powell is a large reservoir on the Colorado River. Lake Powell is formed by a dam with a hydroelectric power plant. At full capacity or full pool, Lake Powell contains 25.16 million acre-feet (maf) of water. Currently, Lake Powell is at 36% full capacity.

- (e) **Calculate** the number of million acre-feet (maf) of water currently in Lake Powell. **Show** your work.

In 2021, after years of drought, Lake Powell held 7.9 million acre-feet (maf) of water. One million acre-feet is equivalent to 3.26×10^{11} gallons.

- (f) The watershed of the Upper Colorado River contributes an average of 9.60 maf of water to Lake Powell annually. The melted mountain snow found in the watershed contributes 50% of the average river flow into Lake Powell. In 2021, the river flow was 36% of the annual average. Assuming that all resources to that flow contributed the same proportions as in prior years, **calculate** the amount of water (in million acre-feet) that was contributed by the melted mountain snow in 2021. **Show** your work.
- (g) The average household in the United States consumes 5.0×10^4 gallons of water in a year. **Calculate** the number of households that could be supported for one year by the average flow of 9.60 maf of water into Lake Powell. **Show** your work.

Begin your response to this question at the top of a new page in the separate Free Response booklet and fill in the appropriate circle at the top of each page to indicate the question number.

STOP

END OF EXAM

2. As conventional sources of crude oil are depleted, unconventional sources such as oil sands (also known as tar sands) are being utilized. Oils sands contain bitumen, which can be processed into a synthetic crude oil. A region of boreal forest in Alberta, Canada, that covers a deposit of oil sands will be cut and removed during the process of bitumen extraction. It is estimated that the deposit contains 73 billion barrels of recoverable bitumen. The rate of extraction from the deposit will be approximately 1 million barrels of bitumen per day.
- (a) **Identify** one ecological benefit, other than providing habitat, that is provided by forests.
 - (b) **Identify** one economic benefit that is provided by forests.
 - (c) **Describe** TWO environmental consequences, other than those related to the loss of boreal forest habitat, that result from the extraction of bitumen or the transportation of synthetic oil to customers.
 - (d) Assuming the above extraction rate, **calculate** how many days will be needed to extract the recoverable volume of bitumen from the oil sands.
 - (e) **Calculate** how many years will be needed to fully extract the recoverable volume of bitumen from the oil sands.
 - (f) Monthly production of synthetic crude oil is 30 million barrels. Producing one barrel of synthetic crude oil uses two barrels of heated freshwater. **Calculate** the number of barrels of freshwater needed each year to supply this demand.

3. Haiti shares a border with the Dominican Republic on the Caribbean island of Hispaniola.
- (a) The border between the two countries can be seen using satellite images because of the severe deforestation in Haiti.
 - (i) **Provide** one reason why deforestation commonly occurs in a less developed country such as Haiti.
 - (ii) **Describe** one realistic strategy to reduce deforestation in a less developed country.
 - (b) Deforestation can affect water quality. **Identify** one change that can occur in the water quality of streams within a watershed that has been deforested. **Explain** how deforestation can lead to this change.
 - (c) **Identify** TWO environmental benefits, other than those related to water quality, of maintaining forest ecosystems.

The table below contains demographic data for Haiti in 1995 and 2015.

DEMOGRAPHIC DATA FOR HAITI

Measure	1995	2015
Fertility rate (number of children per woman)	5.2	2.7
Life expectancy (years)	55	64
Infant mortality (deaths per 1,000)	85	48

- (d) **Identify** and **discuss** one factor in a less developed country that could contribute significantly to a change in life expectancy, similar to what occurred in Haiti from 1995 to 2015.
- (e) **Identify** and **discuss** one economic or cultural factor in a less developed country that could contribute significantly to a change in the fertility rate, similar to what occurred in Haiti from 1995 to 2015.