



2. As the human population continues to grow, the demand for animal protein is expected to double by 2050. Animal protein can be obtained from conventional livestock and other sources. The production of all animal protein has environmental costs, as shown in the following graphs. In Figure 1, global warming potential (GWP) is measured in kilograms of carbon dioxide equivalent ($\text{kg CO}_2\text{-eq.}$). Carbon dioxide equivalent is a measure of the warming effect of a gas relative to that of carbon dioxide. In Figure 2, land use is measured in meters squared (m^2).

Global Warming Potential and Land Use from Animal Protein Production



- (a) Based on the data in the graphs, **identify** the amount of land required to produce 1 kilogram of chicken protein.

Mealworms, the larvae of the darkling beetle, consume grains and decaying plant material. Darkling beetles are an *r*-selected species, and the entire mealworm can be consumed by humans for protein.

- (b) Based on the information provided, **identify** the type of survivorship curve exhibited by the darkling beetle.

- (c) **Explain** why the reproductive strategy of the darkling beetle is an advantage for using mealworms as an alternative protein source for the rapidly growing human population.

An individual's diet choices affect greenhouse gas emissions and land use.

- (d) Based on the data in the graphs, **explain** whether producing 1 kilogram of chicken protein or 1 kilogram of pork protein would cause less environmental damage.

Cattle emit methane from fermentation in their digestive systems. The other animals do not emit methane.

- (e) Based on the data in the graphs, **explain** why the production of 1 kilogram of beef protein has a different impact on global warming than the production of 1 kilogram of protein from any of the other animals studied would have.

Agricultural practices can have negative impacts on rivers and streams.

- (f) **Describe** how water quality can be altered by cattle grazing that occurs near a stream or river.
- (g) **Propose** a solution to reduce the negative impacts on waterways that result from cattle grazing, while still allowing cattle to graze.
- (h) Crop production can cause soil erosion. **Describe** a sustainable agricultural practice used to reduce soil erosion.
- (i) **Justify** the use of the sustainable practice described in part (h) by describing an additional advantage, other than the reduction of soil erosion.
- (j) Crop production around the world is affected by climate change. **Describe** how crop production could be negatively affected by climate change.

Begin your response to this question at the top of a new page in the separate Free Response booklet and fill in the appropriate circle at the top of each page to indicate the question number.

ENVIRONMENTAL SCIENCE


SECTION II

Time—1 hour and 10 minutes

3 Questions


Directions: Answer all three questions, which are weighted equally; the suggested time is about 22 minutes for answering each question. Write all your answers in the Free Response booklet. Where calculations are required, clearly show how you arrived at your answer. Where explanation or discussion is required, support your answers with relevant information and/or specific examples. You may plan your answers in this orange booklet, but no credit will be given for anything written in this booklet. **You will only earn credit for what you write in the separate Free Response booklet.**

1. The following diagram shows how an area changed over time. The dominant vegetation in each stage is labeled in the diagram. Each roman numeral represents a stage in the ecological process.



- (a) **Identify** the ecological change over time illustrated in the diagram.
- (b) **Identify** the stage in which pioneer species first appear.
- (c) **Describe** how soil forms from stage II to stage III.
- (d) **Identify** a cultural ecosystem service that may be provided by a mature forest.

Researchers are interested in investigating the effect of the removal of trees on the water quality of nearby waterways. To conduct their experiment, the researchers locate two identical forest watersheds, both with a river running through them. Forest A is an intact forest and forest B is a clear-cut forest. Four stream monitoring stations are set up in each forest study area. Each station monitors a variety of water quality indicators, including water temperature, at various locations along the river. The experimental setup is shown in the following diagrams. The arrows shown in the rivers represent the direction of water flow. The triangles in the river show the location of the stream monitoring stations.



- (e) **Identify** a testable hypothesis being investigated by the researchers.
- (f) **Identify** the independent variable in the experiment.
- (g) **Describe** the purpose of forest A in the experiment.
- (h) **Explain** why the water temperature might change because of clear-cutting trees in forest B.


After the experiment was completed, land developers constructed a golf course in the clear-cut area of forest B. To prepare the land, developers removed any remaining stumps and newly grown trees and began planting and fertilizing new grass. The researchers are interested in the effect the golf course may have on the water quality of the river. To monitor the river in forest B, the researchers repeated the experiment once the golf course had been constructed.

- (i) **Explain** why an indicator of water quality other than temperature could be altered by the golf course in forest B.
- (j) **Explain** how the aquatic organisms living in the lake illustrated in the diagram of forest B are likely to be affected by the new golf course.

Begin your response to this question at the top of a new page in the separate Free Response booklet and fill in the appropriate circle at the top of each page to indicate the question number.

2. Asian carp were first introduced in the southern United States in the 1970s by fish farmers wanting to control algal growth in fish ponds. Asian carp can eat up to 40% of their body weight in algae each day, and a female carp can produce up to 1 million eggs every year.

Distribution of the Asian Carp in the United States



- (a) Based on the information given, **identify** a body of water invaded by Asian carp that is labeled on the map.
- (b) Based on the information shown in the map, **describe** the change in the distribution of Asian carp since their introduction in the 1970s.

The Great Lakes are a group of five major freshwater inland lakes that all share the same watershed.

- (c) **Describe** one impact the introduction of Asian carp could have on the ecosystem services provided by the Great Lakes region.
- (d) **Propose** a realistic solution to help reduce the spread of the Asian carp from their current distribution.
- (e) **Justify** the solution proposed in part (d) by providing an additional advantage.

The following is an example of a simple food chain in the Great Lakes.

Algae → Zooplankton → Macroinvertebrates → Yellow perch → Rainbow trout

- (f) **Identify** the primary consumer in the Great Lakes food chain.
- (g) **Describe** what the arrows in the aquatic food chain represent.

(h) **Describe** one possible effect of the introduction of the Asian carp on the Great Lakes food chain.

The blue pike, a fish that was endemic to the Great Lakes, was commercially fished from the late 1800s until their population crashed in 1958. In 1970, the species was declared to be extinct.

(i) **Describe** one way that overfishing of blue pike illustrates the tragedy of the commons.

(j) One potential solution to reduce overfishing is to use aquaculture. **Describe** one disadvantage of this solution.

Begin your response to this question at the top of a new page in the separate Free Response booklet and fill in the appropriate circle at the top of each page to indicate the question number.