

1.3 Representing Motion

Name: _____ Class: _____ Date: _____

Total: 14 marks

KEY WORDS slope 斜率 • area 面积 • free fall 自由落体 • kinematic equations 运动学方程
• velocity 速度

Objective

Build the skills to answer exam questions on **representing motion** — motion graphs and the equations for constant acceleration.

You must be able to:

- read **position-time**, **velocity-time**, and **acceleration-time** graphs
- use the **slope** 斜率: slope of an x - t graph = velocity; slope of a v - t graph = acceleration
- use the **area** 面积: area under a v - t graph = displacement; area under an a - t graph = change in velocity
- apply the **kinematic equations** 运动学方程 for constant acceleration
- analyse **free fall** 自由落体 with $g = 9.8 \text{ m s}^{-2}$

1 Worked examples

Study these first. Each one shows the method for a question type used later.

■ The kinematic equations (constant a)

$$v = v_0 + at, \quad \Delta x = v_0 t + \frac{1}{2}at^2,$$

$$v^2 = v_0^2 + 2a \Delta x, \quad \Delta x = \frac{1}{2}(v_0 + v)t.$$

Choose the one that is **missing the quantity you don't know**.

■ Slopes and areas

On a v - t graph the **slope** is the acceleration and the **area** under the line is the displacement. A line rising from 0 to 12 m s^{-1} over 4.0 s: slope = 3.0 m s^{-2} ; area = $\frac{1}{2}(4.0)(12) = 24 \text{ m}$.

■ Free fall

Take down as positive, so $a = g = 9.8 \text{ m s}^{-2}$. A ball dropped from rest for 2.0 s: $\Delta x = \frac{1}{2}(9.8)(2.0)^2 = 19.6 \text{ m}$ and $v = 9.8 \times 2.0 = 19.6 \text{ m s}^{-1}$.

2 Practice

2.1 A car accelerates from rest at 2.0 m s^{-2} for 6.0 s . Find its final velocity. [1]

2.2 For the same car, find the distance it travels in that time. [2]

2.3 On a velocity-time graph, state what (a) the slope and (b) the area under the line represent. [2]

2.4 A stone is dropped from rest. Find its speed after 3.0 s (take $g = 9.8 \text{ m s}^{-2}$). [1]

3 Exam-style questions

3.1 The area under a velocity-time graph gives the [1]

A	B
C	D

- A acceleration
- B displacement
- C speed
- D jerk

3.2 A ball is thrown straight up. At its highest point, its [1]

A	B
C	D

- A velocity and acceleration are both zero
- B velocity is zero and acceleration is 9.8 m s^{-2} downward
- C velocity is maximum and acceleration is zero
- D velocity and acceleration both point up

3.3 A car travelling at 25 m s^{-1} brakes to rest in a distance of 50 m.

(a) Find its acceleration. [2]

(b) Find the time it takes to stop. [1]

3.4 A ball is thrown vertically upward at 20 m s^{-1} (take $g = 10 \text{ m s}^{-2}$).

(a) Find the time to reach the top. [1]

(b) Find the maximum height reached. [2]