

2.5 Newton's Second Law

Name: _____ Class: _____ Date: _____

Total: 11 marks

KEY WORDS incline 斜面 • newton's second law 牛顿第二定律 • components 分量
• friction 摩擦力 • force 力

Objective

Build the skills to answer exam questions on **Newton's second law** — relating force, mass, and acceleration.

You must be able to:

- state **Newton's second law** 牛顿第二定律, $\vec{a} = \frac{\sum \vec{F}}{m}$
- explain that acceleration is **proportional** to net force and **inversely proportional** to mass
- resolve forces into perpendicular **components** 分量 on an inclined surface
- apply the second law to connected objects joined by a string

1 Worked examples

Study these first. Each one shows the method for a question type used later.

■ Applying $F = ma$

A net force of 12 N on a 3.0 kg block gives $a = \frac{12}{3.0} = 4.0 \text{ m s}^{-2}$ in the direction of the net force.

■ On an incline

Resolve weight along and perpendicular to the slope. On a frictionless incline of angle θ , the acceleration down the slope is $a = g \sin \theta$.

■ Connected masses

For masses joined by a string, apply $F = ma$ to the **whole system** to find a , then to **one mass** to find the tension. A 2.0 kg and a 3.0 kg block pulled by 10 N:
 $a = \frac{10}{5.0} = 2.0 \text{ m s}^{-2}$.

2 Practice

2.1 A net force of 20 N acts on a 4.0 kg trolley. Find its acceleration. [1]

2.2 A 1500 kg car accelerates at 2.0 m s^{-2} . Find the net force on it. [1]

2.3 A block slides down a frictionless ramp at 30° (take $g = 10 \text{ m s}^{-2}$). Find its acceleration along the ramp. [2]

2.4 State how the acceleration changes if the net force doubles while the mass stays the same. [1]

3 Exam-style questions

3.1 If the net force on an object doubles and its mass halves, the acceleration becomes [1]

A	B
C	D

- A** unchanged
- B** twice as large
- C** four times as large
- D** half as large

3.2 A 2.0 kg object accelerates at 5.0 m s^{-2} . The net force on it is [1]

A	B
C	D

- A** 0.4 N
- B** 2.5 N
- C** 7.0 N
- D** 10 N

3.3 Two blocks, 2.0 kg and 3.0 kg, sit in contact on a frictionless floor. A 15 N force pushes on the 2.0 kg block.

(a) Find the acceleration of the pair. [2]

(b) Find the contact force between the blocks. [2]