

3.4 Conservation of Energy

Name: _____ Class: _____ Date: _____

Total: 10 marks

KEY WORDS thermal energy 热能 • conservative forces 保守力 • conserved 守恒
• nonconservative forces 非保守力 • energy 能量

Objective

Build the skills to answer exam questions on **conservation of energy**.

You must be able to:

- state the **law of conservation of energy** 能量守恒定律 for an isolated system
- distinguish **conservative forces** 保守力 from **nonconservative forces** 非保守力 such as friction
- apply **conservation of mechanical energy** when no nonconservative force does work
- account for energy transferred to **thermal energy** 热能 when friction acts

1 Worked examples

Study these first. Each one shows the method for a question type used later.

■ Mechanical energy is conserved without friction

With only conservative forces, $KE_i + PE_i = KE_f + PE_f$. A pendulum trades potential for kinetic energy and back.

■ With friction, track the heat

$KE_i + PE_i = KE_f + PE_f + E_{thermal}$. Friction converts mechanical energy into thermal energy — the total is still conserved.

■ Example

A 1.0 kg ball is dropped 5.0 m (take $g = 10 \text{ m s}^{-2}$). All the PE becomes KE:
 $v = \sqrt{2gh} = \sqrt{2(10)(5.0)} = 10 \text{ m s}^{-1}$.

2 Practice

2.1 A 2.0 kg cart starts from rest at the top of a 3.0 m high frictionless ramp (take $g = 10 \text{ m s}^{-2}$). Find its speed at the bottom. [2]

2.2 State the main energy change when friction acts on a moving block. [1]

2.3 A pendulum bob swings down through a height of 0.20 m. Find its speed at the lowest point (take $g = 10 \text{ m s}^{-2}$). [2]

3 Exam-style questions

3.1 As a skier descends a frictionless slope, her gravitational potential energy is [1]

A	B
C	D

A increased

B converted to kinetic energy

C unchanged

D converted to thermal energy

3.2 When friction acts on a moving object, its mechanical energy is [1]

A	B
C	D

A created

B exactly conserved

C partly converted to thermal energy

D destroyed

3.3 A 0.50 kg ball is dropped from 8.0 m and hits the ground at 10 m s^{-1} (take $g = 10 \text{ m s}^{-2}$).

(a) Find the initial gravitational potential energy. [1]

(b) Find the kinetic energy on landing. [1]

(c) Find the energy converted to thermal and other forms. [1]