


Begin your response to **QUESTION 4** on this page.

4. (7 points, suggested time 13 minutes)

A simple pendulum consists of a small sphere that hangs from a string with negligible mass. The top end of the string is fixed. The sphere is pulled to Point A so that the string makes a small angle θ with the vertical, as shown. The sphere is then released from rest and swings through its lowest point at Point B. The work done on the sphere by Earth between points A and B is W_E .

The pendulum is then taken to Planet X. The mass of Planet X is the same as the mass of Earth, but the radius of Planet X is greater than the radius of Earth. The sphere is again brought to Point A (displaced θ from the vertical), released from rest, and swings through its lowest point at Point B. The work done on the sphere by Planet X between points A and B is W_X .

(a) **Justify** why W_X is less than W_E .

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 4** on this page.

A new pendulum is made by hanging the same small sphere from a different string with negligible mass. The new string is slightly elastic, and the length of the string may increase or decrease depending on the tension applied to the string. On Earth, when the sphere is again displaced θ from the vertical and released from rest, the new pendulum oscillates with period T_E .

The new pendulum is then taken to a different planet, Planet Y. The radius of Planet Y is the same as the radius of Earth, but the mass of Planet Y is larger than the mass of Earth. On Planet Y, when the sphere is again displaced from the vertical and released from rest, the new pendulum oscillates with period T_Y .

- (b) In a clear, coherent paragraph-length response that may also contain drawings, **explain** how T_Y could be larger than T_E but also could be smaller than T_E .

GO ON TO THE NEXT PAGE.