


Question 2: Version J

2. Two small disks, R and S, are on a straight, horizontal track.

At time $t = 0$, Disk R of mass m_0 is located at position $x = 0$. Disk R is initially moving with speed v_0 in the $+x$ -direction. Disk S of mass $3m_0$ is initially at rest, as shown in the top view in Figure 1. Frictional forces are negligible.

At $t = t_1$, the disks collide. Immediately after the collision, Disk R moves with speed $\frac{1}{2}v_0$ in the $-x$ -direction.

Figure 1




Top View

Note: Figure not drawn to scale.

Part A

The momentum-vector diagram in Figure 2 represents the momentums of Disk R and Disk S before the collision.


Figure 2



Draw arrows on Figure 3 to represent the momentum vectors of Disk R and Disk S after the collision.

- Each arrow must start on, and point away from, each dot.
- If the momentum of either disk is zero, write “ $p = 0$ ” next to the dot.
- Draw the length of each arrow to represent the magnitude of each momentum, consistent with the scale used in Figure 2.

Figure 3



Part B


Starting with conservation of linear momentum, **derive** an expression for the kinetic energy of Disk S immediately after the collision. Express your final answer in terms of m_0 , v_0 , and physical constants, as appropriate. Begin your derivation by writing the fundamental physics principle or an equation from the reference information.

Part C

The graph shown in Figure 4 represents the positions x as functions of time t from $t = 0$ until $t = t_1$ for each of the following:

- Disk R
- Disk S
- The center of mass of the two-disk system

On the graph shown in Figure 4, **draw** three lines that represent the positions x of Disk R, Disk S, and the center of mass of the two-disk system as functions of t from $t = t_1$ to $t = 2t_1$. Distinctly **label** each line.

Figure 4**Part D**

During the collision, the magnitudes of the changes in momentum of Disk R and Disk S are $|\Delta p_R|$ and $|\Delta p_S|$, respectively.

Indicate whether $|\Delta p_R|$ is greater than, less than, or equal to $|\Delta p_S|$ by writing one of the following.

- $|\Delta p_R| > |\Delta p_S|$
- $|\Delta p_R| < |\Delta p_S|$
- $|\Delta p_R| = |\Delta p_S|$


Briefly **justify** your response by referencing a fundamental physics principle.

Question 1: Version J

1. A student has a cart of mass m_c and a block of mass $\frac{1}{5}m_c$, as shown in Figure 1.

- At time $t = 0$, the cart is moving to the right across a horizontal surface with constant speed v_c , and the student releases the block from rest.
- At $t = t_1$, the block collides with and sticks to the top of the cart. The block does not slide on the cart.
- At $t = t_2$, the block-cart system continues to move to the right with constant speed v_f .

Figure 1



A.

- i. On the axes shown in Figure 2, **sketch** a graph of the magnitude p_x of the x -component of the momentum of the block-cart system as a function of time t from $t = 0$ until $t > t_2$.




Figure 2

- ii. **Derive** an expression for the speed v_f of the block-cart system after time $t = t_2$ in terms of m_c , v_c , and physical constants, as appropriate. Begin your derivation by writing a fundamental physics principle or an equation from the reference information.
- iii. **Derive** an expression for the change in the kinetic energy ΔK in the block-cart system from $t = 0$ to $t = t_2$ in terms of m_c , v_c , and physical constants, as appropriate. Begin your derivation by writing a fundamental physics principle or an equation from the reference information.
- B.** Consider the case where a new block is dropped and collides with the top of the cart. The new block slides along the cart during the collision but does not slide off the cart. The time interval from when the new block collides with the cart and moves together with the cart is Δt . During Δt there is a frictional force between the new block and the cart.
- Indicate** whether the x -component of the momentum of the new block-cart system increases, decreases, or remains constant during Δt .

_____ Increases

_____ Decreases

_____ Remains constant

Justify your response.

Begin your response to **QUESTION 5** on this page.




Figure 1

5. (7 points, suggested time 13 minutes)

At time $t = 0$, Block A slides along a horizontal surface toward Block B, which is initially at rest, as shown in Figure 1. The masses of blocks A and B are 6 kg and 2 kg, respectively. The blocks collide elastically at $t = 1.0$ s, and as a result, the magnitude of the change in kinetic energy of Block B is 9 J. All frictional forces are negligible.

(a) **Determine** the speed of Block B immediately after the collision.

GO ON TO THE NEXT PAGE.


Continue your response to **QUESTION 5** on this page.

Figure 2

The graph shown in Figure 2 represents the positions x of Block A, Block B, and the center of mass of the two-block system as functions of t between $t = 0$ and $t = 1.0$ s.

- (b) On the graph in Figure 2, **draw** and **label** three lines to represent the positions of Block A, Block B, and the center of mass of the two-block system as functions of t between $t = 1.0$ s and $t = 2.0$ s. Each line should be distinctly labeled.
- (c) Consider if in the original scenario, instead of colliding elastically, the blocks collided and stuck together. **Describe** how the line drawn for the center of mass in part (b) would change, if at all. Briefly **justify** your response.

GO ON TO THE NEXT PAGE.


STOP

END OF EXAM


(7 points, suggested time 13 minutes)

A student has a piece of clay and a rubber sphere, both of the same mass. Both objects are thrown horizontally at the same speed at identical blocks that are at rest at the edge of identical tables, as shown, where friction between the blocks and the table is negligible. After the collisions, both blocks fall to the floor.


In Case A, the clay sticks to Block A after the collision. In Case B, the rubber sphere bounces off of Block B after the collision.




Case A: Clay and Block A
Before Collision



Case B: Rubber Sphere and Block B
Before Collision



Case A: Momentum of
Clay-Block System
Immediately After Collision



Case B: Momentum of
Sphere-Block System
Immediately After Collision

- (a) In the figure at left above, the arrow represents the momentum immediately after the collision for the clay-block system in Case A. In the figure at right above, draw an arrow starting on the dot to represent the momentum of the sphere-block system immediately after the collision in Case B. If the momentum is zero, write “zero” next to the dot. The momentum, if it is not zero, must be represented by an arrow starting on, and pointing away from, the dot. The length of the vector, if not zero, should reflect the magnitude of the momentum relative to Case A.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 4** on this page.

- (b) After the clay and Block A collide, Block A lands a horizontal distance d_A from the edge of the table. Does Block B land on the floor at a horizontal distance from the edge of the table that is greater than, less than, or equal to d_A ? In a clear, coherent, paragraph-length response that may also contain equations and/or drawings, explain your reasoning. Neglect any frictional effects due to the table or air resistance.

GO ON TO THE NEXT PAGE.


3. (12 points, suggested time 25 minutes)

(a) A student of mass M_S , standing on a smooth surface, uses a stick to push a disk of mass M_D . The student exerts a constant horizontal force of magnitude F_H over the time interval from time $t = 0$ to $t = t_f$ while pushing the disk. Assume there is negligible friction between the disk and the surface.


i. Assuming the disk begins at rest, determine an expression for the final speed v_D of the disk relative to the surface. Express your answer in terms of F_H , t_f , M_S , M_D , and physical constants, as appropriate.

ii. Assume there is negligible friction between the student's shoes and the surface. After time t_f , the student slides with speed v_S . Derive an equation for the ratio v_D / v_S . Express your answer in terms of M_S , M_D , and physical constants, as appropriate.

(b) Assume that the student's mass is greater than that of the disk ($M_S > M_D$). On the grid below, sketch graphs of the speeds of both the student and the disk as functions of time t between $t = 0$ and $t = 2t_f$. Assume that neither the disk nor the student collides with anything after $t = t_f$. On the vertical axis, label v_D and v_S . Label the graphs "S" and "D" for the student and the disk, respectively.

**GO ON TO THE NEXT PAGE.**

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.



Disk

Block

(c) The disk is now moving at a constant speed v_1 on the surface toward a block of mass M_B , which is at rest on the surface, as shown above. The disk and block collide head-on and stick together, and the center of mass of the disk-block system moves with speed v_{cm} .


- i. Suppose the mass of the disk is much greater than the mass of the block. Estimate the velocity of the center of mass of the disk-block system. Explain how you arrived at your prediction without deriving it mathematically.
- ii. Suppose the mass of the disk is much less than the mass of the block. Estimate the velocity of the center of mass of the disk-block system. Explain how you arrived at your prediction without deriving it mathematically.
- iii. Now suppose that neither object's mass is much greater than the other but that they are not necessarily equal. Derive an equation for v_{cm} . Express your answer in terms of v_1 , M_D , M_B , and physical constants, as appropriate.
- iv. Consider the scenario from part (c)(i), where the mass of the disk was much greater than the mass of the block. Does your equation for v_{cm} from part (c)(iii) agree with your reasoning from part (c)(i) ?
____ Yes ____ No
Explain your reasoning by addressing why, according to your equation, v_{cm} becomes (or approaches) a certain value when M_D is much greater than M_B .

GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

PHYSICS 1**Section II****Time—1 hour and 30 minutes****5 Questions**


Directions: Questions 1, 4, and 5 are short free-response questions that require about 13 minutes each to answer and are worth 7 points each. Questions 2 and 3 are long free-response questions that require about 25 minutes each to answer and are worth 12 points each. Show your work for each part in the space provided after that part.




1. (7 points, suggested time 13 minutes)

Identical blocks 1 and 2 are placed on a horizontal surface at points A and E, respectively, as shown. The surface is frictionless except for the region between points C and D, where the surface is rough. Beginning at time t_A , block 1 is pushed with a constant horizontal force from point A to point B by a mechanical plunger. Upon reaching point B, block 1 loses contact with the plunger and continues moving to the right along the horizontal surface toward block 2. Block 1 collides with and sticks to block 2 at point E, after which the two-block system continues moving across the surface, eventually passing point F.

- (a) On the axes below, sketch the speed of the center of mass of the two-block system as a function of time, from time t_A until the blocks pass point F at time t_F . The times at which block 1 reaches points A through F are indicated on the time axis.





- (b) The plunger is returned to its original position, and both blocks are removed. A uniform solid sphere is placed at point A, as shown. The sphere is pushed by the plunger from point A to point B with a constant horizontal force that is directed toward the sphere's center of mass. The sphere loses contact with the plunger at point B and continues moving across the horizontal surface toward point E. In which interval(s), if any, does the sphere's angular momentum about its center of mass change? Check all that apply.

☐ A to B ☐ B to C ☐ C to D ☐ D to E ☐ None

Briefly explain your reasoning.

2. (12 points, suggested time 25 minutes)

A new kind of toy ball is advertised to “bounce perfectly elastically” off hard surfaces. A student suspects, however, that no collision can be perfectly elastic. The student hypothesizes that the collisions are very close to being perfectly elastic for low-speed collisions but that they deviate more and more from being perfectly elastic as the collision speed increases.

- (a) Design an experiment to test the student’s hypothesis about collisions of the ball with a hard surface. The student has equipment that would usually be found in a school physics laboratory.
- What quantities would be measured?
 - What equipment would be used for the measurements, and how would that equipment be used?
 - Describe the procedure to be used to test the student’s hypothesis. Give enough detail so that another student could replicate the experiment.
- (b) Describe how you would represent the data in a graph or table. Explain how that representation would be used to determine whether the data are consistent with the student’s hypothesis.
- (c) A student carries out the experiment and analysis described in parts (a) and (b). The student immediately concludes that something went wrong in the experiment because the graph or table shows behavior that is elastic for low-speed collisions but appears to violate a basic physics principle for high-speed collisions.
- Give an example of a graph or table that indicates nearly elastic behavior for low-speed collisions but appears to violate a basic physics principle for high-speed collisions.
 - State one physics principle that appears to be violated in the graph or table given in part (c)i. Several physics principles might appear to be violated, but you only need to identify one.

Briefly explain what aspect of the graph or table indicates that the physics principle is violated, and why.