





. (7 points, suggested time 13 minutes)

A block of unknown mass is attached to a long, lightweight string that is wrapped several turns around a pulley mounted on a horizontal axis through its center, as shown. The pulley is a uniform solid disk of mass M and radius R . The rotational inertia of the pulley is described by the equation $I = \frac{1}{2}MR^2$. The pulley can rotate about its center with negligible friction. The string does not slip on the pulley as the block falls.

When the block is released from rest and as the block travels toward the ground, the magnitude of the tension exerted on the block by the string is F_T .

- (a) Determine an expression for the magnitude of the angular acceleration α_D of the disk as the block travels downward. Express your answer in terms of M , R , F_T , and physical constants as appropriate.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 4** on this page.

Scenarios 1 and 2 show two different pulleys. In Scenario 1, the pulley is the same solid disk referenced in part (a). In Scenario 2, the pulley is a hoop that has the same mass M and radius R as the disk. Each pulley has a lightweight string wrapped around it several turns and is mounted on a horizontal axle, as shown. Each pulley is free to rotate about its center with negligible friction.

In both scenarios, the pulleys begin at rest. Then both strings are pulled with the same constant force F_A for the same time interval Δt , causing the pulleys to rotate without the string slipping. After time interval Δt , the change in angular momentum of the disk is equal to the change in angular momentum of the hoop, but the change in rotational kinetic energy for the disk is greater than that of the hoop.

- (b) Consider scenarios 1 and 2 at the end of time interval Δt . In a clear, coherent paragraph-length response that may also contain equations and drawings, explain why the change in angular momentum of both pulleys is the same but the change in rotational kinetic energy is greater for the disk.

GO ON TO THE NEXT PAGE.




Figure 1

(7 points, suggested time 13 minutes)

A rod with a sphere attached to the end is connected to a horizontal mounted axle and carefully balanced so that it rests in a position vertically upward from the axle. The center of mass of the rod-sphere system is indicated with a $\otimes$, as shown in Figure 1. The sphere is lightly tapped, and the rod-sphere system rotates clockwise with negligible friction about the axle due to the gravitational force.

A student takes a video of the rod rotating from the vertically upward position to the vertically downward position. Figure 2 shows five frames (still shots) that the student selected from the video.

Note: these frames are not equally spaced apart in time.




Figure 2

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 5** on this page.

(a) Use the frames of the video shown in Figure 2 to answer the following questions.

- i. In which frame is the angular acceleration of the rod-sphere system the greatest? Justify your answer.

- ii. In which frame is the rotational kinetic energy of the rod-sphere system the greatest? Briefly justify your answer.




Figure 3

(b) The rod-sphere system has mass M and length L , and the center of mass is located a distance $\frac{3}{4}L$ from the axle, shown in Figure 3.


- i. Derive an expression for the change in kinetic energy of the rod-sphere-Earth system from the moment shown in Frame A to the moment shown in Frame E. Express your answer in terms of M , L , and fundamental constants, as appropriate.

- ii. Briefly explain why the rod and sphere gain kinetic energy, even if Earth is not included in the system.

GO ON TO THE NEXT PAGE.

STOP

END OF EXAM



. (12 points, suggested time 25 minutes)

A wheel is mounted on a horizontal axle. A light string is attached to the wheel's rim and wrapped around it several times, and a small block is attached to the free end of the string, as shown in the figure. When the block is released from rest and begins to fall, the wheel begins to rotate with negligible friction.

Two students are discussing how different forms of energy change as the block falls. One student says that the kinetic energy of the block increases as it falls. The second student says that this is because gravitational potential energy is converted to kinetic energy. The students decide to test whether the decrease in gravitational potential energy is equal to the increase in the block's kinetic energy from when the block starts moving to immediately before it reaches the floor.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 3** on this page.

- (a) Design an experimental procedure that the students could use to compare the increase in the block's translational kinetic energy with the decrease in the gravitational potential energy of the block-Earth system as the block falls.

In the table, list the quantities that would be measured in your experiment. Define a symbol to represent each quantity and list the equipment that would be used to measure each quantity. You do not need to fill in every row. If you need additional rows, you may add them to the space just below the table.


In the space to the right of the table, describe the overall procedure. Provide enough detail so that another student could replicate the experiment, including any steps necessary to reduce experimental uncertainty. As needed, use the symbols defined in the table.

If needed, you may include a simple diagram of the setup with your procedure.

Quantity to Be Measured	Symbol for Quantity	Equipment for Measurement	Procedure (and diagram, if needed)


- (b) Explain how the students could determine the kinetic energy of the block immediately before it reaches the floor using the quantities you indicated in the table in part (a).

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 3** on this page.


(c) The graph above represents both the change in the gravitational potential energy of the block-wheel-Earth system and the translational kinetic energy gained by the block as functions of the block's falling distance d . On the graph, draw a line or curve to represent the rotational kinetic energy of the wheel as a function of the block's falling distance d .

(d) The students also measure the angular velocity ω of the wheel as the block falls and determine the rotational kinetic energy K_R of the wheel. The students then make a graph of K_R as a function of ω^2 , as shown.



- On the above graph, draw a straight line that best represents the data.
- Using the line you drew for part (d)(i), calculate an experimental value for the rotational inertia of the wheel.


GO ON TO THE NEXT PAGE.



(7 points, suggested time 13 minutes)


A cylinder of mass m_0 is placed at the top of an incline of length L_0 and height H_0 , as shown above, and released from rest. The cylinder rolls without slipping down the incline and then continues rolling along a horizontal surface.

(a) On the grid below, sketch a graph that represents the total kinetic energy of the cylinder as a function of the distance traveled by the cylinder as it rolls down the incline and continues to roll across the horizontal surface.



GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.



The cylinder is again placed at the top of the incline. A block, also of mass m_0 , is placed at the top of a separate rough incline of length L_0 and height H_0 , as shown above. When the cylinder and block are released at the same instant, the cylinder begins to roll without slipping while the block begins to accelerate uniformly. The cylinder and the block reach the bottoms of their respective inclines with the same translational speed.


(b) In terms of energy, explain why the two objects reach the bottom of their respective inclines with the same translational speed. Provide your answer in a clear, coherent paragraph-length response that may also contain figures and/or equations.

GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

PHYSICS 1**Section II****5 Questions****Time—90 minutes**

Directions: Questions 1, 4 and 5 are short free-response questions that require about 13 minutes each to answer and are worth 7 points each. Questions 2 and 3 are long free-response questions that require about 25 minutes each to answer and are worth 12 points each. Show your work for each part in the space provided after that part.




1. (7 points, suggested time 13 minutes)

A wooden wheel of mass M , consisting of a rim with spokes, rolls down a ramp that makes an angle θ with the horizontal, as shown above. The ramp exerts a force of static friction on the wheel so that the wheel rolls without slipping.

(a)

- i. On the diagram below, draw and label the forces (not components) that act on the wheel as it rolls down the ramp, which is indicated by the dashed line. To clearly indicate at which point on the wheel each force is exerted, draw each force as a distinct arrow starting on, and pointing away from, the point at which the force is exerted. The lengths of the arrows need not indicate the relative magnitudes of the forces.



- ii. As the wheel rolls down the ramp, which force causes a change in the angular velocity of the wheel with respect to its center of mass?

Briefly explain your reasoning.

- (b) For this ramp angle, the force of friction exerted on the wheel is less than the maximum possible static friction force. Instead, the magnitude of the force of static friction exerted on the wheel is 40 percent of the magnitude of the force or force component directed opposite to the force of friction. Derive an expression for the linear acceleration of the wheel's center of mass in terms of M , θ , and physical constants, as appropriate.

- (c) In a second experiment on the same ramp, a block of ice, also with mass M , is released from rest at the same instant the wheel is released from rest, and from the same height. The block slides down the ramp with negligible friction.

i. Which object, if either, reaches the bottom of the ramp with the greatest speed?

____ Wheel ____ Block ____ Neither; both reach the bottom with the same speed.

Briefly explain your answer, reasoning in terms of forces.

ii. Briefly explain your answer again, now reasoning in terms of energy.