

Begin your response to **QUESTION 2** on this page.




Figure 1

2. (12 points, suggested time 25 minutes)

A student hangs a spring of unknown spring constant  $k$  vertically by attaching one end to a stand, as shown in Figure 1. The other end of the spring has a small loop from which small cylinders can be hung. In addition to the spring, the student has access only to a variety of cylinders of unknown masses, a stopwatch, and a digital scale.

(a) Design an experimental procedure the student could use to determine the spring constant  $k$  of the spring.

In the following table, list the quantities that would be measured using only the provided equipment in your experiment. Define a symbol to represent each quantity.

In the space below the table, **describe** the overall procedure. Provide enough detail so that another student could replicate the experiment, including any steps necessary to reduce experimental uncertainty. As needed, use the symbols defined in the table. If needed, you may include a simple diagram of the setup with your procedure.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 2** on this page.

| Quantity to Be Measured | Symbol for Quantity | Equipment for Measurement |
|-------------------------|---------------------|---------------------------|
|                         |                     | Stopwatch                 |
|                         |                     | Digital scale             |
|                         |                     |                           |

Procedure (and diagram, if needed)

(b)

i. **Indicate** the quantities that could be plotted to produce a linear graph whose slope can be used to determine the spring constant  $k$  of the spring.

Vertical axis: \_\_\_\_\_ Horizontal axis: \_\_\_\_\_

ii. Briefly **describe** how the slope of the graph would be analyzed to determine the spring constant  $k$  of the spring.

GO ON TO THE NEXT PAGE.




Continue your response to **QUESTION 2** on this page.

Figure 2

In a different experiment, the student attaches one end of a spring to a force sensor that is attached to a wall. The other end of the spring is attached to a cart with mass  $m = 0.25$  kg. The student places a motion detector to the right of the cart, as shown in Figure 2, and pulls the cart to the right a small distance so that the spring is stretched. The student releases the cart from rest, and the cart-spring system oscillates.

The following graphs show the velocity  $v$  of the cart and the force  $F$  exerted on the cart by the spring as functions of time  $t$ .



GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 2** on this page.


(c)

i. Using the data in the velocity-time graph, **calculate** the change in kinetic energy of the cart from  $t = 0.5$  s to  $t = 2.0$  s. Show your steps and substitutions.

ii. Using the data in the force-time graph, **estimate** the change in momentum of the cart from  $t = 0.5$  s to  $t = 2.5$  s. Briefly **explain** how you arrived at your estimation.

iii. Do the data from the velocity-time graph confirm your estimation from part (c)(ii) ? Briefly **explain**.

GO ON TO THE NEXT PAGE.




(7 points, suggested time 13 minutes)

A spring of unknown spring constant  $k_0$  is attached to a ceiling. A lightweight hanger is attached to the lower end of the spring, and a motion detector is placed on the floor facing upward directly under the hanger, as shown in the figure above. The bottom of the hanger is 1.00 m above the motion detector.

A 0.50 kg object is placed on the hanger and allowed to come to rest at the equilibrium position. The spring is then stretched downward a distance  $d_0$  from equilibrium and released at time  $t = 0$ . The motion detector records the height of the bottom of the hanger as a function of time. The output from the motion detector is shown in the graph on the following page.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 5** on this page.



(a) Using the information given and information taken from the graph, calculate the spring constant.

(b) At time 0.75 s, the object-spring-Earth system has a total kinetic energy  $K_0$  and a total potential energy  $U_0$ . At 1.13 s, the object-spring-Earth system again has a total kinetic energy  $K_0$  and a total potential energy  $U_0$ .

i. Explain how a feature of the graph indicates that the total kinetic energy of the system is the same at these two times.

ii. Briefly explain why the total potential energy of the system is the same at these two times.


GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 5** on this page.


(c) The experiment is repeated with a spring of spring constant  $4k_0$  and that has the same length as the original spring. The 0.50 kg object is hung from the new spring and allowed to come to rest at a new equilibrium position.


i. Determine the new equilibrium position above the motion detector.

ii. The object is again pulled down the same distance  $d_0$  from the equilibrium position and released. On the following graph, draw a curve representing the motion of the object after it is released. Label the vertical axis with an appropriate numerical scale. A grid for scratch (practice) work is also provided.



The following graph is provided for scratch work only and will not be graded.


**GO ON TO THE NEXT PAGE.****STOP****END OF EXAM**



5. (7 points, suggested time 13 minutes)

A tuning fork vibrating at 512 Hz is held near one end of a tube of length  $L$  that is open at both ends, as shown above. The column of air in the tube resonates at its fundamental frequency. The speed of sound in air is 340 m/s.


- (a) Calculate the length  $L$  of the tube.
- (b) The column of air in the tube is still resonating at its fundamental frequency. On the axes below, sketch a graph of the maximum speed of air molecules as they oscillate in the tube, as a function of position  $x$ , from  $x = 0$  (left end of tube) to  $x = L$  (right end of tube). (Ignore random thermal motion of the air molecules.)



- (c) The right end of the tube is now capped shut, and the tube is placed in a chamber that is filled with another gas in which the speed of sound is 1005 m/s. Calculate the new fundamental frequency of the tube.

**STOP**

**END OF EXAM**




5. (7 points, suggested time 13 minutes)

The figure above on the left shows a uniformly thick rope hanging vertically from an oscillator that is turned off. When the oscillator is on and set at a certain frequency, the rope forms the standing wave shown above on the right.  $P$  and  $Q$  are two points on the rope.

- (a) The tension at point  $P$  is greater than the tension at point  $Q$ . Briefly explain why.
- (b) A student hypothesizes that increasing the tension in a rope increases the speed at which waves travel along the rope. In a clear, coherent paragraph-length response that may also contain figures and/or equations, explain why the standing wave shown above supports the student's hypothesis.


**STOP**

**END OF EXAM**



5. (7 points, suggested time 13 minutes)


The figure above shows a string with one end attached to an oscillator and the other end attached to a block. The string passes over a massless pulley that turns with negligible friction. Four such strings, *A*, *B*, *C*, and *D*, are set up side by side, as shown in the diagram below. Each oscillator is adjusted to vibrate the string at its fundamental frequency  $f$ . The distance between each oscillator and pulley  $L$  is the same, and the mass  $M$  of each block is the same. However, the fundamental frequency of each string is different.



The equation for the velocity  $v$  of a wave on a string is  $v = \sqrt{\frac{F_T}{m/L}}$ , where  $F_T$  is the tension of the string and  $m/L$  is the mass per unit length (linear mass density) of the string.

- What is different about the four strings shown above that would result in their having different fundamental frequencies? Explain how you arrived at your answer.
- A student graphs frequency as a function of the inverse of the linear mass density. Will the graph be linear? Explain how you arrived at your answer.

- The frequency of the oscillator connected to string *D* is changed so that the string vibrates in its second harmonic. On the side view of string *D* below, mark and label the points on the string that have the greatest average vertical speed.



**STOP**

**END OF EXAM**