





. (7 points, suggested time 13 minutes)

A block of unknown mass is attached to a long, lightweight string that is wrapped several turns around a pulley mounted on a horizontal axis through its center, as shown. The pulley is a uniform solid disk of mass M and radius R . The rotational inertia of the pulley is described by the equation $I = \frac{1}{2}MR^2$. The pulley can rotate about its center with negligible friction. The string does not slip on the pulley as the block falls.

When the block is released from rest and as the block travels toward the ground, the magnitude of the tension exerted on the block by the string is F_T .

- (a) Determine an expression for the magnitude of the angular acceleration α_D of the disk as the block travels downward. Express your answer in terms of M , R , F_T , and physical constants as appropriate.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 4** on this page.

Scenarios 1 and 2 show two different pulleys. In Scenario 1, the pulley is the same solid disk referenced in part (a). In Scenario 2, the pulley is a hoop that has the same mass M and radius R as the disk. Each pulley has a lightweight string wrapped around it several turns and is mounted on a horizontal axle, as shown. Each pulley is free to rotate about its center with negligible friction.

In both scenarios, the pulleys begin at rest. Then both strings are pulled with the same constant force F_A for the same time interval Δt , causing the pulleys to rotate without the string slipping. After time interval Δt , the change in angular momentum of the disk is equal to the change in angular momentum of the hoop, but the change in rotational kinetic energy for the disk is greater than that of the hoop.

- (b) Consider scenarios 1 and 2 at the end of time interval Δt . In a clear, coherent paragraph-length response that may also contain equations and drawings, explain why the change in angular momentum of both pulleys is the same but the change in rotational kinetic energy is greater for the disk.

GO ON TO THE NEXT PAGE.




Figure 1

(7 points, suggested time 13 minutes)

A rod with a sphere attached to the end is connected to a horizontal mounted axle and carefully balanced so that it rests in a position vertically upward from the axle. The center of mass of the rod-sphere system is indicated with a $\otimes$, as shown in Figure 1. The sphere is lightly tapped, and the rod-sphere system rotates clockwise with negligible friction about the axle due to the gravitational force.

A student takes a video of the rod rotating from the vertically upward position to the vertically downward position. Figure 2 shows five frames (still shots) that the student selected from the video.

Note: these frames are not equally spaced apart in time.




Figure 2

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 5** on this page.

(a) Use the frames of the video shown in Figure 2 to answer the following questions.

- i. In which frame is the angular acceleration of the rod-sphere system the greatest? Justify your answer.

- ii. In which frame is the rotational kinetic energy of the rod-sphere system the greatest? Briefly justify your answer.




Figure 3

(b) The rod-sphere system has mass M and length L , and the center of mass is located a distance $\frac{3}{4}L$ from the axle, shown in Figure 3.

- i. Derive an expression for the change in kinetic energy of the rod-sphere-Earth system from the moment shown in Frame A to the moment shown in Frame E. Express your answer in terms of M , L , and fundamental constants, as appropriate.

- ii. Briefly explain why the rod and sphere gain kinetic energy, even if Earth is not included in the system.


GO ON TO THE NEXT PAGE.

STOP

END OF EXAM

PHYSICS 1**Section II****Time—1 hour and 30 minutes****5 Questions**


Directions: Questions 1, 4, and 5 are short free-response questions that require about 13 minutes each to answer and are worth 7 points each. Questions 2 and 3 are long free-response questions that require about 25 minutes each to answer and are worth 12 points each. Show your work for each part in the space provided after that part.




1. (7 points, suggested time 13 minutes)

Identical blocks 1 and 2 are placed on a horizontal surface at points A and E, respectively, as shown. The surface is frictionless except for the region between points C and D, where the surface is rough. Beginning at time t_A , block 1 is pushed with a constant horizontal force from point A to point B by a mechanical plunger. Upon reaching point B, block 1 loses contact with the plunger and continues moving to the right along the horizontal surface toward block 2. Block 1 collides with and sticks to block 2 at point E, after which the two-block system continues moving across the surface, eventually passing point F.

- (a) On the axes below, sketch the speed of the center of mass of the two-block system as a function of time, from time t_A until the blocks pass point F at time t_F . The times at which block 1 reaches points A through F are indicated on the time axis.





- (b) The plunger is returned to its original position, and both blocks are removed. A uniform solid sphere is placed at point A, as shown. The sphere is pushed by the plunger from point A to point B with a constant horizontal force that is directed toward the sphere's center of mass. The sphere loses contact with the plunger at point B and continues moving across the horizontal surface toward point E. In which interval(s), if any, does the sphere's angular momentum about its center of mass change? Check all that apply.

_____ A to B _____ B to C _____ C to D _____ D to E _____ None

Briefly explain your reasoning.