

9 Temperature, gases, and heat – Part 1

Name: _____ Class: _____ Date: _____

Vocabulary

Write each word three times.

English	Write it 3 times (English)		
Thermodynamics	_____	_____	_____
kinetic theory	_____	_____	_____
Temperature	_____	_____	_____
pressure	_____	_____	_____
ideal gas	_____	_____	_____
Heat	_____	_____	_____
thermal equilibrium	_____	_____	_____
conduction	_____	_____	_____
convection	_____	_____	_____
radiation	_____	_____	_____
specific heat	_____	_____	_____

Recall

You already know a lot about heat and gases:


- Hot things feel hot because their tiny particles move fast; cold things move slowly.
- A gas pushes out on its container – a balloon stays full because the air pushes on it.
- Heat always flows from the hot object to the cold one, never the other way.

This sheet gives these ideas exact formulas. Each idea has a worked example.

New ideas

■ 1 Temperature and moving particles

Thermodynamics is the study of heat and energy in large groups of particles. The **kinetic theory** says a gas is countless tiny particles in fast, random motion.



1. many identical molecules, in continuous **random motion**
2. their own volume is **tiny** next to the container
3. straight lines between hits - intermolecular forces **ignored**
4. collisions are **elastic**: no kinetic energy is lost

poor at very high pressure (volume matters) or very low temperature (forces matter)

Temperature measures the **average kinetic energy** of those particles: hotter means faster-moving. The **pressure** a gas makes comes from its particles hitting the container walls.

■ 2 The ideal gas law

An **ideal gas** obeys


$$PV = nRT,$$

linking pressure P , volume V , amount n , and temperature T (always in **kelvin**). Change one quantity and the law tells you how another responds.

Worked example. A sealed rigid can of gas is at 1.0×10^5 Pa and 300 K. Heated to 450 K at fixed volume, $\frac{P}{T}$ stays constant, so $P_2 = P_1 \frac{T_2}{T_1} = 1.0 \times 10^5 \times \frac{450}{300} = 1.5 \times 10^5$ Pa.

■ 3 Heat and how it moves

Heat Q is energy that flows because of a temperature difference – always from hot to cold. When two objects reach the same temperature they are in **thermal equilibrium** and the flow stops. Heat travels by **conduction** (through solids), **convection** (moving fluids), and **radiation** (waves, even through empty space).



■ 4 Specific heat

The **specific heat** c is the heat needed to warm 1 kg of a substance by 1°C :

$$Q = mc \Delta T.$$

Worked example. Heating 2.0 kg of water ($c = 4200$) from 20°C to 80°C needs $Q = mc \Delta T = 2.0 \times 4200 \times 60 = 5.0 \times 10^5 \text{ J}$.

Watch out: always convert temperatures to **kelvin** ($T_K = T_C + 273$) before using a gas law – using Celsius gives nonsense.

Practice

Try these in order. The methods are all above.

2.1 State what temperature measures, in terms of the particles of a gas. [1]

2.2 Convert 27°C to kelvin. [1]

2.3 A rigid can of gas at $2.0 \times 10^5 \text{ Pa}$ and 300 K is cooled to 150 K . Find the new pressure. [3]

2.4 Name the three ways heat can travel, and give the one that works through empty space. [2]

2.5 How much heat warms 0.50 kg of water ($c = 4200$) by 30°C ? [2]

2.6 Explain why coastal towns have milder temperatures than inland ones (think about water's specific heat). [2]