

PHYSICS 2

SECTION II

Time—1 hour and 30 minutes

4 Questions

Directions: Questions 1 and 4 are short free-response questions that require about 20 minutes each to answer and are worth 10 points each. Questions 2 and 3 are long free-response questions that require about 25 minutes each to answer and are worth 12 points each. Show your work for each part in the space provided after that part.




Figure 1

(10 points, suggested time 20 minutes)

A rectangular tank with a mirrored bottom is filled with water (index of refraction n_w). A beam of light passes from air (index of refraction n_a) into the water at angle θ_i from the normal, as shown in Figure 1. Index of refraction n_w is greater than index of refraction n_a .

- (a) On the following diagram, sketch the entire path of the beam as the beam enters, travels through, and then exits the water.



GO ON TO THE NEXT PAGE.


Continue your response to **QUESTION 1** on this page.

Figure 2

Sugar is then added to the water, resulting in a mixture that has a different index of refraction than water. A student considers two models, Model A and Model B, for how the sugar mixes with water. The models are shown in Figure 2.

Model A: The sugar is uniformly mixed throughout the water, resulting in a mixture with index of refraction n_m such that $n_m > n_w$.

Model B: Layers are formed of varying concentrations of sugar in the water. There are three distinct layers of equal volume. The top layer is only water (index of refraction n_w). The middle layer has the same concentration of sugar as the mixture in Model A (index of refraction n_m). The bottom layer has the highest concentration of sugar (index of refraction n_b).

- (b) Consider Model A. Briefly describe how the observed wavelength of light changes, if at all, as the beam travels from air into the mixture.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 1** on this page.

- (c) Relevant angles between the beam and the normal for the various layers present in models A and B are defined in the following table.

Model A		Model B	
θ_i	Incident angle of the beam in air	θ_i	Incident angle of the beam in air
θ_1	Angle the beam makes with the normal in the mixture in Model A	θ_2	Angle the beam makes with the normal in the top layer in Model B
		θ_3	Angle the beam makes with the normal in the middle layer in Model B
		θ_4	Angle the beam makes with the normal in the bottom layer in Model B


- i. Determine an expression for θ_4 in terms of θ_i , n_a , and n_b .

- ii. Rank the angles from greatest to least, with 1 being greatest. If two angles are the same value, give them the same ranking.


_____ θ_1 _____ θ_2 _____ θ_3 _____ θ_4

Briefly explain your reasoning using appropriate physics principles and/or mathematical models.


GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 1** on this page.

Original Tank



Model A



Model B

For the original tank filled with water, the beam is observed to exit the surface of the water a horizontal distance d_w from the entry point. For models A and B, the horizontal distances are d_A and d_B , respectively.

- (d) Determine whether d_A and d_B are each greater than, less than, or equal to d_w . It is NOT necessary to compare d_A to d_B . Briefly justify your answer.

GO ON TO THE NEXT PAGE.

Begin your response to QUESTION 1 on this page.


PHYSICS 2

SECTION II

Time—1 hour and 30 minutes

4 Questions

Directions: Questions 1 and 4 are short free-response questions that require about 20 minutes each to answer and are worth 10 points each. Questions 2 and 3 are long free-response questions that require about 25 minutes each to answer and are worth 12 points each. Show your work for each part in the space provided after that part.



(10 points, suggested time 20 minutes)

Students are investigating electromagnetic wave phenomena in transparent media. They use a string to support a stationary thin, rectangular block of mass m_b , volume V_b , and density ρ_b . The block has two narrow slits in its center and is submerged in a glass tank containing water with density ρ_w , as shown above.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 1** on this page.

(a)

i. On the dot below, which represents the block, draw and label the forces that are exerted on the block. Each force must be represented by a distinct arrow starting on, and pointing away from, the dot.




ii. Derive an expression for the force exerted on the block by the string in terms of the given quantities and physical constants, as appropriate.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 1** on this page.

- (b) A monochromatic laser beam is incident perpendicular to the wall of the tank. The beam passes through the slits in the block. An interference pattern is formed on the screen inside the tank. The water is then replaced with a clear fluid with a greater index of refraction than that of water. In a coherent, paragraph-length response, describe how the greater index of refraction of the new fluid affects the interference pattern. Explain your reasoning in terms of speed, frequency, and wavelength of the light.




- (c) The block is replaced by a triangular prism, as shown above. The path of the beam is indicated by the dotted line, and the beam reaches the screen at point P . The fluid is then removed from the tank, and the prism is surrounded by air. Predict whether the beam will reach the side of the tank above point P , at point P , or below point P when the prism is surrounded by air. Support your answer using physics principles.

GO ON TO THE NEXT PAGE.

4. (10 points, suggested time 20 minutes)

A student notices many air bubbles rising through the water in a large fish tank at an aquarium.

- (a) In the figure below, the circle represents one such air bubble, and two incoming rays of light, A and B, are shown. Ray B points toward the center of the circle. On the diagram, draw the paths of rays A and B as they go through the bubble and back into the water. Your diagram should clearly show what happens to the rays at each interface.



- (b) The bubble has a volume V_1 , the air inside it has density ρ_A , and the water around it has density ρ_W . The bubble starts at rest and has a speed v_f when it has risen a height h . Assume that the change in the bubble's volume is negligible. Derive an expression for the mechanical energy dissipated by drag forces as the bubble rises this distance. Express your answer in terms of the given quantities and fundamental constants, as appropriate.
- (c) At a particular instant, one bubble is 4.5 m below the water's surface. The surface of the water is at sea level, and the density of the water is 1000 kg/m^3 .
- Determine the absolute pressure in the bubble at this location.
 - The bubble has a volume V_1 when it is 4.5 m below the water's surface. Assume that the temperature of the air in the bubble remains constant as it rises. In terms of V_1 , calculate the volume of the bubble when it is just below the surface of the water.
 - If the air in the bubble cooled as it rose, the volume of the bubble would be less than the value calculated in part (c)(ii). Use physics principles to briefly explain why.

STOP

END OF EXAM