

# 12.3 Magnetic Fields of Current-Carrying Wires and the Biot-Savart Law

Name: \_\_\_\_\_ Class: \_\_\_\_\_ Date: \_\_\_\_\_

Total: 9 marks

KEY WORDS biot-savart law 毕奥 • magnetic field 磁场

## Objective

Build the skills to answer exam questions on **magnetic fields of current-carrying wires and the Biot-Savart law**.

You must be able to:

- state the **Biot-Savart law** 毕奥-萨伐尔定律,  $d\vec{B} = \frac{\mu_0}{4\pi} \frac{I d\vec{l} \times \hat{r}}{r^2}$
- calculate the field of a **long straight wire**,  $B = \frac{\mu_0 I}{2\pi r}$
- find the field at the **centre of a circular loop**,  $B = \frac{\mu_0 I}{2R}$
- use the **right-hand rule** for field direction

## 1 Worked examples

Study these first. Each one shows the method for a question type used later.

### ■ The Biot-Savart law

$d\vec{B} = \frac{\mu_0}{4\pi} \frac{I d\vec{l} \times \hat{r}}{r^2}$  gives the field of a small current element; integrate over the wire for the total field ( $\mu_0 = 4\pi \times 10^{-7}$ ).

### ■ Standard results

- Long straight wire:  $B = \frac{\mu_0 I}{2\pi r}$ .
- Centre of a circular loop of radius  $R$ :  $B = \frac{\mu_0 I}{2R}$ .

### ■ Direction

Curl the right hand around the wire with the thumb along the current; the fingers give the field direction.

## 2 Practice

---

**2.1** Find the field 0.050 m from a long wire carrying 8.0 A ( $\mu_0 = 4\pi \times 10^{-7}$ ). [2]

---

---

**2.2** State the rule used to find the direction of the field around a straight wire. [1]

---

**2.3** Write the field at the centre of a circular loop of radius  $R$  carrying current  $I$ . [1]

---

## 3 Exam-style questions

---

**3.1** The magnetic field of a long straight wire is [1]

- |   |   |
|---|---|
| A | B |
| C | D |
- A**  $\frac{\mu_0 I}{2\pi r}$   
**B**  $\frac{\mu_0 I}{2R}$   
**C**  $\mu_0 n I$   
**D**  $BIL$

**3.2** The Biot-Savart law gives the field of a [1]

- |   |   |
|---|---|
| A | B |
| C | D |
- A** current element  
**B** point charge  
**C** capacitor  
**D** resistor

**3.3** A circular loop of radius 0.10 m carries 5.0 A ( $\mu_0 = 4\pi \times 10^{-7}$ ).

(a) Find the magnetic field at its centre. [2]

(b) State how the field changes if the current is doubled. [1]