6.1 Rotational Kinetic Energy

Name: _____ Class: _____ Date: _____ **Total: 14 marks**

KEY WORDS rotational kinetic energy

Objective

Build the skills to answer exam questions on **rotational kinetic energy**.

You must be able to:

- use **rotational kinetic energy** : $K = \frac{1}{2}I\omega^2$
- recognise it as the analogue of $\frac{1}{2}mv^2$
- add translational and rotational kinetic energy for a rolling object
- find K, I, or ω given the others

1 Key knowledge and worked examples

Study these first. Each one shows the method for a question type used later – follow the steps and you can do the Practice and Exam-style questions yourself.

■ **The rotational KE formula**

$$K = \frac{1}{2}I\omega^2.$$

A disc with $I = 0.5$ kg m^2 spinning at $\omega = 4$ rad s^{-1} stores

$$K = \frac{1}{2}(0.5)(4)^2 = 4 \text{ J}.$$

■ **The square of omega**

Because ω is squared, doubling the spin rate quadruples the stored energy.

■ **Rolling: two kinds of KE**

A rolling object has both, so its total is

$$K = \frac{1}{2}mv^2 + \frac{1}{2}I\omega^2.$$

2 Practice

Now apply the methods above.

2.1 A wheel with $I = 2$ kg m^2 spins at 3 rad s^{-1}. Find its rotational KE. [2]

2.2 If a spinning object's angular speed triples, by what factor does its rotational KE change? [1]

2.3 State the two terms in the total kinetic energy of a rolling object. [2]

2.4 A flywheel stores 18 J of rotational KE at $I = 4$ kg m^2. Find its angular speed. [2]

3 Exam-style questions

3.1 Rotational kinetic energy is [1]

☐ **A** $\frac{1}{2}mv^2$
B $\frac{1}{2}I\omega^2$
C $I\omega$
D $\frac{1}{2}kx^2$

3.2 A rolling ball's total kinetic energy is [1]

☐ **A** only $\frac{1}{2}mv^2$
B only $\frac{1}{2}I\omega^2$
C $\frac{1}{2}mv^2 + \frac{1}{2}I\omega^2$
D zero

3.3 A solid disc has $I = 0.8$ kg m^2 and spins at 5 rad s^{-1}.

(a) Find its rotational kinetic energy. [2]

(b) State its energy if the spin rate doubles. [1]

3.4 A rolling wheel has translational KE = 12 J and rotational KE = 6 J.

(a) Find the total kinetic energy. [1]

(b) State the fraction stored in rotation. [1]