


Energy & Momentum of Rotating Systems


AP Physics C: Mechanics ■ Unit 6

AP Physics C: Mechanics



What we will cover

- 1 Rotational kinetic energy
- 2 Work and power of a torque
- 3 Angular momentum and impulse
- 4 Conservation of angular momentum
- 5 Rolling without slipping
- 6 Orbiting satellites





Rotational kinetic energy

Energy of spin

$$K_{rot} = \frac{1}{2} I \omega^2$$

A rolling body has **both**: $K = \frac{1}{2} m v^2 + \frac{1}{2} I \omega^2$.

A **flywheel** stores energy in its spin – and $K \propto \omega^2$.

Torque and work

$$W = \int \tau d\theta, \quad P = \tau \omega$$

Area under a τ - θ graph; $W_{net} = \Delta K_{rot}$.

$\tau = 15$, $\Delta\theta = 8$: $W = 120$ J; at $\omega = 20$,
 $P = 300$ W.

The whole energy toolkit, translated

kinetic energy

$$\frac{1}{2}mv^2 \longrightarrow \frac{1}{2}I\omega^2$$

work

$$W = \int F dx \longrightarrow W = \int \tau d\theta$$

power

$$P = Fv \longrightarrow P = \tau\omega$$

momentum

$$\vec{p} = m\vec{v} \longrightarrow \vec{L} = I\vec{\omega}$$

the substitution

$$F \rightarrow \tau, \quad x \rightarrow \theta, \quad v \rightarrow \omega, \quad m \rightarrow I$$

There is nothing new to learn in rotational energy – only a dictionary. Every translational result you already have comes across intact.



Angular momentum and impulse

Angular momentum is the rotational twin of momentum:

Point particle and rigid body

$$\vec{L} = \vec{r} \times \vec{p}, \quad L = I\omega$$

$$\vec{\tau}_{net} = \frac{d\vec{L}}{dt}$$

Angular impulse $\int \tau dt = \Delta L$.

Worked example

$L_i = 30$; $\tau = 5 \text{ N m}$ for 4 s:

$$\Delta L = 5 \times 4 = 20$$

$$L_f = 30 + 20 = 50 \text{ kg m}^2/\text{s}$$

Conservation of angular momentum

With **zero net external torque**, angular momentum is conserved – even as the shape changes:

Before = after

$$I_i \omega_i = I_f \omega_f$$

A skater pulls their arms in: I drops, so ω rises. (Kinetic energy rises too – paid by muscular work.)



Skater spin


Worked example – a bullet strikes a pivoted rod

A 0.020 kg bullet at 300 m/s hits the tip of a uniform rod ($M = 1.5$ kg, $l = 0.60$ m) pivoted at its centre, and embeds. Find ω .

- angular momentum about the pivot is conserved; **linear momentum is not** (the pivot exerts a force)
- before: $L = mvr = (0.020)(300)(0.30) = 1.8 \text{ kg m}^2/\text{s}$
- after:
$$I = \frac{1}{12}Ml^2 + mr^2 = 0.045 + 0.0018 = 0.0468$$
- $\omega = L/I = 38 \text{ rad/s}$

Choose **angular** momentum whenever there is a pivot. The pivot's own force acts through the axis, so it exerts no torque and drops out.


Pulling mass inward lowers I – one



$$L = I\omega \text{ conserved (no external torque): } I_1\omega_1 = I_2\omega_2$$

Pulling mass inward lowers I , so ω rises to conserve $L = I\omega$

Pulling mass inward lowers I – two



$$m v l = I_{\text{sys}} \omega \quad (\text{about pivot; } KE \text{ not conserved — inelastic})$$

Angular momentum about the pivot is conserved as the bullet embeds in the rod



Rolling without slipping


Rolling without slipping links translation and rotation:

The condition


$$v_{cm} = \omega r, \quad a_{cm} = \alpha r$$

Static friction produces the rolling but does **no work** (the contact point is momentarily at rest).

Down a ramp: the solid sphere wins, the hoop last.



Rolling race: the shape with the smallest – one



smallest $\beta \Rightarrow$ largest a (sphere wins)
 rolling race: smaller I reaches bottom first

Rolling race: the shape with the smallest I/mr^2 reaches the bottom first

Where the energy goes when a shape rolls

the split

a rolling body has **both** translational and rotational kinetic energy

write $I = \beta mr^2$

β is the shape factor: hoop 1, disc $\frac{1}{2}$, solid sphere $\frac{2}{5}$, spherical shell $\frac{2}{3}$

racing down an incline

$$a_{\text{cm}} = \frac{g \sin \theta}{1 + \beta}, \text{ from } mgh = \frac{1}{2}mv^2(1 + \beta)$$

so

$$v = \sqrt{\frac{2gh}{1 + \beta}} - \text{mass and radius both cancel}$$

the winner

the **smallest** β : solid sphere, then disc, then shell, then hoop

what β measures

how much **rotational inertia** sits far from the axis

Mass and radius cancel, so the race is decided by shape alone – and $\Delta x_{\text{cm}} = r\Delta\theta$ ties the centre's displacement to the **angular displacement**.

The rolling race, decided by one number

write $I = \beta mr^2$

β is the shape factor: hoop 1, disc $\frac{1}{2}$, solid sphere $\frac{2}{5}$, shell $\frac{2}{3}$

energy conservation

$$mgh = \frac{1}{2}mv^2(1 + \beta), \text{ so } v = \sqrt{\frac{2gh}{1 + \beta}}$$

so the winner


the **smallest** β – the solid sphere, then the disc, then the shell, then the hoop

and the mass?

cancels completely: size and mass do not matter, **only the shape**

β measures how much mass sits far from the axis. Mass at the rim has to be spun up as well as moved forward, and that energy is taken from the translation.

Rolling race: the shape with the smallest – two



$$K = \frac{1}{2}mv^2 + \frac{1}{2}I\omega^2 = \frac{1}{2}mv^2\left(1 + \frac{I}{mr^2}\right)$$

rolling no slip: $v = r\omega \quad \cdot \quad a = r\alpha$

In rolling without slipping the contact point is at rest, so $v = r\omega$




Orbiting satellites

Gravity supplies the centripetal force. Set them equal:

Orbital speed and Kepler

$$v = \sqrt{\frac{GM}{r}}, \quad T^2 \propto r^3$$

- energy $E = K + U$, with $U = -GMm/r$
- elliptical orbits conserve energy *and* angular momentum



A bigger orbit means
a **slower** speed, longer period.

The energy of an orbit is negative, and that matters

gravitational potential energy

$$U_g = -\frac{GMm}{r}, \text{ taking zero at infinity}$$

a circular orbit

$$K = -\frac{1}{2}U, \text{ so the total mechanical energy is } E = K + U = -\frac{GMm}{2r}$$

negative means bound

the satellite cannot escape; raising it to a larger orbit means *increasing* E toward zero

escape velocity


$$\text{set } E = 0: v_{\text{esc}} = \sqrt{\frac{2GM}{r}} = \sqrt{2} \times \text{the circular orbital speed}$$

Kepler's second law

equal areas in equal times – which is conservation of angular momentum in disguise

Larger orbits are **slower** ($v = \sqrt{GM/r}$) yet carry **more** energy. Speeding a satellite up moves it to a higher, slower orbit – the counter-intuitive result AP loves to test.

Motion of Orbiting Satellites



Gravity provides the centripetal force that keeps a satellite in orbit

Worked example – a satellite in low Earth orbit

A satellite orbits at $r = 7.0 \times 10^6$ m, with $GM = 4.0 \times 10^{14} \text{ m}^3/\text{s}^2$. Find its speed and escape speed.

- $v = \sqrt{GM/r} = \sqrt{4.0 \times 10^{14} / 7.0 \times 10^6} = 7.6 \times 10^3 \text{ m/s}$
- $v_{\text{esc}} = \sqrt{2} v = 1.1 \times 10^4 \text{ m/s}$
- period $T = 2\pi r/v \approx 5.8 \times 10^3 \text{ s}$, about 97 minutes

Exam skill. Orbit FRQs are energy-and-angular-momentum problems in disguise. Write $E = \frac{1}{2}mv^2 - \frac{GMm}{r}$ and $L = mvr \sin \theta$ at the **two points of interest** and solve the pair – never assume the circular-orbit formulae apply to an ellipse.

Exam tips

- Compute the **moment of inertia** $I = \int r^2 dm$ and shift axes with the **parallel-axis theorem** $I = I_{cm} + Md^2$.
- Rotational kinetic energy is $\frac{1}{2}I\omega^2$; a rolling body has both translational and rotational KE.
- Conserve **angular momentum** $L = I\omega$ when net external torque is zero (a spinning skater pulling in).
- Use energy conservation for rolling-without-slipping problems ($v = r\omega$ ties the two motions).
- Know standard I values (hoop, disk, rod, sphere) and where the axis is.



Unit 6 – key takeaways

1

Rotational energy

$$K_{rot} = \frac{1}{2} I \omega^2; W = \int \tau d\theta$$

2

Angular momentum

$$\vec{L} = \vec{r} \times \vec{p} = I\omega; \vec{\tau} = d\vec{L}/dt$$

3

Conservation

$$\text{no external torque} \Rightarrow I_i \omega_i = I_f \omega_f$$


4

Orbits

$$v = \sqrt{GM/r}, \text{ Kepler's } T^2 \propto r^3$$

Check yourself

- 1 Write the net torque as a rate of change of angular momentum.
- 2 A skater pulls their arms in – what happens to ω ?
- 3 Does the static friction on a rolling wheel do work?
- 4 A satellite moves to a bigger orbit – faster or slower?



Answers 1. $\vec{\tau} = d\vec{L}/dt$ 2. it speeds up 3. no 4. slower