

1.2 Displacement, Velocity, and Acceleration

Name: _____ Class: _____ Date: _____

Total: 15 marks

KEY WORDS acceleration 加速度 • velocity 速度 • integration 积分 • derivatives 导数

Objective

Build the skills to answer exam questions on **displacement, velocity, and acceleration** — the calculus links between them.

You must be able to:

- use **velocity** 速度 as the derivative of position, $v = \frac{dx}{dt}$
- use **acceleration** 加速度 as the derivative of velocity, $a = \frac{dv}{dt}$
- go the other way by **integration** 积分: $x = \int v \, dt$ and $v = \int a \, dt$
- read velocity as the slope of an x - t graph and displacement as the area under a v - t graph

1 Worked examples

Study these first. Each one shows the method for a question type used later — follow the steps and you can do the Practice and Exam-style questions yourself.

Derivatives: position to velocity to acceleration

Differentiate to move down the chain. If $x(t) = 3t^2$, then

$$v = \frac{dx}{dt} = 6t, \quad a = \frac{dv}{dt} = 6.$$

At $t = 2$ s: $v = 12 \text{ m s}^{-1}$, $a = 6 \text{ m s}^{-2}$.

Integration: acceleration back to velocity

Integrate to move up the chain, adding the starting value. If $a = 4$ (constant) and $v_0 = 2$, then

$$v = \int a \, dt = 4t + 2.$$

Slopes and areas on graphs

- The **slope** of an x - t graph is the velocity.
- The **area** under a v - t graph is the displacement.

A v - t graph holds a steady 5 m s^{-1} for 4 s: displacement = $5 \times 4 = 20 \text{ m}$ (the area).

2 Practice

Now apply the methods above.

2.1 A particle has $x(t) = 2t^3$. Find $v(t)$. [2]

2.2 A particle has $v(t) = 6t$. Find $a(t)$. [1]

2.3 A constant acceleration $a = 3 \text{ m s}^{-2}$ acts, with $v_0 = 5 \text{ m s}^{-1}$. Write $v(t)$. [2]

2.4 A v - t graph is a constant 8 m s^{-1} for 3 s. Find the displacement. [2]

3 Exam-style questions

3.1 The velocity of an object is the _____ of its position with respect to time. [1]

A	B
C	D

- A integral
B derivative
C reciprocal
D square

3.2 On a position-time graph, the velocity at a point is given by the [1]

- | | |
|---|---|
| A | B |
| C | D |
- A** area under the curve
B slope of the tangent
C height of the curve
D curvature

3.3 A particle moves with $x(t) = t^3 - 6t$.

(a) Find $v(t)$. [2]

(b) Find the acceleration $a(t)$. [1]

3.4 A particle starts from rest with acceleration $a(t) = 6t$.

(a) Find $v(t)$. [2]

(b) Find the velocity at $t = 2$ s. [1]