

3.4 Conservation of Energy

Name: _____ Class: _____ Date: _____

Total: 12 marks

KEY WORDS mechanical energy 机械能 • thermal energy 热能

• conservation of mechanical energy 机械能守恒 • friction 摩擦力 • potential energy 势能

Objective

Build the skills to answer exam questions on **conservation of energy**.

You must be able to:

- apply **conservation of mechanical energy** 机械能守恒 when no friction acts
- convert between kinetic and potential energy
- use $K_i + U_i = K_f + U_f$
- account for energy lost to friction as thermal energy

1 Worked examples

Study these first. Each one shows the method for a question type used later — follow the steps and you can do the Practice and Exam-style questions yourself.

■ Mechanical energy is conserved

With no friction, the total of kinetic plus potential energy stays constant:

$$K_i + U_i = K_f + U_f.$$

■ A falling object

Drop a 2 kg mass from 5 m ($g = 10 \text{ m s}^{-2}$). All the PE becomes KE:

$$mgh = \frac{1}{2}mv^2 \Rightarrow v = \sqrt{2gh} = \sqrt{2(10)(5)} = 10 \text{ m s}^{-1}.$$

■ When friction acts

Friction removes mechanical energy as heat:

$$K_i + U_i = K_f + U_f + (\text{energy lost to friction}).$$

2 Practice

Now apply the methods above.

2.1 State the condition under which mechanical energy is conserved. [1]

2.2 A ball is dropped from 2 m ($g = 10 \text{ m s}^{-2}$). Find its speed just before landing. [2]

2.3 At the top of a swing, a pendulum has 6 J of PE and no KE. State its KE at the bottom (no friction). [1]

2.4 Where does mechanical energy go when friction acts? [1]

3 Exam-style questions

3.1 In the absence of friction, as an object falls its [1]

A	B
C	D

- A** total mechanical energy increases
- B** total mechanical energy stays constant
- C** kinetic energy decreases
- D** potential energy increases

3.2 A cart at the top of a frictionless track has PE = 40 J and KE = 0. At the bottom its KE is [1]

A	B
C	D

- A** 0
- B** 20 J
- C** 40 J
- D** 80 J

3.3 A 0.5 kg ball is released from rest at height 1.8 m on a frictionless ramp ($g = 10 \text{ m s}^{-2}$).

(a) Find its speed at the bottom. [2]

(b) State the assumption you made. [1]

3.4 A 1 kg block slides down a ramp, starting with 20 J of PE, and reaches the bottom with 14 J of KE.

(a) State how much energy was lost to friction. [1]

(b) Explain where that energy went. [1]