

## 4.2 Change in Momentum and Impulse

Name: \_\_\_\_\_ Class: \_\_\_\_\_ Date: \_\_\_\_\_

Total: 16 marks

**KEY WORDS** impulse 冲量 • kinetic energy 动能

### Objective

Build the skills to answer exam questions on **change in momentum and impulse**.

**You must be able to:**

- calculate **impulse** 冲量:  $J = F \Delta t$
- use the impulse-momentum theorem:  $J = \Delta p$
- explain why a longer contact time reduces the force for a given change in momentum
- find force, time, or change in momentum given the other two

### 1 Worked examples

Study these first. Each one shows the method for a question type used later — follow the steps and you can do the Practice and Exam-style questions yourself.

#### ■ Impulse equals change in momentum

$$J = F \Delta t = \Delta p = m \Delta v.$$

A 50 N force acting for 0.2 s delivers  $J = 50 \times 0.2 = 10 \text{ kg m s}^{-1}$ , changing the momentum by  $10 \text{ kg m s}^{-1}$ .

#### ■ Longer time, smaller force

For a fixed change in momentum, a longer contact time means a smaller force:

$$F = \frac{\Delta p}{\Delta t}.$$

This is why crumple zones and airbags help – they stretch  $\Delta t$ , cutting the force.

#### ■ Finding the change in velocity

A 2 kg ball receives an impulse of  $12 \text{ kg m s}^{-1}$ :  $\Delta v = J/m = 12/2 = 6 \text{ m s}^{-1}$ .

## 2 Practice

---

Now apply the methods above.

**2.1** A 20 N force acts for 3 s. Find the impulse. [2]

---

---

**2.2** State what impulse equals (the theorem). [1]

---

**2.3** A 4 kg object receives an impulse of  $16 \text{ kg m s}^{-1}$ . Find its change in velocity. [2]

---

---

**2.4** Explain why a longer collision time reduces the force. [2]

## 3 Exam-style questions

---

**3.1** The impulse of a force equals the change in [1]

|   |   |
|---|---|
| A | B |
| C | D |

**A** kinetic energy

**B** momentum

**C** mass

**D** acceleration

**3.2** For a fixed change in momentum, increasing the contact time [1]

|   |   |
|---|---|
| A | B |
| C | D |

**A** increases the force

**B** decreases the force

**C** leaves the force unchanged

**D** reverses the force

**3.3** A 0.15 kg ball hits a wall at  $10 \text{ m s}^{-1}$  and bounces back at  $8 \text{ m s}^{-1}$  (opposite direction).

(a) Find the change in momentum. [2]

(b) If the contact lasts 0.02 s, find the average force. [2]

**3.4** A 60 kg person jumps down and lands, changing momentum by  $300 \text{ kg m s}^{-1}$ .

(a) Find the force if they stop in 0.1 s (stiff legs). [1]

(b) Find the force if they stop in 0.5 s (bent knees). [2]