

7.5 Simple and Physical Pendulums

Name: _____ Class: _____ Date: _____

Total: 13 marks**KEY WORDS** simple pendulum 单摆 • physical pendulum 物理摆 • period 周期 • amplitude 振幅

Objective

Build the skills to answer exam questions on **simple and physical pendulums**.**You must be able to:**

- use the **simple pendulum** 单摆 period: $T = 2\pi\sqrt{L/g}$
- use the **physical pendulum** 物理摆 period: $T = 2\pi\sqrt{I/(mgd)}$
- identify d as the pivot-to-centre-of-mass distance and I as the inertia about the pivot
- recognise the simple pendulum as a special case of the physical one

1 Worked examples

Study these first. Each one shows the method for a question type used later — follow the steps and you can do the Practice and Exam-style questions yourself.

■ The simple pendulum

A point mass on a light string of length L :

$$T = 2\pi\sqrt{\frac{L}{g}}.$$

■ The physical pendulum

Any rigid body swinging about a pivot:

$$T = 2\pi\sqrt{\frac{I}{mgd}},$$

where I is about the pivot and d reaches the centre of mass.

■ One is a special case of the other

For a point mass at distance L , $I = mL^2$ and $d = L$, so $T = 2\pi\sqrt{mL^2/(mgL)} = 2\pi\sqrt{L/g}$.

2 Practice

Now apply the methods above.

2.1 A simple pendulum has $L = 0.25$ m ($g = 9.8$ m s⁻²). Find its period. [2]**2.2** In $T = 2\pi\sqrt{I/(mgd)}$, what is d ? [1]**2.3** State whether making a pendulum's bob heavier changes its period. [1]**2.4** For a physical pendulum, about which axis must I be measured? [1]

3 Exam-style questions

3.1 The period of a simple pendulum increases if you [1]

A	B
C	D

A increase the mass**B** increase the length**C** increase the amplitude**D** paint it black**3.2** The simple-pendulum formula is a special case of the [1]

A	B
C	D

A mass-spring formula**B** physical-pendulum formula**C** ideal gas law**D** work-energy theorem**3.3** A uniform rod of length L is pivoted at one end, with $I = \frac{1}{3}mL^2$ and $d = L/2$.(a) Write its period in terms of L and g . [2]

(b) State whether it swings faster or slower than a simple pendulum of the same length. [1]

3.4 A pendulum has $L = 1.0$ m ($g = 9.8$ m s⁻²).

(a) Find its period. [2]

(b) State the effect of moving it to the Moon (smaller g). [1]