

AP Physics C: Mechanics

Scoring Guidelines

AP[®] PHYSICS 2017 SCORING GUIDELINES

General Notes About 2017 AP Physics Scoring Guidelines


1. The solutions contain the most common method of solving the free-response questions and the allocation of points for this solution. Some also contain a common alternate solution. Other methods of solution also receive appropriate credit for correct work.
2. The requirements that have been established for the paragraph length response in Physics 1 and Physics 2 can be found on AP Central at <https://secure-media.collegeboard.org/digitalServices/pdf/ap/paragraph-length-response.pdf>.
3. Generally, double penalty for errors is avoided. For example, if an incorrect answer to part (a) is correctly substituted into an otherwise correct solution to part (b), full credit will usually be awarded. One exception to this may be cases when the numerical answer to a later part should be easily recognized as wrong, e.g., a speed faster than the speed of light in vacuum.
4. Implicit statements of concepts normally receive credit. For example, if use of the equation expressing a particular concept is worth one point, and a student's solution embeds the application of that equation to the problem in other work, the point is still awarded. However, when students are asked to derive an expression it is normally expected that they will begin by writing one or more fundamental equations, such as those given on the exam equation sheet. For a description of the use of such terms as "derive" and "calculate" on the exams, and what is expected for each, see "The Free-Response Sections—Student Presentation" in the *AP Physics; Physics C: Mechanics, Physics C: Electricity and Magnetism Course Description* or "Terms Defined" in the *AP Physics 1: Algebra-Based and AP Physics 2: Algebra-Based Course and Exam Description*.
5. The scoring guidelines typically show numerical results using the value $g = 9.8 \text{ m/s}^2$, but use of 10 m/s^2 is of course also acceptable. Solutions usually show numerical answers using both values when they are significantly different.
6. Strict rules regarding significant digits are usually not applied to numerical answers. However, in some cases answers containing too many digits may be penalized. In general, two to four significant digits are acceptable. Numerical answers that differ from the published answer due to differences in rounding throughout the question typically receive full credit. Exceptions to these guidelines usually occur when rounding makes a difference in obtaining a reasonable answer. For example, suppose a solution requires subtracting two numbers that should have five significant figures and that differ starting with the fourth digit (e.g., 20.295 and 20.278). Rounding to three digits will lose the accuracy required to determine the difference in the numbers, and some credit may be lost.

SCORING GUIDELINES

Question 1

15 points total

(a) 3 points



For correctly drawing and labeling the vectors for the weight of the block and the tension for block M_1 with the tension larger than the weight

1 point

For correctly drawing and labeling the vectors for the weight of the block and the tension for block M_2 with the weight larger than the tension

1 point

For correctly drawing tension on the two blocks as equal in magnitude

1 point

Note: If any extraneous vectors are drawn, only a maximum of two points may be earned.

(b) 3 points

For correctly applying Newton's second law to block 1

1 point

$$F_T - M_1g = M_1a$$

For correctly applying Newton's second law to block 2

1 point

$$M_2g - F_T = M_2a$$

For combining the two equations above in such a way that will lead to the correct equation

1 point

$$M_2g - M_1g = (M_1 + M_2)a$$

$$a = \frac{(M_2 - M_1)}{(M_1 + M_2)}g$$

(c) 1 point

For indicating variables that will create a straight line whose slope can be used to determine g

1 point

Example: Vertical axis: a

$$\text{Horizontal axis: } \frac{(M_2 - M_1)}{(M_1 + M_2)}$$


Note: Full credit is earned if axes are reversed, or if the student uses another acceptable combination.

2017 SCORING GUIDELINES

Question 1 (continued)

Distribution of points

(d) 3 points



For using a correct scale that uses more than half the grid and for correctly labeling the axes, including units as appropriate

1 point

For correctly plotting the data

1 point

For drawing a straight best-fit line consistent with the plotted data

1 point

(e) 2 points

For correctly calculating the slope using the best-fit line and not the data points, unless the points fall on the best-fit line

1 point

(Note: Credit may be given for the linear regression only if the student states linear regression is used.)

$$\text{slope} = \frac{(y_2 - y_1)}{(x_2 - x_1)} = \frac{(3.0 - 1.0) \text{ m/s}^2}{(0.31 - 0.12)} = 10.5 \text{ m/s}^2$$

For correctly relating the slope to g

1 point

$$g = \text{slope} = 10.5 \text{ m/s}^2 \text{ (using linear regression yields } 10.1 \text{ m/s}^2 \text{)}$$

2017 SCORING GUIDELINES

Question 1 (continued)

Distribution
of points

(f) 2 points

Correct answer: “Lower”

For including a correct statement about the acceleration of the blocks
 For including a correct statement about the forces on the blocks

1 point
 1 point

Example: Because the block M_1 is on the table, the net force on the system increases, and therefore the acceleration increases. Because the acceleration of M_2 increases, the net force on M_2 must increase. Therefore, there must be a greater difference in the magnitude of the two forces on M_2 . Because the weight of block M_2 stays the same, the retarding force — the tension in the string — must decrease.

(g) 1 point

For a correct justification

1 point

Example: Block M_1 will experience friction with the table. The acceleration of the system will decrease and this will decrease the slope of the line; therefore, the value of g is determined by the experiment.

SCORING GUIDELINES

Question

15 points total

Distribution
of points

(a)

i. 2 points

For correctly using conservation of energy for the block moving down the incline

1 point

$$U_g = K$$

$$mgh = \frac{1}{2}mv^2$$

For a correct answer

1 point

$$v = \sqrt{2gh}$$

ii. 1 point

Correct answer: “Greater than”

For a correct justification

1 point

Example: The speed is proportional to the square root of the change in height. So if the height is reduced by a factor of 2, the speed is reduced by a factor of $\sqrt{2} \approx 1.41$. Therefore, the speed halfway down the ramp is more than half the speed at the bottom of the ramp.

Note: If the incorrect selection is made, the justification cannot earn credit.

(b) 1 point

For correctly using conservation of energy, consistent with part (a), for the block compressing the spring

1 point

$$U_g = U_s$$

$$mgh = \frac{1}{2}kx_{\max}^2$$

$$x_{\max} = \sqrt{\frac{2mgh}{k}}$$

(c) 2 points

For indicating a simple harmonic motion approach

1 point

For a correct answer

1 point

$$t = \frac{1}{4}T = \left(\frac{1}{4}\right)\left(2\pi\sqrt{\frac{m}{k}}\right) = \frac{\pi}{2}\sqrt{\frac{m}{k}}$$

2017 SCORING GUIDELINES

Question continued)

Distribution
of points

(d)

i. 3 points

For correctly applying Newton's second law for the horizontally sliding block

For correctly indicating that the direction of F_{net} is opposite the direction of motion

$$F_{net} = ma$$

$$-\beta v^2 = ma$$

For expressing the equation as a differential equation

$$-\beta v^2 = m \frac{dv}{dt}$$

1 point

1 point

1 point

ii. 3 points

For correctly separating variables

$$-\frac{\beta}{m} dt = \frac{1}{v^2} dv$$

For correctly integrating the equation above

$$\int -\frac{\beta}{m} dt = \int \frac{1}{v^2} dv$$

$$-\frac{\beta}{m} [t] = \left[-\frac{1}{v} \right]$$

For using the correct limits or constant of integration

$$-\frac{\beta}{m} [t]_0^t = \left[-\frac{1}{v} \right]_{v_0}^{v(t)}$$

$$-\frac{\beta t}{m} = -\frac{1}{v(t)} - \left(-\frac{1}{v_0} \right)$$

$$\frac{1}{v(t)} = \frac{1}{v_0} + \frac{\beta t}{m}$$

1 point

1 point

1 point


2017 SCORING GUIDELINES

Question continued)

Distribution
of points

(e)

3 points



For a displacement curve that is concave down and approaching a horizontal asymptote

For a velocity curve that is concave up and has the horizontal axis as an asymptote

For an acceleration curve that is concave down and has the horizontal axis as an asymptote

1 point

1 point

1 point

Note: If an incorrect nonlinear velocity graph is generated, 1 point is earned if the position and acceleration graphs are consistent with the velocity graph.

Note: Full credit is earned if all three graphs are flipped vertically.

2017 SCORING GUIDELINES

Question 3

15 points total

Distribution
of points

(a) 2 points

For correctly applying conservation of energy to the cylinder rolling down the incline

$$U_{g_top} = K_{table}$$

$$K_{table} = mgh = (0.50 \text{ kg})(9.8 \text{ m/s}^2)(1.0 \text{ m})(\sin 30^\circ)$$

For a correct answer with units

$$K_{table} = 2.45 \text{ J (or } 2.5 \text{ J using } g = 10 \text{ m/s}^2)$$

1 point

1 point

(b) 3 points

For correctly setting the kinetic energy of the cylinder equal to the sum of both the linear and rotational kinetic energy

$$K = \frac{1}{2}mv^2 + \frac{1}{2}I\omega^2$$

For correctly substituting into the above equation for the linear velocity and moment of inertia of the cylinder

$$K = \frac{1}{2}MR^2\omega^2 + \frac{1}{2}\left(\frac{1}{2}MR^2\right)\omega^2 = \frac{1}{2}M(R\omega)^2 + \frac{1}{4}M(R\omega)^2 = \frac{3}{4}M(R\omega)^2$$

$$\omega = \sqrt{\frac{4K}{3MR^2}} = \sqrt{\frac{(4)(2.45 \text{ J})}{(3)(0.50 \text{ kg})(0.10 \text{ m})^2}}$$

For correct substitution into the equation above

$$\omega = 25.6 \text{ rad/s (or } 26.0 \text{ rad/s using } g = 10 \text{ m/s}^2)$$

1 point

1 point

1 point

(c) 2 points

For using a correct expression for the ratio of the rotational kinetic energy to the total kinetic energy of the cylinder

$$\frac{K_{rot}}{K_{tot}} = \frac{\frac{1}{2}I\omega^2}{\frac{1}{2}mv^2 + \frac{1}{2}I\omega^2 + mgh} = \frac{\frac{1}{4}M(R\omega)^2}{\frac{3}{4}M(R\omega)^2 + Mgh_{table}} = \frac{(R\omega)^2}{3(R\omega)^2 + 4gh_{table}}$$

1 point

For substituting into the above equation

$$\frac{K_{rot}}{K_{tot}} = \frac{[(0.10 \text{ m})(25.6 \text{ rad/s})]^2}{(3)(0.10 \text{ m})^2(25.6 \text{ rad/s})^2 + (4)(9.81 \text{ m/s}^2)(0.75 \text{ m})}$$

1 point

$$\frac{K_{rot}}{K_{tot}} = 0.133 \text{ (or } 0.135 \text{ using } g = 10 \text{ m/s}^2)$$

2017 SCORING GUIDELINE

Question 3 (continued)

Distribution
of points

(c) (continued)

Alternate Solution

For using a correct expression for the ratio of the rotational kinetic energy to the total potential energy of the cylinder

Alternate Points

1 point

$$\frac{K_{rot}}{K_{tot}} = \frac{\frac{1}{2}I\omega^2}{U_{total}} = \frac{\frac{1}{4}M(R\omega)^2}{Mg(h+y)}$$

For substituting into the above equation

1 point

$$\frac{K_{rot}}{U_{total}} = \frac{\frac{1}{4}(R\omega)^2}{g(h+y)} = \frac{\frac{1}{4}[(0.10 \text{ m})(25.6 \text{ rad/s})]^2}{(9.81 \text{ m/s}^2)((1.0 \text{ m})\sin(30^\circ) + (0.75 \text{ m}))}$$

$$\frac{K_{rot}}{U_{total}} = 0.13 \text{ (or } 0.135 \text{ using } g = 10 \text{ m/s}^2)$$

(d) 2 points

For correctly using motion in the vertical direction to calculate the time for the cylinder to reach the floor

1 point

$$y = y_0 + v_{0y}t + \frac{1}{2}a_yt^2$$

$$y - y_0 = 0 - \frac{1}{2}gt^2$$

Determine the time for the cylinder to reach the floor

$$y - y_0 = -\frac{1}{2}gt^2$$

$$0 - y = -\frac{1}{2}gt^2$$

$$t = \sqrt{\frac{2y}{g}} = \sqrt{\frac{(2)(0.75 \text{ m})}{(9.81 \text{ m/s}^2)}} = 0.39 \text{ s}$$

For correctly using the equation for constant speed in the horizontal direction

1 point

$$x - x_0 = v_{0x}t$$

$$x - x_0 = R\omega t$$

$$x - x_0 = R\omega\sqrt{\frac{2y}{g}}$$

$$D = R\omega t = (0.10 \text{ m})(25.6 \text{ rad/s})(0.39 \text{ s})$$

$$D = 1.0 \text{ m}$$

2017 SCORING GUIDELINE**Question 3 (continued)****Distribution
of points**

(e)

i. 2 points

For selecting “Equal to” and attempting a relevant justification
 For a correct justification

1 point

1 point

Example: Because the sphere falls the same height as the cylinder and because they have the same mass, the sphere-Earth system has the same initial potential energy and, therefore, the same total kinetic energy when it reaches the floor.

ii. 2 points

For selecting “Less than” and attempting a relevant justification
 For a correct justification

1 point

1 point

Example: Because the rotational inertia of the sphere is less than the rotational inertia of the cylinder, the sphere will rotate faster and, because $v = r\omega$, will move with a greater linear speed. Because the mass is the same and the linear speed is greater, the sphere will have a greater linear kinetic energy. Because the total kinetic energies of the sphere and cylinder are the same, the sphere must have less rotational kinetic energy.

iii. 2 points

For selecting “Greater than” and attempting a relevant justification
 For a correct justification

1 point

1 point

Example: Because the sphere has a greater linear speed as it leaves the table, it will travel a greater horizontal distance before it reaches the floor.

AP[®] Physics C: Mechanics

2016 Scoring Guidelines

© 2016 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.


Visit the College Board on the Web: www.collegeboard.org.

AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

2016 SCORING GUIDELINES

15 points total

3 points




For correctly drawing and labeling the force of tension
 For correctly drawing and labeling the force of friction
 For correctly drawing and labeling both forces in the vertical direction
 Note: A maximum of two points may be earned if there are any extraneous vectors.

Distribution
of points

1 point
 1 point
 1 point

3 points



For a correct scale that uses more than half the grid
 For correctly plotting the given data
 For drawing a straight line consistent with the given data
 Note: Full credit can be earned if the axes are switched.

1 point
 1 point
 1 point

SCORING GUIDELINES

Question 1 (continued)

Distribution
of points

2 points

For correctly calculating slope using the best-fit straight line and not data points

1 point

$$\text{slope} = \frac{(y_2 - y_1)}{(x_2 - x_1)} = \frac{(0.70 - 0.16)}{(0.60 - 0.35)} \text{ kg}^{-1} = 2.16 \text{ kg}^{-1}$$

Note: Linear regression gives slope = 2.12 kg^{-1} .

For correctly calculating the mass of the cart using the slope

1 point

$$m = \frac{1}{\text{slope}} = \frac{1}{(2.16 \text{ kg}^{-1})}$$

Correct answer:

$$m = 0.463 \text{ kg (Note: linear regression gives } m = 0.472 \text{ kg)}$$

1 point

For an answer with correct units consistent with the x-intercept of the graph from

1 point

(b) i.

$$f = 0.272 \text{ N}$$

1 point

Applying Newton's second law and substituting the values from part (b)

$$\sum F = ma = F_a - f$$

$$a = \frac{F_a - f}{m} = \frac{0.45 \text{ N} - 0.272 \text{ N}}{0.463 \text{ kg}}$$

For an answer with correct units consistent with part (b), either from the graph or calculated using the mass and frictional force.

1 point

$$a = 0.376 \text{ m/s}^2$$

SCORING GUIDELINES

Question 1 (continued)

Distribution
of points

3 points

For using a correct equation to solve for the speed of the cart when the string breaks, with an acceleration consistent with part (c) i.

$$v_2 = v_1 + at$$

$$v_2 = 0 + (0.376 \text{ m/s}^2)(2.0 \text{ s})$$

$$v_2 = 0.752 \text{ m/s}$$

For recognizing that the acceleration of the cart after the string breaks is due to the frictional force determined in part (b)

Use a correct equation to solve for the time for the cart to stop after the string breaks

$$v_2 = v_1 + at$$

$$0 = v_1 - \frac{f}{m}t$$

For using the final velocity before the string breaks as the initial velocity for the cart stopping in the correct equation for time

$$t = \frac{mv_1}{f}$$

$$t = \frac{(0.463 \text{ kg})(0.752 \text{ m/s})}{(0.272 \text{ N})}$$

Correct answer

$$t = 1.28 \text{ s}$$

Alternate solution

Alternate points

For setting the magnitude of the impulse before the string breaks equal to the magnitude of the impulse after the string breaks

$$F_1 t_1 = F_2 t_2$$

For correctly using the proper force (e.g., the tension force minus the friction force) for F_1

For correctly using the proper force (e.g., the friction force) for F_2

Correct answer

$$t = 1.28 \text{ s}$$

1 point

For selecting “Equal to”

1 point

For selecting “Greater than”

1 point

1 point

SCORING GUIDELINES

Question

Distribution
of points

15 points total

2 points

Block of Mass $2M$ F_{N1}  $2Mg$ Block of Mass $3M$ F_{N2}  $3Mg$

For correctly drawing and labeling vectors on the block of mass $2M$

1 point

For correctly drawing and labeling vectors on the block of mass $3M$ using symbols for the vectors that are physically correct and different from those on the block of mass $2M$

1 point

Note: A maximum of one point can be earned if there are any extraneous vectors.

2 points

Using a proper expression for conservation of momentum

$$p_1 = p_2$$

For correctly substituting into the above equation

1 point

$$3Mv_0 = (3M + 2M)v_f$$

For a correct answer

1 point

$$v_f = \frac{3}{5}v_0$$

(1 point

Using a proper expression for kinetic energy of the two-block system

$$K = \frac{1}{2}mv^2$$

$$K = \frac{1}{2}(5M)\left(\frac{3}{5}v_0\right)^2$$

For an answer consistent with part (b)

1 point

$$K = \frac{9}{10}Mv_0^2$$

SCORING GUIDELINES

Question continued)	Distribution of points
4 points	
For a correct expression of the conservation of energy $\Delta K_{\text{system}} + \Delta U_{\text{system}} = 0$	1 point
$K_0 = U_{\text{final}}$ For attempting to integrate the spring force equation $\frac{9}{10} M v_0^2 = - \int_{x_1}^{x_2} B x^3 dx$ $\frac{9}{10} M v_0^2 = \int_{x_1}^{x_2} B x^3 dx$	1 point
For using the correct limits of integration or an appropriate constant of integration $\frac{9}{10} M v_0^2 = \int_0^D B x^3 dx$ $\frac{9}{10} M v_0^2 = \left[\frac{B x^4}{4} \right]_0^D$	1 point
For an answer consistent with the speed from (b) or the kinetic energy from part (c) $D = \sqrt[4]{\frac{18 M v_0^2}{5 B}}$	1 point
2 points	
For selecting the correct answer “Right” with a reasonable attempt at a justification If the incorrect selection is made, no points are earned for the justification.	1 point
For an indication that at maximum compression the block of mass $2M$ has an acceleration to the right due to the forces acting on the block of mass $2M$ or an acceleration to the right due to the external spring force acting on the system of blocks Example: At maximum compression the two-block system is instantaneously at rest. The only horizontal external force acting on the system is due to the spring. This force is directed to the right. The system and therefore the block of mass $2M$ is accelerated to the right, which implies that the net force acting on the block of mass $2M$ is also to the right.	1 point


SCORING GUIDELINES

Question continued)	Distribution of points
2 points	
a magnitude of the net force is greater on the block of mass $3M$. If the incorrect selection is made, no points are earned for the justification. For an indication that both blocks will have the same acceleration For a correct justification for why the net force is greater on the block of mass $3M$ Example: Because the blocks stick together, both blocks must have the same acceleration. Because the block of mass $3M$ has more mass, the net force on it must be greater than the net force on the block of mass $2M$.	1 point 1 point
2 points	
For selecting the correct answer “No,” with a reasonable attempt at a justification If the incorrect selection is made, no points are earned for the justification. If the correct answer is selected without any justification, the point is not earned for the selection.	1 point
For an indication that the spring does not apply a linear force and that simple harmonic motion is the resulting motion of a linear restoring force $\left(a = -\frac{k}{m} \Delta x \right)$ Example: Because the blocks are sticking together and are attached to the spring, the spring will apply a restoring force to the blocks. However, because the restoring force exerted by a nonlinear spring is not proportional to the blocks’ displacement from equilibrium, the blocks do not exhibit simple harmonic motion.	1 point

2016 SCORING GUIDELINES

15 points total	Question	Distribution of points
3 points	For an indication that the spring force equals the mass times the centripetal acceleration $F_S = F_C = ma_C$ $kx = mr\omega^2$ correct substitution for x in the equation For a correct substitution for r in the equation that can be used to solve for k $k\left(\frac{d}{2}\right) = m(2d)\omega^2$ Correct answer: $k = 4m\omega^2$	1 point 1 point 1 point
1 point	Substitute values for the block into the equation for rotational inertia $I = \sum mr^2 = m(2d)^2$ For a correct answer or an answer consistent with the radius used in part (a) $I = 4md^2$	1 point
2 points	For an indication that the rotational inertia of the system must include the platform, the rod, and the object $I = I_P + I_R + I_O$ $I = \frac{m_P R^2}{2} + \frac{m_R d^2}{3} + mr^2$ For correctly substituting into the equation the rotational inertia of the platform, the rod, and the object consistent with (b) i. $I = \frac{5m(2d)^2}{2} + \frac{3m d^2}{3} + 4md^2$ $I = (10 + 1 + 4)md^2$ Correct answer: $I = 15md^2$	1 point 1 point
1 point	Using a correct expression of angular momentum $L = I\omega$ For an answer consistent with the answer from part (b) ii. $L = 15md^2\omega$	1 point

SCORING GUIDELINES

Question continued)	Distribution of points
3 points For indicating that the spring force equals the mass times the centripetal acceleration $F_S = F_C = ma_C$ $kx = mr\omega^2$ For a correct substitution for k , or a substitution consistent with part (a), in the equation For a correct substitution for r in the equation that can be used to solve for x $(4m\omega^2)x = m(d+x)\omega^2$ $4x = d+x$ $3x = d$ Correct answer: $x = d/3$	1 point 1 point 1 point
2 points electing "Decreasing" the wrong selection is made, no points are earned for the justification. For a correct justification Example: The rotational inertia I of the system decreases while the angular velocity ω stays the same. Because the angular momentum is $I\omega$, it decreases.	1 point 1 point
2 points For selecting "Negative" If the wrong selection is made, no points are earned for the justification. For a correct justification Example: Because the angular velocity is constant and the rotational inertia has decreased, the rotational kinetic energy has decreased. Therefore, work must be negative.	1 point 1 point
1 point 	1 point

n arrow starting on the box and pointing down and to the right