

Linear Momentum

AP Physics C: Mechanics

Linear Momentum

Linear momentum 动量 is mass times velocity:

$$\vec{p} = m\vec{v}.$$

It is a **vector** 矢量 pointing along the velocity, and it is *the* quantity for analysing **collisions** 碰撞 and **explosions** 爆炸. A collection of objects can be treated as one system moving with the velocity of its **center of mass** 质心:

$$\vec{v}_{\text{cm}} = \frac{\sum m_i \vec{v}_i}{\sum m_i},$$

so the system's total momentum is its total mass times \vec{v}_{cm} –one object's worth of book-keeping for any number of parts.

Change in Momentum and Impulse

Newton's second law is really a statement about momentum:


$$\vec{F}_{\text{net}} = \frac{d\vec{p}}{dt},$$

which reduces to $m\vec{a}$ when the mass is constant –and handles a *changing* mass ($\vec{F} = \frac{dm}{dt}\vec{v}$, a rocket or a chain piling onto a scale) when it is not. The **impulse** 冲量 delivered by a force is its time integral, and the **impulse–momentum theorem** 冲量-动量定理 says it equals the change in momentum:

$$\vec{J} = \int_{t_1}^{t_2} \vec{F}_{\text{net}} dt = \Delta\vec{p}.$$

Impulse is a vector along the net force. Read it off graphs both ways:

- On a force–time graph, impulse is the **area** under the curve.
- On a momentum–time graph, the net force is the **slope** 斜率 at each instant.



Impulse is the area under the force-time curve, equal to the average force times the contact time

Spreading the same Δp over a longer time lowers the force –that is why airbags, crumple zones, and soft landings work, and why you bend your knees when you land.


Worked example. A time-varying force $F(t) = 10t$ N acts on a 2.0 kg object for 2.0 s. The impulse is $J = \int_0^2 10t \, dt = [5t^2]_0^2 = 20 \text{ N} \cdot \text{s}$, so the speed changes by $\Delta v = J/m = 10 \text{ m/s}$.

Conservation of Linear Momentum

Internal forces come in Newton’s-third-law pairs, so they cancel inside any system: they can shuffle momentum *between* parts but never change the total. Momentum only enters or leaves a system through a net *external* force ($\vec{J} = \Delta\vec{p}$). So, with **zero net external force**, total momentum is **conserved** 守恒:

$$\sum \vec{p}_{\text{before}} = \sum \vec{p}_{\text{after}}.$$

Momentum is conserved in *every* collision, however violent –choose the system large enough that the collision forces are internal. Apply conservation separately along each axis; AP asks for quantitative work in one or two dimensions.



A head-on collision: total momentum before equals total momentum after

Worked example (explosion). A 6.0 kg shell at rest splits into a 2.0 kg piece moving at 9.0 m/s east and a 4.0 kg piece. Total momentum stays zero, so the heavy piece moves west at $v = \frac{2.0(9.0)}{4.0} = 4.5$ m/s. The kinetic energy came from stored (chemical or spring) energy – momentum conservation does not require kinetic-energy conservation.

Elastic and Inelastic Collisions


All collisions conserve momentum; they differ in what happens to the **kinetic energy** 动能:

type	momentum	kinetic energy
elastic collision 弹性碰撞	conserved	total conserved (individual shares may change)
inelastic collision 非弹性碰撞	conserved	decreases –some becomes heat, sound, deformation 形变
perfectly inelastic collision 完全非弹性碰撞	conserved	largest possible loss –the objects stick and share one velocity

Strategy: *always* write momentum conservation first; add the kinetic-energy equation only when the problem says "elastic". Two useful elastic facts: equal masses in a 1D elastic collision simply exchange velocities, and in the center-of-mass frame each object just reverses its velocity.

Worked example. A 1000 kg car at 20 m/s strikes a stationary 1500 kg car and they lock together –perfectly inelastic. Momentum: $v = \frac{1000(20)}{2500} = 8.0$ m/s. Kinetic energy falls from 2.0×10^5 J to $\frac{1}{2}(2500)(8.0)^2 = 8.0 \times 10^4$ J: about 60% is lost, even though momentum is exactly conserved.

Worked example (2D). A puck moving east at 4.0 m/s strikes an identical puck at rest; after the glancing hit, one moves at 2.0 m/s at 60° north of east. Conserve each axis: east–west, $m(4.0) = m(2.0) \cos 60^\circ + mv_x$, so $v_x = 3.0$ m/s; north–south, $0 = m(2.0) \sin 60^\circ - mv_y$, so $v_y = 1.7$ m/s. The second puck moves at $\sqrt{3.0^2 + 1.7^2} = 3.5$ m/s, about 30° south of east.



A glancing collision, resolved along two perpendicular axes

Exam skill. On FRQs, justify with the *condition*, not the slogan: "the net external force on the two-puck system is zero during the collision, so its total momentum is constant." If asked whether the collision is elastic, *compute* the kinetic energy before and after and compare –never assume.

Exam tips

- Impulse equals the momentum change: $\vec{J} = \int \vec{F} dt = \Delta\vec{p}$, and it is the area under a force–time graph.
- **Momentum is conserved** whenever the net external force is zero —always the go-to for collisions.
- Distinguish **elastic** (kinetic energy conserved) from **inelastic** (objects stick) collisions.
- Apply conservation to each component (x and y) separately in 2-D.
- Connect to center of mass: total momentum = $M\vec{v}_{cm}$.