

4. Directions:

- Unless otherwise specified, the domain of a function f is assumed to be the set of all real numbers x for which $f(x)$ is a real number. Angle measures for trigonometric functions are assumed to be in radians.
- Solutions to equations must be real numbers. Determine the exact value of any expression that can be obtained without a calculator. For example, $\log_2 8$, $\cos\left(\frac{\pi}{2}\right)$, and $\sin^{-1}(1)$ can be evaluated without a calculator.
- Unless otherwise specified, combine terms using algebraic methods and rules for exponents and logarithms, where applicable. For example, $2x + 3x$, $5^2 \cdot 5^3$, $\frac{x^5}{x^2}$, and $\ln 3 + \ln 5$ should be rewritten in equivalent forms.
- For each part of the question, show the work that leads to your answers.

A. The functions g and h are given by

$$g(x) = 2 \log_3 x$$

$$h(x) = 4 \cos^2 x$$

- Solve $g(x) = 4$ for values of x in the domain of g .
- Solve $h(x) = 3$ for values of x in the interval $\left[0, \frac{\pi}{2}\right)$.

B. The functions j and k are given by

$$j(x) = \log_2 x + 3 \log_2 2$$

$$k(x) = \frac{6}{\tan x (\csc^2 x - 1)}$$

- Rewrite $j(x)$ as a single logarithm base 2 without negative exponents in any part of the expression. Your result should be of the form $\log_2(\text{expression})$.
- Rewrite $k(x)$ as an expression in which $\tan x$ appears exactly once and no other trigonometric functions are involved.

C. The function m is given by $m(x) = e^{2x} - e^x - 12$. Find all input values in the domain of m that yield an output value of 0.

STOP

END OF EXAM

4. Directions:

- Unless otherwise specified, the domain of a function f is assumed to be the set of all real numbers x for which $f(x)$ is a real number. Angle measures for trigonometric functions are assumed to be in radians.
- Solutions to equations must be real numbers. Determine the exact value of any expression that can be obtained without a calculator. For example, $\log_2 8$, $\cos\left(\frac{\pi}{2}\right)$, and $\sin^{-1}(1)$ can be evaluated without a calculator.
- Unless otherwise specified, combine terms using algebraic methods and rules for exponents and logarithms, where applicable. For example, $2x + 3x$, $5^2 \cdot 5^3$, $\frac{x^5}{x^2}$, and $\ln 3 + \ln 5$ should be rewritten in equivalent forms.
- For each part of the question, show the work that leads to your answers.

(A) The functions g and h are given by

$$g(x) = e^{(x+3)}$$

$$h(x) = \arcsin\left(\frac{x}{2}\right).$$

(i) Solve $g(x) = 10$ for values of x in the domain of g .

(ii) Solve $h(x) = \frac{\pi}{4}$ for values of x in the domain of h .

(B) The functions j and k are given by

$$j(x) = \log_{10}(8x^5) + \log_{10}(2x^2) - 9\log_{10}x$$

$$k(x) = \left(\frac{1 - \sin^2 x}{\sin x} \right) \sec x.$$

(i) Rewrite $j(x)$ as a single logarithm base 10 without negative exponents in any part of the expression. Your result should be of the form $\log_{10}(\text{expression})$.

(ii) Rewrite $k(x)$ as a single term involving $\tan x$.

(C) The function m is given by

$$m(x) = \cos^{-1}(\tan(2x)).$$

Find all values in the domain of m that yield an output value of 0.

Write your responses to this question only on the designated pages in the separate Free Response booklet. Write your solution to each part in the space provided for that part.

STOP

END OF EXAM