

1.7 Summary Statistics for a Quantitative Variable

Name: _____ Class: _____ Date: _____ **Total: 11 marks**

KEY WORDS mean 均值 • median 中位数 • standard deviation 标准差 • resistant 稳健
• interquartile range 四分位距

Objective

Build the skills to answer exam questions on **summary statistics for a quantitative variable**.

You must be able to:

- calculate the **mean** 均值 $\bar{x}$ and the **median** 中位数
- calculate the **range**, **interquartile range** 四分位距 $IQR = Q_3 - Q_1$, and **standard deviation** 标准差 s_x
- explain why the median and IQR are **resistant** 稳健 to outliers

1 Worked examples

Study these first. Each one shows the method for a question type used later.

■ Measures of center

- Mean** $\bar{x} = \frac{\sum x}{n}$.
- Median** — the middle value when the data are ordered.

■ Measures of spread

- Range** = max – min.
- IQR** = $Q_3 - Q_1$.
- Standard deviation** s_x — the typical distance from the mean.

■ Resistance

The **median** and **IQR** are **resistant** to outliers; the **mean** and **standard deviation** are **not**.

2 Practice

2.1 Find the mean of {3, 5, 7, 9}. [2]

2.2 Find the median of {2, 6, 4, 8, 10}. [2]

2.3 State which measures of center and spread resist outliers. [1]

3 Exam-style questions

3.1 The interquartile range equals [1]

- A

B

C

D

A max – min

B $Q_3 - Q_1$

C mean – median

D $Q_1 - Q_3$

3.2 Which of these is resistant to outliers? [1]

- A

B

C

D

A the mean

B the standard deviation

C the median

D the range

3.3 For the data {4, 8, 6, 10, 12}:

- (a) find the mean. [2]

(b) find the median. [1]

(c) find the range. [1]