

6.11 Carrying Out a Test for the Difference of Two Population Proportions

Name: _____ Class: _____ Date: _____

Total: 9 marks

KEY WORDS significance level 显著性水平 • p-value P 值 • test statistic 检验统计量

Objective

Build the skills to answer exam questions on **carrying out a test for a difference of two proportions**.

You must be able to:

- compute the test statistic z using the pooled standard error
- find the **p-value** and compare it to α
- state a conclusion in context

1 Worked examples

Study these first. Each one shows the method for a question type used later.

■ The full procedure

1. State $H_0 : p_1 = p_2$ and H_a ; check conditions.
2. Compute $z = \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}_c(1 - \hat{p}_c) \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}}$.
3. Find the **p-value** from z .
4. Compare to α : reject H_0 if p-value $< \alpha$, else fail to reject.
5. State the conclusion **in context**.

■ A worked test

$\hat{p}_1 = 0.60$ ($n_1 = 50$), $\hat{p}_2 = 0.45$ ($n_2 = 60$). Pooled $\hat{p}_c = \frac{30+27}{110} = 0.518$. Then

$$z = \frac{0.60 - 0.45}{\sqrt{0.518(0.482) \left(\frac{1}{50} + \frac{1}{60} \right)}} \approx \frac{0.15}{0.096} \approx 1.57,$$

a two-sided p-value $\approx 0.12 > 0.05$ – fail to reject H_0 .

2 Practice

2.1 State the steps to carry out a two-proportion test. [2]

2.2 A two-proportion test gives $z = 2.5$. State whether this is likely significant at $\alpha = 0.05$. [1]

2.3 State what to compare the p-value to. [1]

3 Exam-style questions

3.1 After computing z , you find the [1]

A	B
C	D

- A slope
- B p-value
- C mean
- D residual

3.2 A two-proportion test rejects H_0 when the p-value is [1]

A	B
C	D

- A above α
- B below α
- C exactly zero
- D exactly one

3.3 A two-proportion test gives a p-value of 0.03 at $\alpha = 0.05$.

(a) State the decision. [1]

(b) State whether the two proportions differ significantly. [1]

(c) State the conclusion in context. [1]