

7.3 Justifying a Claim About a Population Mean Based on a Confidence Interval

Name: _____ Class: _____ Date: _____

Total: 8 marks

Objective

Build the skills to answer exam questions on **justifying a claim about a mean from a confidence interval**.

You must be able to:

- use a confidence interval to judge whether a claimed mean is **plausible**
- recognise a value inside the interval as plausible and one outside as not supported

1 Worked examples

Study these first. Each one shows the method for a question type used later.

■ Judging a claim about μ

The confidence interval gives the plausible values of the population mean:

- a claimed value **inside** the interval is **plausible**;
- a claimed value **outside** the interval is **not supported** by the data.

■ Example

A 95% interval of (48, 52) contains 50, so " $\mu = 50$ " is plausible; it does not contain 55, so " $\mu = 55$ " is not supported.

2 Practice

2.1 State how to judge a claim about μ using a confidence interval. [1]

2.2 A 95% CI for μ is (48, 52). State whether the claim " $\mu = 50$ " is plausible. [1]

2.3 State what a claimed μ outside the interval implies. [1]

3 Exam-style questions

3.1 A claimed mean that lies inside the confidence interval is [1]

A	B
C	D

- A** rejected
- B** plausible
- C** impossible
- D** certainly correct

3.2 A 95% CI for μ is $(60, 70)$. The claim " $\mu = 55$ " is [1]

A	B
C	D

- A** plausible
- B** not supported
- C** certainly true
- D** unknown

3.3 A 95% CI for the mean is $(10.2, 11.8)$.

- (a) State whether " $\mu = 11$ " is plausible. [1]
- (b) State whether " $\mu = 12$ " is plausible. [1]
- (c) State the confidence level. [1]