

AP[®] STATISTICS
2017 SCORING GUIDELINES

Question 2

Intent of Question

The primary goals of this question were to assess a student's ability to (1) construct and interpret a confidence interval for a population proportion and (2) use a confidence interval for a proportion to find a confidence interval for a dollar amount that can be calculated using that proportion.

Solution

Part (a):

Step 1: Identify the appropriate confidence interval by name or formula and check appropriate conditions.

The appropriate procedure is a one-sample z-interval for a population proportion p . In this case, the population is all customers of the restaurant who ask for a water cup, and p is the proportion of that population who will fill the cup with a soft drink. The appropriate formula is $\hat{p} \pm z^* \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$.

Conditions:

1. Random sample
2. Large sample (number of successes $n\hat{p} \geq 10$ and number of failures $n(1-\hat{p}) \geq 10$)

For condition 1 the stem of the problem states that a random sample of customers who asked for a water cup was used.

For condition 2 the number of successes (filled cup with soft drink) is 23 and the number of failures is 57, both of which are greater than 10.

Step 2: Correct mechanics

The sample proportion is $\hat{p} = \frac{23}{80} = 0.2875$. The confidence interval is

$$\begin{aligned} 0.2875 \pm 1.96 \sqrt{\frac{0.2875(1-0.2875)}{80}} \\ = 0.2875 \pm 1.96(0.0506) \quad \text{or } 0.1883 \text{ to } 0.3867. \\ = 0.2875 \pm 0.0992 \end{aligned}$$

Step 3: Interpretation

We can be 95 percent confident that in the population of all customers of the restaurant who ask for a water cup, the proportion who will fill it with a soft drink is between 0.1883 and 0.3867.

Part (b):

Using the confidence interval in part (a), a 95 percent interval estimate for the number of customers in June who asked for a water cup but then filled it with a soft drink is $3,000 \times 0.1883$ to $3,000 \times 0.3857$, or 565 to 1,160. At a cost of \$0.25 per customer, a 95 percent interval estimate for the cost to the restaurant in June is \$141.25 to \$290.00.

2017 SCORING GUIDELINES

Question 2 (continued)

Scoring

This question is scored in four sections. Section 1 consists of step 1 in part (a), section 2 consists of step 2 in part (a), section 3 consists of step 3 in part (a), and section 4 consists of part (b). Each section is scored as essentially correct (E), partially correct (P), or incorrect (I).

Section 1 is scored as follows:

Essentially correct (E) if the one-sample z-interval for a proportion is identified (either by name or formula) *AND* both conditions (random sampling and large sample) are adequately addressed.

Partially correct (P) if the response identifies the correct procedure *BUT* adequately addresses only one of the two required conditions;

OR

if the response does not identify the correct procedure *BUT* adequately addresses both required conditions.

Incorrect (I) if the response does not meet the criteria for E or P.

Notes:

- Stating the large sample condition without verifying it is not sufficient. The response must use specific numerical values to adequately address the condition.
- If the response includes additional inappropriate conditions, such as $n \geq 30$ or requiring a normal population, then the response earns at most P for section 1.
- Stating and checking a condition about the size of the sample relative to the size of the population is appropriate but not required.
- Any statement of hypotheses, description of the population, or definition of the parameter should be considered extraneous. However, if such statements are included and incorrect, they are considered as poor communication in terms of holistic scoring.

Section 2 is scored as follows:

Essentially correct (E) if the response gives the correct 95 percent confidence interval. Supporting work is not required, but if included, it must be correct.

Partially correct (P) if the response gives a correct confidence interval with incorrect (but appropriate) supporting work shown;

OR

if the response gives an incorrect but reasonable confidence interval with appropriate supporting work shown — for instance, if a value other than 1.96 is used for the critical value or a value other than $\frac{23}{80}$ is used for $\hat{p}$.

Incorrect (I) if the response does not meet the criteria for E or P.

Notes:

- Appropriate supporting working must have the form:
 $\text{proportion} \pm (\text{critical value})(\text{SE/SD of proportion}).$

2017 SCORING GUIDELINES

Question 2 (continued)

- A confidence interval that has both endpoints outside the interval from 0 to 1 is considered unreasonable.

Section 3 is scored as follows:

Essentially correct (E) if the response provides an appropriate interpretation of the interval that includes the following three components:

1. Conveys inference about a population proportion
2. Demonstrates a clear understanding that the parameter is the proportion of the water cup population that fills the cup with a soft drink
3. Mentions 95 percent confidence and interprets it correctly using words such as “We can be 95 percent confident” or “With 95 percent confidence”

Partially correct (P) if the response provides an appropriate interpretation of the interval that includes the first component *AND* only one of the other two components;

OR

if the response provides a correct interpretation of the confidence *level* in context without interpreting the specific interval.

Incorrect (I) if the response does not meet the criteria for E or P.

Notes:

- Clear indication of an inference to the random sample of 80 customers rather than to the population does not satisfy the first component and is scored I.
- Stating values that are unrealistic as proportions or percentages (including blanks) in the interpretation lowers the score one level (from E to P, or from P to I).
- When both the interpretation and the level of the interval are given, only the interpretation is scored. If the interpretation of the confidence level is incorrect, it is considered as poor communication in terms of holistic scoring.
- Any interpretation that implies the interval has a 95 percent chance (or possibility or probability) of capturing the population proportion is scored I.

Section 4 is scored as follows:

Essentially correct (E) if the response gives a correct interval estimate for the cost to the restaurant *AND* shows enough work to indicate how the interval was found.

Partially correct (P) if the response gives the correct interval estimate for the *number* of customers (565 to 1,160) who would fill the water cup with soda, including showing work, but does not multiply by \$0.25 to find the interval for the cost;

OR

if the response gives the correct interval estimate for the *expected cost* to the restaurant for an individual who asked for a water cup (\$0.05 to \$0.10), including showing work, but does not multiply by 3,000 to find the interval for the total cost;

OR

if the response makes a reasonable attempt to find the endpoints of the interval as $(0.1883)(3,000)(0.25)$ and $(0.3867)(3,000)(0.25)$, but makes an error such as using the sample size of 80 instead of the population size of 3,000;

2017 SCORING GUIDELINES

Question 2 (continued)

OR

if the response gives the correct interval without showing how it was found.

Incorrect (I) if the response does not meet the criteria for E or P.

Notes:

- The response in section 4 earns an E if it follows correctly from an incorrect interval found in part (a), even if the interval is unreasonable.
- Units (\$) are not required to earn an E.

Each essentially correct (E) section counts as 1 point, and a partially correct (P) section counts as $\frac{1}{2}$ point.

4 Complete Response

3 Substantial Response

2 Developing Response

1 Minimal Response

If a response is between two scores (for example, $2\frac{1}{2}$ points), use a holistic approach to decide whether to score up or down, depending on the strength of the response and communication.

**AP® STATISTICS
2016 SCORING GUIDELINES**

Question 5

Intent of Question

The primary goals of this question were to assess a student's ability to (1) construct and interpret a confidence interval for a population proportion; (2) explain why one of the conditions for inference is necessary; and (3) explain why a suggested procedure for constructing a confidence interval is incorrect.

Solution

Part (a):

The appropriate procedure is a one-sample z-interval for a population proportion. The problem stated the conditions for inference have been met, so they do not need to be checked. A 95 percent

confidence interval for the population proportion is given as $\hat{p} \pm z^* \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$, which is

$$0.37 \pm 1.96 \sqrt{\frac{(0.37)(0.63)}{1,048}} \approx 0.37 \pm 0.03 = (0.34, 0.40).$$

We are 95 percent confident that the population proportion of all adults in the U.S. who would have chosen the economy statement is between 0.34 and 0.40.

Part (b):

The condition is necessary because the formula for the confidence interval relies on the fact that the binomial distribution can be approximated by a normal distribution which then results in the sampling distribution of $\hat{p}$ being approximately normal. The approximation does not work well unless both $n\hat{p}$ and $n(1-\hat{p})$ are at least 10.

Part (c):

The suggested procedure is not appropriate because one of the requirements for using a two-sample z-interval for a difference between proportions is that the two proportions are based on two independent samples. In the situation described the two proportions come from a single sample and thus are not independent.

Scoring

This question is scored in three sections. Section 1 consists of the mechanics of the confidence interval in part (a), section 2 consists of interpreting the confidence interval in part (a), and section 3 consists of parts (b) and (c).

Section 1 is scored as follows:

Essentially correct (E) if the response includes the following two components:

1. States the correct procedure by name or formula
2. Calculates the confidence interval

Partially correct (P) if the response includes only one of the two components.

Incorrect (I) if the response does not meet the criteria for E or P.

**AP® STATISTICS
2016 SCORING GUIDELINES**

Question 5 (continued)

Notes:

- A formula with correct values is sufficient for component 2.
- Component 2 can never be satisfied if the response contains an incorrect formula.

Section 2 is scored as follows:

Essentially correct (E) if the response includes the following four components for the interval interpretation:

1. Estimates a proportion
2. Infers about the population
3. States 95 percent confidence
4. Includes context

Partially correct (P) if the response satisfies components 1 and 2 *AND* satisfies only one of components 3 or 4;

OR

if the response gives a correct interpretation of the confidence *level* in context without interpreting the specific interval.

Incorrect (I) if the response does not meet the criteria for E or P.

Notes:

- Any indication of an inference to the adults sampled rather than the population does not satisfy component 2 and is scored as I.
- Stating values that are unrealistic as proportions (including blanks) in the interpretation lowers the score by one level (from E to P or from P to I).
- When both the interpretation of the interval and the level are given, only the interpretation of the interval is scored.

Section 3 is scored as follows:

Essentially correct (E) if the response includes the following two components:

1. Part (b) states that the condition implies the sampling distribution of $\hat{p}$ is approximately normal *OR* the normal approximation to the binomial distribution is appropriate.
2. Part (c) indicates the procedure is not appropriate because the two proportions come from a single sample (dependent) rather than two (independent) samples.

Partially correct (P) if the response includes only one of the two components.

Incorrect (I) if the response does not meet the criteria for E or P.

Notes:

- Referring to the sampling distribution needing to be normal without explicitly stating $\hat{p}$ satisfies component 1.
- Referring to normal approximation must specify "to binomial" to satisfy component 1.
- Discussing the two-sample z-interval versus the two-proportion z-interval is extraneous information.

**AP® STATISTICS
2016 SCORING GUIDELINES****Question 5 (continued)**

- In part (c) component 2 can never be satisfied if a procedure that is not possible for one sample of categorical data is suggested.

4 Complete Response

Three parts essentially correct

3 Substantial Response

Two parts essentially correct and one part partially correct

2 Developing Response

Two parts essentially correct and no parts partially correct

OR

One part essentially correct and one or two parts partially correct

OR

Three parts partially correct

1 Minimal Response

One part essentially correct

OR

No parts essentially correct and two parts partially correct