

2017

AP[®]

CollegeBoard

AP Statistics

Scoring Guidelines

2017 SCORING GUIDELINES

Question 1

Intent of Question

The primary goals of this question were to assess a student's ability to (1) explain statistical terms used when describing the relationship between two variables; (2) interpret the slope of a linear regression equation; and (3) calculate a value of y when given a regression equation, a value of x , and a residual.

Solution**Part (a):**

In the context of a scatterplot in which y represents weight and x represents length, the following are defined.

A positive relationship means that wolves with higher values of length also tend to have higher weights.

A linear relationship means that as length increases by one meter, weight tends to change by a constant amount, on average.

A strong relationship means that the data points fall close to a line (or curve).

Part (b):

The slope of 35.02 indicates that two wolves that differ by one meter in length are predicted to differ by 35.02 kilograms in weight, with the longer wolf having the greater weight.

Part (c):

In general, a residual is equal to actual weight minus predicted weight, or equivalently,

$$\text{actual weight} = \text{predicted weight} + \text{residual}.$$

For the wolf with length 1.4 meters and residual of -9.67 , the predicted weight is

$$-16.46 + 35.02(1.4) = 32.568 \text{ kilograms}.$$

Therefore, the actual weight of the wolf is $32.568 + (-9.67) = 22.898$ kilograms.

Scoring

Parts (a), (b), and (c) are scored as essentially correct (E), partially correct (P), or incorrect (I).

Part (a) is scored as follows:

Essentially correct (E) if the response includes the following four components:

1. A reasonable definition of positive
2. A reasonable definition of linear
3. A reasonable definition of strong
4. At least one definition in context

2017 SCORING GUIDELINES

Question 1 (continued)

Partially correct (P) if the response includes only three of the four components.

Incorrect (I) if the response does not meet the conditions for E or P.

Notes:

- The description of a positive relationship should clearly indicate that relatively low values of one variable tend to appear with relatively low values of the other variable, and relatively high values of the first variable tend to appear with relatively high values of the other variable.
 - Examples of acceptable responses:
 - As length increases, so does weight.
 - Longer wolves weigh more.
 - The points on the graph go up as you move from left to right.
 - Examples of unacceptable responses:
 - As length goes up, weight changes.
 - Both length and weight get bigger.
 - The correlation is greater than 0.
- The description of a linear relationship can take one of two approaches: the data pattern (data points exhibit the pattern of a line in the graph) or the constant rate of change (as the explanatory variable changes, the response variable exhibits a constant rate of change).
 - Examples of acceptable responses:
 - The points generally follow a straight line.
 - The relationship between x and y is straight.
 - Length and width have a constant slope.
 - Examples of unacceptable responses:
 - The points all line up.
 - You can draw a straight line through the points.
 - There is a positive correlation.
 - Every increase in x yields a 35.02 increase in y .
- The description of strong should indicate how close points are to a line.
 - Examples of acceptable responses:
 - Observed values are close to predicted values.
 - Deviations from the least-squares regression line are small.
 - The correlation coefficient is close to 1.
 - Examples of unacceptable responses:
 - All the points are close together.
 - The scatterplots are clustered together.
 - There is a high positive correlation.
- Context can be shown by referring to length and weight or by using meters and kilograms.
- Sketches and graphs can be used to help clarify definitions, but a sketch alone cannot satisfy a definition component.

Part (b) is scored as follows:

Essentially correct (E) if the response includes the following three components:

1. The correct value of 35.02 for the slope.
2. An interpretation that includes an increase of a specified amount of weight for each unit increase in length.
3. An indication that the relationship is not exact by using words such as “on average” or “predicted weight.”

2017 SCORING GUIDELINES

Question 1 (continued)

Partially correct (P) if the response includes only two of the three components.

Note: If the response identifies the slope as -16.46 (the intercept value), the second component is satisfied only if the response states that for each one-meter increase in length there is a *decrease* in predicted or average weight of 16.46 kilograms.

Incorrect (I) if the response does not meet the criteria for E or P.

Part (c) is scored as follows:

Essentially correct (E) if the response includes the following two components:

1. A correct computation for the predicted value 32.568 kilograms.
2. A correct computation for the actual weight 22.9 kilograms using the given residual and the predicted value.

Partially correct (P) if the response provides a correct computation for the predicted value but is not able to complete the correct calculation of the actual weight, including if the residual is defined in the wrong direction as $(\text{predicted weight}) - (\text{actual weight})$ to give an answer of 42.24 kilograms;

OR

if the response provides an incorrect value for the predicted weight, but then uses that value correctly to determine the actual weight as $(\text{predicted weight}) + \text{residual} = (\text{predicted weight}) + (-9.67)$;

OR

if the response provides a correct answer for the actual weight but does not give sufficient information to determine how it was calculated.

Incorrect (I) if the response does not meet the criteria for E or P.

Notes:

- The expression $-16.46 + 35.02(1.4)$ is enough to satisfy the first component.
- The equation $-16.46 + 35.02(1.4) - 9.67 = 22.9$ satisfies both components.
- Arithmetic mistakes are overlooked if they do not lead to an unreasonable answer (such as a negative value). For example, $32.568 + (-9.67) = 21.9$ satisfies the second component.

2017 SCORING GUIDELINES**Question 1 (continued)****4 Complete Response**

Three parts essentially correct

3 Substantial Response

Two parts essentially correct and one part partially correct

2 Developing Response

Two parts essentially correct and no parts partially correct

OR

One part essentially correct and one or two parts partially correct

OR

Three parts partially correct

1 Minimal Response

One part essentially correct

OR

No parts essentially correct and two parts partially correct

2017 SCORING GUIDELINES

Question 2

Intent of Question

The primary goals of this question were to assess a student's ability to (1) construct and interpret a confidence interval for a population proportion and (2) use a confidence interval for a proportion to find a confidence interval for a dollar amount that can be calculated using that proportion.

Solution

Part (a):

Step 1: Identify the appropriate confidence interval by name or formula and check appropriate conditions.

The appropriate procedure is a one-sample z -interval for a population proportion p . In this case, the population is all customers of the restaurant who ask for a water cup, and p is the proportion of that population who will fill the cup with a soft drink. The appropriate formula is $\hat{p} \pm z^* \sqrt{\frac{\hat{p}(1 - \hat{p})}{n}}$.

Conditions:

1. Random sample
2. Large sample (number of successes $n\hat{p} \geq 10$ and number of failures $n(1 - \hat{p}) \geq 10$)

For condition 1 the stem of the problem states that a random sample of customers who asked for a water cup was used.

For condition 2 the number of successes (filled cup with soft drink) is 23 and the number of failures is 57, both of which are greater than 10.

Step 2: Correct mechanics

The sample proportion is $\hat{p} = \frac{23}{80} = 0.2875$. The confidence interval is

$$\begin{aligned} & 0.2875 \pm 1.96 \sqrt{\frac{0.2875(1 - 0.2875)}{80}} \\ & = 0.2875 \pm 1.96(0.0506) \quad \text{or } 0.1883 \text{ to } 0.3867. \\ & = 0.2875 \pm 0.0992 \end{aligned}$$

Step 3: Interpretation

We can be 95 percent confident that in the population of all customers of the restaurant who ask for a water cup, the proportion who will fill it with a soft drink is between 0.1883 and 0.3867.

Part (b):

Using the confidence interval in part (a), a 95 percent interval estimate for the number of customers in June who asked for a water cup but then filled it with a soft drink is $3,000 \times 0.1883$ to $3,000 \times 0.3867$, or 565 to 1,160. At a cost of \$0.25 per customer, a 95 percent interval estimate for the cost to the restaurant in June is \$141.25 to \$290.00.

2017 SCORING GUIDELINES

Question 2 (continued)

Scoring

This question is scored in four sections. Section 1 consists of step 1 in part (a), section 2 consists of step 2 in part (a), section 3 consists of step 3 in part (a), and section 4 consists of part (b). Each section is scored as essentially correct (E), partially correct (P), or incorrect (I).

Section 1 is scored as follows:

Essentially correct (E) if the one-sample z -interval for a proportion is identified (either by name or formula) *AND* both conditions (random sampling and large sample) are adequately addressed.

Partially correct (P) if the response identifies the correct procedure *BUT* adequately addresses only one of the two required conditions;

OR

if the response does not identify the correct procedure *BUT* adequately addresses both required conditions.

Incorrect (I) if the response does not meet the criteria for E or P.

Notes:

- Stating the large sample condition without verifying it is not sufficient. The response must use specific numerical values to adequately address the condition.
- If the response includes additional inappropriate conditions, such as $n \geq 30$ or requiring a normal population, then the response earns at most P for section 1.
- Stating and checking a condition about the size of the sample relative to the size of the population is appropriate but not required.
- Any statement of hypotheses, description of the population, or definition of the parameter should be considered extraneous. However, if such statements are included and incorrect, they are considered as poor communication in terms of holistic scoring.

Section 2 is scored as follows:

Essentially correct (E) if the response gives the correct 95 percent confidence interval. Supporting work is not required, but if included, it must be correct.

Partially correct (P) if the response gives a correct confidence interval with incorrect (but appropriate) supporting work shown;

OR

if the response gives an incorrect but reasonable confidence interval with appropriate supporting work shown — for instance, if a value other than 1.96 is used for the critical value or a value other than $\frac{23}{80}$ is used for $\hat{p}$.

Incorrect (I) if the response does not meet the criteria for E or P.

Notes:

- Appropriate supporting working must have the form:
proportion $\pm$ (critical value)(SE/SD of proportion).

2017 SCORING GUIDELINES

Question 2 (continued)

- A confidence interval that has both endpoints outside the interval from 0 to 1 is considered unreasonable.

Section 3 is scored as follows:

Essentially correct (E) if the response provides an appropriate interpretation of the interval that includes the following three components:

1. Conveys inference about a population proportion
2. Demonstrates a clear understanding that the parameter is the proportion of the water cup population that fills the cup with a soft drink
3. Mentions 95 percent confidence and interprets it correctly using words such as “We can be 95 percent confident” or “With 95 percent confidence”

Partially correct (P) if the response provides an appropriate interpretation of the interval that includes the first component *AND* only one of the other two components;

OR

if the response provides a correct interpretation of the confidence *level* in context without interpreting the specific interval.

Incorrect (I) if the response does not meet the criteria for E or P.

Notes:

- Clear indication of an inference to the random sample of 80 customers rather than to the population does not satisfy the first component and is scored I.
- Stating values that are unrealistic as proportions or percentages (including blanks) in the interpretation lowers the score one level (from E to P, or from P to I).
- When both the interpretation and the level of the interval are given, only the interpretation is scored. If the interpretation of the confidence level is incorrect, it is considered as poor communication in terms of holistic scoring.
- Any interpretation that implies the interval has a 95 percent chance (or possibility or probability) of capturing the population proportion is scored I.

Section 4 is scored as follows:

Essentially correct (E) if the response gives a correct interval estimate for the cost to the restaurant *AND* shows enough work to indicate how the interval was found.

Partially correct (P) if the response gives the correct interval estimate for the *number* of customers (565 to 1,160) who would fill the water cup with soda, including showing work, but does not multiply by \$0.25 to find the interval for the cost;

OR

if the response gives the correct interval estimate for the *expected cost* to the restaurant for an individual who asked for a water cup (\$0.05 to \$0.10), including showing work, but does not multiply by 3,000 to find the interval for the total cost;

OR

if the response makes a reasonable attempt to find the endpoints of the interval as $(0.1883)(3,000)(0.25)$ and $(0.3867)(3,000)(0.25)$, but makes an error such as using the sample size of 80 instead of the population size of 3,000;

2017 SCORING GUIDELINES**Question 2 (continued)**

OR

if the response gives the correct interval without showing how it was found.

Incorrect (I) if the response does not meet the criteria for E or P.

Notes:

- The response in section 4 earns an E if it follows correctly from an incorrect interval found in part (a), even if the interval is unreasonable.
- Units (\$) are not required to earn an E.

Each essentially correct (E) section counts as 1 point, and a partially correct (P) section counts as $\frac{1}{2}$ point.

- | | |
|----------|-----------------------------|
| 4 | Complete Response |
| 3 | Substantial Response |
| 2 | Developing Response |
| 1 | Minimal Response |

If a response is between two scores (for example, $2\frac{1}{2}$ points), use a holistic approach to decide whether to score up or down, depending on the strength of the response and communication.

2017 SCORING GUIDELINES

Question 3

Intent of Question

The primary goals of this question were to assess a student's ability to (1) calculate a probability from a normal distribution; (2) calculate a weighted probability from two individual probabilities; and (3) calculate a conditional probability for dependent events when individual and joint probabilities are provided.

Solution**Part (a):**

Let X denote the diameter of a randomly selected melon from Distributor J. X has an approximately normal distribution with mean 133 mm and standard deviation 5 mm.

The z-score for a diameter of 137 mm is $z = \frac{137 - 133}{5} = \frac{4}{5} = 0.8$.

Therefore, $P(X > 137) = P(Z > 0.8) = 1 - 0.7881 = 0.2119$.

Part (b):

Define events:

J : melon is from Distributor J


K : melon is from Distributor K

G : melon diameter is greater than 137 mm

$$\begin{aligned} P(G) &= P(G | J) \times P(J) + P(G | K) \times P(K) \\ &= (0.2119)(0.7) + (0.8413)(0.3) \\ &= 0.1483 + 0.2524 \\ &= 0.4007 \end{aligned}$$

For a randomly selected melon from the grocery store,

OR



From the tree diagram, $P(G) = P(G \text{ and } J) + P(G \text{ and } K) = 0.1483 + 0.2524 = 0.4007$.

2017 SCORING GUIDELINES

Question 3 (continued)

Part (c):

Using the events defined in part (b), the requested probability is

$$P(J | G) = \frac{P(J \text{ and } G)}{P(G)} = \frac{P(G | J)P(J)}{P(G)} = \frac{(0.2119)(0.7)}{0.4007} = \frac{0.1483}{0.4007} = 0.3701.$$

Scoring

Parts (a), (b), and (c) are scored as essentially correct (E), partially correct (P), or incorrect (I).

Part (a) is scored as follows:

Essentially correct (E) if the response provides the following three components:

1. Normality and parameters: Indicates use of a normal (or approximately normal) distribution and clearly identifies the correct parameter values. (Showing correct components in a z-score, labeling the mean and standard deviation in a normalcdf calculator statement, or drawing a normal curve with sufficient values marked to indicate the mean and standard deviation are all sufficient methods for identifying the correct parameter values.)
2. Boundary and direction: Uses the correct boundary value of $x = 137$ or $z = 0.8$ and the correct direction. (Showing correct boundary and direction in a probability statement using "greater than," labeling the lower limit and upper limit in a normalcdf calculator statement, drawing a normal curve with 137 labeled and an indication that the area of interest is to the right of 137, or a conclusion in words using "greater than.")
3. Probability: Reports the correct normal probability consistent with the response's setup described in components 1 and 2.

Partially correct (P) if the response correctly provides only two of the three components.

Incorrect (I) if the response gives the answer with no work shown or otherwise does not satisfy the criteria for E or P.

Notes:

- An inconsistency in calculations lowers the score for part (a) by one level (from E to P, or from P to I). For instance, if the response states the requested probability as $P(X > 137)$, but actually computes $P(X < 137)$.
- An error in statistical notation, such as using s instead of σ for the standard deviation or $\bar{x}$ instead of μ for the mean, does not satisfy component 1.
- A correct direction can be obtained by using a left direction and an upper bound of 137 and then subtracting the value of the cumulative probability from 1. However, simply showing the calculation $1 - 0.7881 = 0.2119$ does NOT give a correct direction.
- Consider any steps of a hypothesis test as extraneous work.
- Standard notations such as $N(133,5)$ or $N(133,25)$ satisfy the first component.
- A sketch of the normal curve with 133 and 138 in the appropriate positions satisfies the first component.
- Use of a value other than 137 does not satisfy the second component.
- If the only error in part (a) is the reversal of the numerator for the z-score $(133 - 137)$, the response is scored as P.

2017 SCORING GUIDELINES

Question 3 (continued)

- For calculator notation:
 - ncdf is sufficient to indicate use of the normal distribution.
 - Stating the lower and upper bounds is sufficient for identifying boundary and direction.
 - Minimum notation, such as “u” for upper, is sufficient for identifying boundary.
 - The minimum value for an upper bound must be at least 4 standard deviations above the mean.

Part (b) is scored as follows:

Essentially correct (E) if the probability is computed correctly, *AND* work is shown that includes correct numerical values using a formula, end results from a tree diagram, or some other appropriate method.

Partially correct (P) if the response provides a correct strategy for finding the probability, such as a formula or tree diagram, but uses one or more incorrect values or does not use the end result from the tree;

OR

if the response gives the correct probability but not enough work is shown to determine how it was found.

Incorrect if the response does not meet the criteria for E or P.

Notes:

- A response that uses 0.3 for J and 0.7 for K earns a P if no other mistakes are made.
- Tree diagrams:
 - If used, the multiplication is implied, but the addition must be indicated.
 - If not used, both the multiplication and the addition must be indicated.
- The shown work might be found in part (c).
- Probabilities can be shown in a table. Addition is implied if “Total” is labeled, as shown.

Intersection	G	Not G	Total
J	0.1483	0.5517	0.7
K	0.2524	0.0576	0.3
Total	0.4007	0.5993	

Part (c) is scored as follows:

Essentially correct (E) if the probability is computed correctly *AND* work is shown that illustrates how the probability was found.

Partially correct (P) if the response provides a strategy for finding the probability for dependent variables, such as the appropriate formula or end results from the correct tree diagram, but uses one or more incorrect values or does not use the end result from the tree;

OR

if the response gives the correct probability but not enough work is shown to determine how it was found.

Incorrect (I) if the response does not meet the criteria for E or P.

2017 SCORING GUIDELINES**Question 3 (continued)**

Notes:

- Part (c) earns an E if a correct answer follows from incorrect values used from part (a) and/or part (b), provided all values are between 0 and 1, inclusive.
- A transcription error in a response can be ignored if correct work is shown.
- For any part, if the resulting probability or part of the calculation of the probability uses a value that is not between 0 and 1, inclusive, the score is lowered by one level (from E to P, or from P to I).
- In any part, unsupported answers are scored as I.

4 Complete Response

Three parts essentially correct

3 Substantial Response

Two parts essentially correct and one part partially correct

2 Developing Response

Two parts essentially correct and no parts partially correct

OR

One part essentially correct and one or two parts partially correct

OR

Three parts partially correct

1 Minimal Response

One part essentially correct

OR

No parts essentially correct and two parts partially correct

2017 SCORING GUIDELINES

Question 4

Intent of Question

The primary goals of this question were to assess a student's ability to use boxplots to (1) compare multiple sets of data; (2) identify which set of data is most likely to have produced a particular summary value; and (3) determine which variable is most useful for classifying a new observation.

Solution**Part (a):**

The median value for the percent of chemical Z in the pottery pieces is similar for all three sites, at about 7 percent. The ranges for the percent of chemical Z are much different for the three sites, with the smallest range of about 2 percent (from 6 percent to 8 percent) at Site II, a range of about 6 percent (from about 4 percent to 10 percent) at Site I, and the largest range of about 8 percent (from about 3 percent to 11 percent) at Site III.

Part (b):

- (i) The piece most likely originated at Site III. Although values outside of the range of data observed in the samples would be possible, using the available data results in approximate minimum and maximum sums of the percents for the three chemicals as shown in the table below. Site III is the only site in which 20.5 falls between the sums of the minimum and maximum values.

	Site I		Site II		Site III	
Chemical	Min	Max	Min	Max	Min	Max
X	6	8	5	7	5	7.5
Y	11	15	1.9	4	6	8
Z	4	10	6	8	3	11
Sum	21	33	12.9	19	14	26.5

- (ii) Chemical Y would be most useful, because the distribution of the percentages of total weights at the three sites do not overlap. The distributions of chemicals X and Z have substantial overlap.

Scoring

This question is scored in three sections. Section 1 consists of part (a), section 2 consists of part (b-i), and section 3 consists of part (b-ii). Each section is scored as essentially correct (E), partially correct (P), or incorrect (I).

Section 1 is scored as follows:

Essentially correct (E) if the response includes the following three components:

1. Recognition that the medians or centers are almost the same for the three sites
2. Recognition that the variability (ranges, IQRs, spread) is different across the three sites
3. Context is included

Partially correct (P) if the response includes only two of the three components.

Incorrect (I) if the response includes at most one of the three components.

2017 SCORING GUIDELINES

Question 4 (continued)

Notes:

- In all sections, comments about shape should be ignored because complete shape information is not obtainable from boxplots.
- Responses are not required to give numerical values. If responses provide numerical values, any reasonable approximation from the boxplots is acceptable.
- Because the boxplots are all symmetric, it is acceptable if the response discusses means instead of medians.
- Any discussion of chemical X and chemical Y is considered extraneous.
- Context is satisfied by any of the following references: site, chemical, weight, total weight, X, Y, or Z.

Section 2 is scored as follows:

Essentially correct (E) if the response includes the following three components:

1. Site III is chosen.
2. Sums of the minimum and maximum are computed for the three chemicals at each site.
3. A reasonable numerical justification is given involving sums of a statistical measure across the three chemicals to choose Site III.

Partially correct (P) if the response includes only two of the three components.

Incorrect (I) if the response includes at most one of the three components.

Notes:

- If the response computes only the sum of the minimums for Site I and the sum of the maximums for Site II and recognizes that this is sufficient, the response is scored E.
- If an alternative measure is used that involves sums of the three chemicals, such as the sum of the medians or the sums of the first quartiles and sums of the third quartiles, instead of the minimum and maximum sums, the second component is not satisfied, but the third component might be satisfied.
 - If the response explicitly or implicitly compares the alternate sum to the other two sites (for example, by indicating that the sum is the closest to 20.5 percent or by listing the sums for all three sites) the response is scored P.
 - If the response does not have an implicit or explicit comparison, the response is scored I.
- If either Site I or Site II is identified as the correct choice, no matter how that choice is justified, the response is scored I.
- The approximate sums of the medians are 27.5 for Site I, 16 for Site II, and 20 for Site III.

Section 3 is scored as follows:

Essentially correct (E) if the response chooses chemical Y *AND* gives a reasonable justification based on the fact that the distributions of chemical Y are distinctive across sites.

Partially correct (P) if the response chooses chemical Y *AND* provides justification based on the boxplots, but does not clearly explain that the distributions of chemical Y are distinctive across sites;

OR

if the response correctly discusses that the distributions of chemical Y are distinctive across sites, but never explicitly chooses chemical Y as the best choice, for instance, by stating only that there is substantial overlap across sites for chemicals X and Z but no overlap for chemical Y.

2017 SCORING GUIDELINES

Question 4 (continued)

Incorrect (I) if the response does not meet the criteria for E or P.

Notes:

- To justify that the distributions of chemical Y are distinctive across sites, the justification must address both location and variability of the boxplots; for example, by stating that the boxplots do not overlap for chemical Y.
- If the response chooses chemical X or Z *OR* chooses chemical Y with no reasonable justification, the response is scored I.
- The justification that the distributions of chemical Y are distinctive across sites:
 - The following are acceptable because both location and variability are addressed. Such responses are scored E.
 - The boxplots for chemical Y do not overlap, or the boxplots for chemicals X and Z overlap.
 - All values of Site I are high, all values of Site II are low, and all values of Site III are in the middle.
 - The ranges never intersect.
 - The boxplots share no data.
 - Has completely different percentages at each site.
 - The following are incomplete justifications and are scored P.
 - The boxplots vary.
 - Chemical Y varies the most.
 - Chemical Y has the greatest variation.
 - The variation between/among sites is the largest.
 - The boxplots are different.
 - The medians/means differ.
 - The medians/means are most variable.
 - There is a difference in the percentages of chemical Y for each site.
 - The distribution of percents differs the most among the sites.

2017 SCORING GUIDELINES**Question 4 (continued)****4 Complete Response**

Three sections essentially correct

3 Substantial Response

Two sections essentially correct and one section partially correct

2 Developing Response

Two sections essentially correct and no sections partially correct

OR

One section essentially correct and one or two sections partially correct

OR

Three sections partially correct

1 Minimal Response

One section essentially correct

OR

No sections essentially correct and two sections partially correct

2017 SCORING GUIDELINES

Question 5

Intent of Question

The primary goal of this question was to assess a student's ability to identify, set up, perform, and interpret the results of an appropriate hypothesis test to address a particular question. More specific goals were to assess a student's ability to (1) state appropriate hypotheses; (2) identify the appropriate statistical test procedure and check appropriate conditions for inference; (3) calculate the appropriate test statistic and p -value; and (4) draw an appropriate conclusion, with justification, in the context of the study.

Solution

Step 1: State a correct pair of hypotheses.

The null hypothesis is that age group at diagnosis and gender are independent (that is, they are not associated) for the population of people currently being treated for schizophrenia.

The alternative hypothesis is that age group at diagnosis and gender are not independent for the population of people currently being treated for schizophrenia.

Step 2: Identify a correct test procedure (by name or formula) and check appropriate conditions.

The appropriate test is a chi-square test of independence.

The conditions for this test are satisfied because:

1. The question states that the sample was randomly selected.
2. The expected counts for the eight cells of the table are at least 5, as seen in the following table, with expected counts shown below observed counts.

		Age at Diagnosis				
		20 to 29	30 to 39	40 to 49	50 to 59	Total
Women	46	40	21	12	119	
	56.91	36.22	17.25	8.62		
Men	53	23	9	3	88	
	42.09	26.78	12.75	6.38		

Step 3: Find the value of the test statistic and the p -value.

The test statistic is calculated as $\chi^2 = \sum \frac{(O - E)^2}{E}$, or

$$\begin{aligned}\chi^2 &= 2.093 + 0.395 + 0.817 + 1.322 \\ &\quad + 2.830 + 0.534 + 1.105 + 1.788 \\ &= 10.884.\end{aligned}$$

The p -value is $P(\chi^2 \geq 10.884) = 0.012$, based on $(4 - 1) \times (2 - 1) = 3$ degrees of freedom.

2017 SCORING GUIDELINES

Question 5 (continued)

Step 4: State the conclusion in context, with linkage to the p -value.

Because the p -value is very small (for instance much smaller than $\alpha = 0.05$), we reject the null hypothesis and conclude that the sample data provide strong evidence that there is an association between age group at diagnosis and gender for the population currently being treated for schizophrenia.

Scoring

This question is scored in three sections. Section 1 consists of steps 1 and 2 (stating the correct hypotheses, identifying the appropriate test procedure, and checking the technical conditions); section 2 consists of step 3 (performing the correct mechanics); and section 3 consists of step 4 (stating a correct conclusion with justification). Sections 1, 2, and 3 are scored as essentially correct (E), partially correct (P), or incorrect (I).

Section 1 is scored as follows:

Essentially correct (E) if the response correctly includes the following three components:

1. States BOTH hypotheses correctly with context included in at least one of them
2. Identifies a chi-square test of independence by name or formula
3. Verifies appropriate conditions that minimally include the condition for the expected counts and do not include any incorrect conditions (such as normality)

Partially correct (P) if the response includes only two of the three components.

Incorrect (I) if the response includes at most one of the three components.

Notes:

- Stating the expected count condition is not sufficient; the condition must be checked by reporting the expected counts, or minimally by showing that the smallest expected count is at least 5.
- The random sample condition was stated in the stem so need not be explicitly checked.
- If the null and alternative hypothesis are correctly stated in terms of population proportions, component 1 is satisfied. For example:

$H_0 : p_1 = p_2 = p_3 = p_4$, where p_i is the population proportion of women at each indicated age group, 1, 2, 3, or 4, who are currently being treated for schizophrenia.

H_a : At least one of the population proportions, p_1, p_2, p_3, p_4 , differs from the other three.

OR

H_a : The population proportions for the four age groups are not all the same.

Section 2 is scored as follows.

Essentially correct (E) if the response correctly calculates the following two values:

1. The value of the chi-square test statistic
2. The p -value, critical value, or p -value range from chi-square table

Partially correct (P) if the response correctly calculates only one of the two values.

2017 SCORING GUIDELINES

Question 5 (continued)

Incorrect (I) if the response does not satisfy the criteria for E or P.

Notes:

- If the response makes an error in one calculation, subsequent calculations are considered correct if they follow correctly from the initial miscalculation.
- With 3 degrees of freedom, the correct critical value is 7.81 for a significance level of 0.05 and 11.34 for a significance level of 0.01.
- Work does not have to be shown for calculations of test statistic or p -value. However, if incorrect work (other than minor arithmetic/transcription errors) is shown it is considered to be an incorrect calculation of the respective component, even if the correct value is given.
- If a response provides a test statistic that is not a chi-square test statistic, section 2 is scored I.

Section 3 is scored as follows.

Essentially correct (E) if the response includes the following three components:

1. A correct conclusion about the alternative hypothesis.
2. Justification of the conclusion based on linkage between the p -value and a reasonable alpha (or linkage between test statistic and critical value).
3. The conclusion is stated in context.

Partially correct (P) if the response includes only two of the three components.

Incorrect (I) if the response includes at most one of the three components.

Notes:

- If the response provides a correct decision, in context, with linkage to the p -value, but the decision is stated in terms of the null hypothesis with no conclusion about the alternative hypothesis, component 1 is not satisfied.
- Incorrect statistical statements are considered incorrect conclusions for the hypothesis test and do not satisfy component 1.
- If the conclusion is consistent with the p -value from section 2, and also in context with justification based on linkage to the p -value, section 3 is scored E.
- If no alpha level is given, the solution must be explicit about the linkage by giving a correct interpretation of the p -value or explaining how the conclusion follows from the p -value. For example, stating that because the p -value is small, we reject the null hypothesis or stating that because the p -value is large, we do not reject the null hypothesis.
- A decision about the null hypothesis (reject H_0 or fail to reject H_0) is not required, but if such a statement is given the scoring of the decision is considered in component 2.

2017 SCORING GUIDELINES**Question 5 (continued)****4 Complete Response**

Three sections essentially correct

3 Substantial Response

Two sections essentially correct and one section partially correct

2 Developing Response

Two sections essentially correct and no sections partially correct

OR

One section essentially correct and one or two sections partially correct

OR

Three sections partially correct

1 Minimal Response

One section essentially correct

OR

No sections essentially correct and two sections partially correct

OR

Section 1 partially correct and the other two sections incorrect

2017 SCORING GUIDELINES

Question 6

Intent of Question

The primary goals of this question were to assess a student's ability to (1) calculate probabilities associated with treatment and control group memberships for two different methods of random assignment and (2) justify which method of random assignment is more appropriate in a given situation.

Solution**Part (a):**

- (i) Let T (tail) represent being assigned to the treatment group and H (head) represent being assigned to the control group for each coin flip. The process stops when either the treatment group or the control group has two members. The outcomes and their probabilities are as follows.

Arrangement	A	B	C	D	E	F
Chip outcomes	TT	THT	THH	HH	HTH	HTT
Calculation	$\left(\frac{1}{2}\right)\left(\frac{1}{2}\right)$	$\left(\frac{1}{2}\right)\left(\frac{1}{2}\right)\left(\frac{1}{2}\right)$	$\left(\frac{1}{2}\right)\left(\frac{1}{2}\right)\left(\frac{1}{2}\right)$	$\left(\frac{1}{2}\right)\left(\frac{1}{2}\right)$	$\left(\frac{1}{2}\right)\left(\frac{1}{2}\right)\left(\frac{1}{2}\right)$	$\left(\frac{1}{2}\right)\left(\frac{1}{2}\right)\left(\frac{1}{2}\right)$
Probability	$\frac{1}{4}$	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{1}{4}$	$\frac{1}{8}$	$\frac{1}{8}$

- (ii) Man 1 and Man 2 are assigned to the same group for arrangements A and D, so the probability is

$$P(A) + P(D) = \frac{1}{4} + \frac{1}{4} = \frac{1}{2}.$$

Part (b):

- (i) Let T represent being assigned to the treatment group and C represent being assigned to the control group for each chip drawn. The process stops when either the treatment group or the control group has two members. The probabilities differ from the coin flip method because chips are drawn *without* replacement. The outcomes and their probabilities are as follows.

Arrangement	A	B	C	D	E	F
Chip outcomes	TT	TCT	TCC	CC	CTC	CTT
Calculation	$\left(\frac{2}{4}\right)\left(\frac{1}{3}\right)$	$\left(\frac{2}{4}\right)\left(\frac{2}{3}\right)\left(\frac{1}{2}\right)$	$\left(\frac{2}{4}\right)\left(\frac{2}{3}\right)\left(\frac{1}{2}\right)$	$\left(\frac{2}{4}\right)\left(\frac{1}{3}\right)$	$\left(\frac{2}{4}\right)\left(\frac{2}{3}\right)\left(\frac{1}{2}\right)$	$\left(\frac{2}{4}\right)\left(\frac{2}{3}\right)\left(\frac{1}{2}\right)$
Probability	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$	$\frac{1}{6}$

- (ii) Man 1 and Man 2 are assigned to the same group for arrangements A and D, so the probability is

$$P(A) + P(D) = \frac{1}{6} + \frac{1}{6} = \frac{1}{3}.$$

2017 SCORING GUIDELINES

Question 6 (continued)

Part (c):

Use the chip method. The chip method gives equal probability to all possible arrangements, but the coin method does not, as shown in the tables from parts (a-i) and (b-i). Furthermore, the coin method is more likely to result in imbalanced treatment groups with regard to students and teachers, based on the probabilities in parts (a-ii) and (b-ii). If food preferences for teachers are different than for students, the imbalance is a problem. For example, if one treatment group consists entirely of students, it would be impossible to know if a difference in the response variable is due to the treatment (type of meal) or the role of the person at the school (teacher or student).

Scoring

Parts (a), (b), and (c) are each scored as essentially correct (E), partially correct (P), or incorrect (I).

Part (a) is scored as follows:

Essentially correct (E) if the response includes the following three components:

1. The correct probability for each arrangement in the table of (i).
2. Appropriate justification is shown in (i).
3. The correct probability is given in (ii).

Partially correct (P) if the response includes only two of the three components.

Incorrect (I) if the response does not meet the criteria for E or P.

Notes:

- For component 2, examples of appropriate justification include the following:
 - Shows calculation of a correct probability for at least one of arrangements A and D *AND* shows calculation of a correct probability for at least one of arrangements B, C, E, and F. Calculations must be linked to a specific arrangement with a label, by proximity to an arrangement in the table, or by following the structure of the table or display of arrangements.
 - Lists correct coin outcomes for at least one of arrangements A and D *AND* lists correct coin outcomes for at least one of arrangements B, C, E, and F (for example, A: TT and B: THT). Lists of outcomes must be linked to a specific arrangement with a label, by proximity to an arrangement in the table, or by following the structure of the table or display of arrangements.
 - Shows a tree diagram that ends with six branches corresponding to the six arrangements. Probabilities or coin outcomes must be listed on the branches, but arrangements do not need to be identified.
 - Shows a tree diagram that ends with eight (or 16) branches where the branches leading to at least one of arrangements A and D are identified, and the branches leading to at least one of arrangements B, C, E, and F are identified. Probabilities or coin outcomes must be listed on the branches.
- If an incorrect probability is given in part (ii), component 3 is satisfied if the probability is between 0 and 1 and:
 - is the result of a minor arithmetic error; *OR*
 - is the result of adding the probabilities of arrangements A and D from the table in (i).

2017 SCORING GUIDELINES

Question 6 (continued)

Part (b) is scored as follows:

Essentially correct (E) if the response includes the following two components:

1. The correct probability for each arrangement in the table of (i).
2. The correct probability is given in (ii).

Partially correct (P) if the response includes only one of the three components.

Incorrect (I) if the response does not meet the criteria for E or P.

Note: If an incorrect probability is given in (ii), component 2 is satisfied if the probability is between 0 and 1 and

- is the result of a minor arithmetic error; *OR*
- is the result of adding the probabilities of arrangements A and D from the table in (i); *OR*
- work is shown that demonstrates the same incorrect approach is used in both parts (a-ii) and (b-ii). For example, component 2 is satisfied if the response forgets to include $P(D)$ and says

$P(A) = \frac{1}{4}$ in (a-ii) and $P(A) = \frac{1}{6}$ in part (b-ii). However, unlabeled answers of $\frac{1}{4}$ in (a-ii) and $\frac{1}{6}$ in (b-ii) would not satisfy component 2 because it is unclear if the same approach is being used.

Part (c) is scored as follows:

Essentially correct (E) if the response chooses the chip method AND includes the following three components:

1. Provides a statistical benefit to the chip method. For example, stating that all arrangements are equally likely.
2. Provides a statistical drawback to the coin method. For example, stating that the coin method is more likely to result in imbalanced treatment groups.
3. States that the responses (opinions, food preferences) of teachers and students might be different.

Partially correct if the response chooses the chip method and includes only one or two of the three components.

Incorrect (I) if the response does not meet the criteria for E or P.

Notes:

- A correct comparative statement can be used to satisfy both components 1 and 2. For example, stating that the coin method is *more likely* to result in a disproportionate number of students in one treatment group, or that the chip method has a *greater chance* of providing roughly equivalent treatment groups.
- Benefits and drawbacks must be about the probabilities of the arrangements (from the tables in (i)), or the probabilities of imbalanced (balanced) groups (from the questions in (ii)), or both.
- If the response chooses the coin method, look back to see if the coin method should be the preferred method based on incorrect work from parts (a) and (b). If so, score part (c) using the corresponding three components.

2017 SCORING GUIDELINES**Question 6 (continued)**

- If the response says that it does not matter which method to use, look back to see if the tables are the same in parts (a-i) and (b-i) or if the probabilities are the same in parts (a-ii) and (b-ii). If so, part (c) is partially correct if the response justifies the decision by stating that the two tables or the two probabilities are the same. To be essentially correct, the response also needs to satisfy component 3.
- If the response does not choose a method, deduct one component from the number of correct components.

4 Complete Response

Three parts essentially correct

3 Substantial Response

Two parts essentially correct and one part partially correct

2 Developing Response

Two parts essentially correct and no parts partially correct

OR

One part essentially correct and one or two parts partially correct

OR

Three parts partially correct

1 Minimal Response

One part essentially correct

OR

No parts essentially correct and two parts partially correct

AP[®] Statistics 2015 Scoring Guidelines

© 2015 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.

Visit the College Board on the Web: www.collegeboard.org.

AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

0

2015 SCORING GUIDELINES**Question 1****Intent of Question**

The primary goals of this question were to assess a student's ability to (1) compare features of two distributions of data displayed in boxplots and (2) identify statistical measures that are important in making decisions based on data sets.

Solution**Part (a):**

The median salary is approximately the same for both corporations. The range and interquartile range of the salaries are greater for Corporation A than for Corporation B. The two highest salaries at Corporation A are outliers while Corporation B has no outliers.

Part (b):

- (i) Five years after starting, at least 3 out of 30 (10%) of the salaries at Corporation A are greater than the maximum salary at Corporation B. If I accept the offer from Corporation A, I might be able to make a higher salary at Corporation A than at Corporation B.
- (ii) Five years after starting, the minimum salary at Corporation B is greater than at Corporation A. In fact, at Corporation A it looks like some people are still making the starting salary of \$36,000 and never received a raise in the five years since they were hired. So if I work at Corporation A, I might never receive a raise in salary.

Scoring

Parts (a) and (b) are scored as essentially correct (E), partially correct (P), or incorrect (I).

Part (a) is scored as follows:

Essentially correct (E) if the response includes the following four components:

1. A correct comparison of center.
2. A correct comparison of spread.
3. A discussion of the outliers for Corporation A.
4. The response is in context.

Partially correct (P) if the response includes only three of the four components.

Incorrect (I) if the response includes at most two of the four components.

Note: Any mention of shape should be ignored because complete shape information cannot be determined from a boxplot.

2015 SCORING GUIDELINES**Question 1 (continued)**

Part (b) is scored as follows:

Essentially correct (E) if the response includes the following four components:

1. In part (b-i) a relevant statistical measure is identified (or described) or a relevant statistical comparison is provided that supports the choice of Corporation A.
2. In part (b-i) an explanation is provided for why the measure or comparison is relevant.
3. In part (b-ii) a relevant statistical measure is identified (or described) or a relevant statistical comparison is provided that supports the choice of Corporation B.
4. In part (b-ii) an explanation is provided for why the measure or comparison is relevant.

Partially correct (P) if the response includes only two or three of the four components.

Incorrect (I) if the response includes none or one of the four components.

Note: If a response does not provide a statistical measure or comparison in part (b-i) or (b-ii), the second and fourth components can still be satisfied if an acceptable explanation is provided that would follow from a relevant statistical measure or comparison. For example, if the response in part (b-i) only states “At Corporation A, I have the potential to earn a higher salary,” the second component is satisfied.

4 Complete Response

Both parts essentially correct

3 Substantial Response

One part essentially correct and one part partially correct

2 Developing Response

One part essentially correct and one part incorrect

OR

Both parts partially correct

1 Minimal Response

One part partially correct and one part incorrect

2015 SCORING GUIDELINES

Question 2

Intent of Question

The primary goals of this question were to assess a student's ability to (1) use confidence intervals to test a question about a proportion and (2) understand the relationship between sample size and margin of error in a confidence interval for a proportion.

Solution**Part (a):**

- (i) No. The confidence interval is $(0.09, 0.21)$, which includes the value of 0.20. Therefore, it is plausible that the computer program is generating discounts with a probability of 0.20, and the confidence interval does not provide convincing statistical evidence that the program is not working as intended.
- (ii) No. The confidence interval includes values from 0.09 to 0.21, so any value in that interval is a plausible value for the probability that the computer is using to generate discounts.

Part (b):

The formula for computing the margin of error for a proportion includes the square root of the sample size in the denominator. For a random sample that is four times the size of the original sample, the margin of error can be determined by dividing the margin of error of the original sample by two. Therefore, the new margin of error is 0.03.

Part (c):

Using the margin of error of 0.03 obtained from the second sample, the confidence interval for p is 0.15 ± 0.03 or $(0.12, 0.18)$. The interval does not include 0.20, and therefore, there is convincing evidence that the computer program is not working as intended and is not generating discounts with a probability of 0.20.

Scoring

This question is scored in four sections. Section 1 consists of part (a-i); section 2 consists of part (a-ii); section 3 consists of part (b); and section 4 consists of part (c). Sections 1, 2, 3, and 4 are scored as essentially correct (E), partially correct (P), or incorrect (I).

Section 1 is scored as follows:

Essentially correct (E) if the response states that because the interval contains 0.20, it does not provide convincing statistical evidence that the computer program is not working as intended.

Partially correct (P) if the response indicates that it is necessary to check whether the value of 0.20 is in the computed interval, but there are errors in implementation. Examples of errors include:

- The response notes that 0.20 is within the interval but does not draw a conclusion.
- The response has an arithmetic error in the computation of the endpoints of the interval but provides a correct conclusion with justification that is consistent with the computed interval.

2015 SCORING GUIDELINES**Question 2 (continued)**

Incorrect (I) if the response does not recognize how to use the confidence interval to check whether the computer is working correctly;

OR

if the response states that the interval shows that the proportion is equal to 0.20;

OR

if the response notes that 0.20 is within the interval and concludes that the program is not working as intended;

OR

if the response otherwise does not meet the criteria for E or P.

Section 2 is scored as follows:

Essentially correct (E) if the response concludes that there is not convincing statistical evidence that the computer program generates the discount with a probability of 0.20 *AND* justifies the conclusion by noting that there are values other than 0.20 in the interval.

Partially correct (P) if the response correctly concludes that there is not convincing evidence that the computer program generates the discount with a probability of 0.20, but provides incomplete reasoning to justify the conclusion.

Examples of incomplete reasoning include:

- stating that 0.20 is a plausible value for the proportion of discounts without giving further explanation;
- indicating that having 0.20 in the interval does not prove that 0.20 is the true proportion; and
- providing a generic statistical argument, such as reference to the fact that the null hypothesis should never be accepted, or stating that not rejecting the null hypothesis is not proof that 0.20 is the true proportion of bills discounted.

Note: If an incorrect interval is computed in part (a-i) that does not contain 0.20, and in part (a-ii) the response concludes that the program is not generating discounts with a probability of 0.20 because 0.20 is not in the interval, section 2 is scored as P.

Incorrect (I) if the response states that there is evidence that the computer program generates the discount with a probability of 0.20;

OR

if the response correctly concludes that there is not convincing evidence that the computer program generates the discount with a probability of 0.20 *AND* provides incorrect or no justification;

OR

if the response otherwise does not meet the criteria for E or P.

Section 3 is scored as follows:

Essentially correct (E) if the response gives the correct value of 0.03 as the new margin of error by using the correct formula or by providing a correct explanation that recognizes that quadrupling the sample size divides the margin of error from the original sample by two.

Note: If the response relies on the margin of error formula and calculates a value that would round to 0.030, the response is scored as E.

2015 SCORING GUIDELINES

Question 2 (continued)

Partially correct (P) if the response uses the correct margin of error formula or provides a correct explanation that recognizes that quadrupling the sample size divides the margin of error from the original sample by two, but calculates an incorrect margin of error that is less than 0.06;

OR

if the response does not calculate a margin of error but provides a correct explanation that recognizes that quadrupling the sample size divides the margin of error from the original sample by two;

OR

if the response gives the correct margin of error without correct justification.

Incorrect (I) if the response does not recognize in any way that the new margin of error depends on the square root of the sample size;

OR

if the response calculates a margin of error that is greater than or equal to 0.06;

OR

if the response otherwise does not meet the criteria for E or P.

Section 4 is scored as follows:

Essentially correct (E) if the conclusion states that there is convincing evidence that the computer program is not working as intended because, based on the new margin of error, the interval does not contain 0.20;

OR

if the conclusion states the intended value of 0.20 is greater than the upper boundary of 0.18;

OR

if the conclusion states the margin of error is smaller than the difference between the sample proportion and the intended long-run proportion of 0.20.

Notes:

- If the margin of error was computed incorrectly in part (b), but a correct answer to part (c) is consistent with this incorrect margin of error, section 4 is scored as E.
- If no specific margin of error or confidence interval was given in parts (b) or (c), but the response provides a complete and correct conclusion with reference to a smaller margin of error (or narrower confidence interval), section 4 is scored as E.

Partially correct (P) if an interval is incorrectly constructed using the margin of error from part (b), but a correct conclusion is given and justified for the computed interval;

OR

if an interval is computed correctly using the margin of error from part (b) and an argument is made based on whether or not 0.20 is in the interval, but the conclusion is incorrect;

OR

if no specific margin of error or confidence interval has been given in parts (b) or (c), but the response concludes that because the margin of error has decreased (or the confidence interval is narrower), 0.20 is not in the interval.

Incorrect (I) if an interval is computed correctly using the margin of error from part (b) and a correct conclusion is given, but no argument is made based on whether or not 0.20 is in the interval;

OR

if the response otherwise does not meet the criteria for E or P.

2015 SCORING GUIDELINES**Question 2 (continued)**

Notes:

- If a response includes a confidence level, the level can be ignored because no confidence level was provided.
- If the response bases a conclusion on the relative location of 0.20 within the interval (for example, 0.20 is near the edge of the interval), the response is scored as I.
- A response that provides additional incorrect explanation lowers the score in section 4 by one level (that is, from E to P, or from P to I).

Each essentially correct (E) section counts as 1 point. Each partially correct (P) section counts as $\frac{1}{2}$ point.

- | | |
|----------|-----------------------------|
| 4 | Complete Response |
| 3 | Substantial Response |
| 2 | Developing Response |
| 1 | Minimal Response |

If a response is between two scores (for example, $2\frac{1}{2}$ points), use a holistic approach to decide whether to score up or down, depending on the overall strength of the response and communication.

SCORING GUIDELINES**Question****Intent of Question**

The primary goals of this question were to assess a student's ability to (1) perform a probability calculation from a discrete random variable; (2) calculate the expected value of a discrete random variable; (3) perform a conditional probability calculation from a discrete random variable; and (4) use probabilistic thinking to make a prediction about how an expected value will change given a condition about the random variable.

Solution**Part (a):**

The probability that at least one ATM is working when the mall opens is:

$$P(X \geq 1) = 0.21 + 0.40 + 0.24 = 0.85.$$

Part (b):

The expected value of the number of ATMs that are working when the mall opens is:

$$E(X) = 0(0.15) + 1(0.21) + 2(0.40) + 3(0.24) = 1.73 \text{ machines.}$$

Part (c):

probability that all three ATMs are working when the mall opens, given that at least one ATM is working is:

$$P(X = 3 \mid X \geq 1) = \frac{P(X = 3 \text{ and } X \geq 1)}{P(X \geq 1)} = \frac{P(X = 3)}{P(X \geq 1)} = \frac{0.24}{0.85} \approx 0.282$$

Part (d):

Given that at least one ATM is working when the mall opens, the expected value of the number of working ATMs would be greater than the expected value calculated in part (b). By eliminating the possibility of 0 working ATMs, the probabilities for 1, 2, and 3 working ATMs all increase proportionally, so the expected value must increase.

Scoring

(b), (c), and (d) are scored as essentially correct (E), partially correct (P), or incorrect (I).

Part (a) is scored as follows:

Essentially correct (E) if the probability is computed correctly with work shown.

Partially correct (P) if the correct answer is given, but no work is shown;

OR

appropriate work is shown but the answer is incorrect or missing;

OR

SCORING GUIDELINES

Question 3 (continued)

one of the incorrect cumulative probabilities $P(X < 1)$, $P(X \leq 1)$, or $P(X > 1)$ is stated AND computed correctly.

Incorrect (I) if the response does not meet the criteria for E or P.

Note: The probability can be calculated as $1 - P(X = 0) = 1 - 0.15 = 0.85$.

Part (b) is scored as follows:

Essentially correct (E) if the expected value is computed correctly with work shown.

Partially correct (P) if the correct answer is given, but no work is shown;

OR

if appropriate work is shown but the answer is incorrect or missing.

Incorrect (I) if the response does not meet the criteria for E or P.

Part (c) is scored as follows:

Essentially correct (E) if the probability is computed correctly, with work shown that includes correct numerical values for both the numerator and denominator.

Partially correct (P) if the response includes a numerator and denominator in calculating the conditional probability, with one (numerator or denominator) correct in numerical value and the other incorrect;

OR

if the correct answer is given, but no work is shown.

Incorrect (I) if the response does not meet the criteria for E or P.

Part (d) is scored as follows:

Essentially correct (E) if the response provides the correct answer (greater than) with a reasonable explanation based on the fact that with $X = 0$ eliminated, the probabilities for $X = 1$, $X = 2$, $X = 3$ all increase;

OR

if the response provides the correct answer (greater than) with a reasonable explanation based on fact that with $X = 0$ eliminated, the balance point of the distribution increases;

OR

if the response provides the correct answer (greater than) and the conditional expected value is computed correctly with work shown.

Partially correct (P) if the response provides the correct answer (greater than) with a weak explanation, such as "Yes, because 0 is eliminated";

OR

SCORING GUIDELINES**Question 3 (continued)**

if the response provides the correct answer (greater than) with a reasonable but incorrect attempt to calculate the conditional expected value using a revised probability distribution with only the values $X = 1$, $X = 2$, and $X = 3$, and corresponding probabilities greater than or equal to those in the original probability distribution.

Incorrect (I) if the response does not provide the correct answer (greater than);

OR

if the response provides the correct answer (greater than) with an incorrect explanation or no explanation.

Note: The conditional expected value is:

$$1\left(\frac{0.21}{0.85}\right) + 2\left(\frac{0.40}{0.85}\right) + 3\left(\frac{0.24}{0.85}\right) \approx 1(0.247) + 2(0.471) + 3(0.282) = 2.04 \text{ machines.}$$

Each essentially correct (E) part counts as 1 point. Each partially correct (P) part counts as $\frac{1}{2}$ point.

- 4 Complete Response**
- 3 Substantial Response**
- 2 Developing Response**
- 1 Minimal Response**

If a response is between two scores (for example, $2\frac{1}{2}$ points), use a holistic approach to decide whether score up or down, depending on the overall strength of the response and communication.

2015 SCORING GUIDELINES

Question 4

Intent of Question

The primary goal of this question was to assess a student's ability to identify, set up, perform, and interpret the results of an appropriate hypothesis test to address a particular question. More specific goals were to assess a student's ability to (1) state appropriate hypotheses; (2) identify the appropriate statistical test procedure and check appropriate conditions for inference; (3) calculate the appropriate test statistic and p -value; and (4) draw an appropriate conclusion, with justification, in the context of the study.

Solution

Step 1: States a correct pair of hypotheses.

Let p_{asp} represent the population proportion of adults similar to those in the study who would have developed colon cancer within the six years of the study if they had taken a low-dose aspirin each day. Similarly, let p_{plac} represent the population proportion of adults similar to those in the study who would have developed colon cancer within the six years of the study if they had taken a placebo each day.

The hypotheses to be tested are $H_0 : p_{\text{asp}} = p_{\text{plac}}$ versus $H_a : p_{\text{asp}} < p_{\text{plac}}$ or equivalently, $H_0 : p_{\text{asp}} - p_{\text{plac}} = 0$ versus $H_a : p_{\text{asp}} - p_{\text{plac}} < 0$.

Step 2: Identifies a correct test procedure (by name or by formula) and checks appropriate conditions.

The appropriate procedure is a two-sample z -test for comparing proportions.

Because this is a randomized experiment, the first condition is that the volunteers were randomly assigned to one treatment group or the other. The condition is satisfied because we are told that the volunteers were randomly assigned to take a low-dose aspirin or a placebo.

The second condition is that the sample sizes are large, relative to the proportions involved. The condition is satisfied because all sample counts are large enough; that is, 15 with colon cancer in aspirin group, 26 with colon cancer in placebo group, $500 - 15 = 485$ cancer-free in aspirin group, and $500 - 26 = 474$ cancer-free in placebo group.

Step 3: Calculates the appropriate test statistic and p -value.

The sample proportions who developed colon cancer are $\hat{p}_{\text{asp}} = \frac{15}{500} = 0.030$ and $\hat{p}_{\text{plac}} = \frac{26}{500} = 0.052$.

The combined sample proportion who developed colon cancer is $\hat{p}_{\text{combined}} = \frac{15 + 26}{500 + 500} = 0.041$.

The test statistic is $z = \frac{0.030 - 0.052}{\sqrt{0.041(1 - 0.041)\left(\frac{1}{500} + \frac{1}{500}\right)}} \approx -1.75$ (-1.7542 from calculator).

The p -value is $P(Z \leq -1.75) = 0.0401$ (0.0397 from calculator), where Z has a standard normal distribution.

2015 SCORING GUIDELINES

Question 4 (continued)

Step 4: States a correct conclusion in the context of the study, using the result of the statistical test.

Because the p -value is less than the given significance level of $\alpha = 0.05$, we reject the null hypothesis. The data provide convincing statistical evidence that the proportion of all adults similar to the volunteers who would develop colon cancer if they had taken a low-dose aspirin every day is less than the proportion of all adults similar to the volunteers who would develop colon cancer if they had not taken a low-dose aspirin every day.

Scoring

Steps 1, 2, 3, and 4 are scored as essentially correct (E), partially correct (P), or incorrect (I).

Step 1 is scored as follows:

Essentially correct (E) if the response identifies correct parameters *AND* both hypotheses are labeled and state the correct relationship between the parameters.

Partially correct (P) if the response identifies correct parameters *OR* states correct relationships, but not both.

Incorrect (I) if the response does not meet the criteria for E or P.

Note: Either defining the parameters in context, or simply using common parameter notation with subscripts clearly relevant to the context, such as p_{asp} and p_{plac} , is sufficient.

Step 2 is scored as follows:

Essentially correct (E) if the response correctly includes the following three components:

1. Identifies the correct test procedure (by name or by formula).
2. Notes that the use of random assignment satisfies the randomness condition.
3. Checks for approximate normality of the test statistic by citing that all four counts are larger than some standard criterion such as 5 or 10.

Partially correct (P) if the response correctly includes only two of the three components.

Incorrect (I) if the response correctly includes at most one of the three components.

Notes:

- For the randomness component, it is (minimally) acceptable to say “random assignment — check” but not acceptable to say “random — check” or “SRS — check.” The important concept here is that it is random assignment, and not random sampling, that is required. If the response implies that the study used a random sample, the randomness component is not satisfied, regardless of whether random assignment is correctly addressed.
- The normality check may use the expected counts under the null hypothesis in place of observed counts.

2015 SCORING GUIDELINES

Question 4 (continued)

Step 3 is scored as follows:

Essentially correct (E) if the response correctly calculates both the test statistic and a p -value that is consistent with the stated alternative hypothesis.

Partially correct (P) if the response correctly calculates the test statistic but not the p -value;

OR

if the response calculates the test statistic incorrectly but then calculates the correct p -value for the computed test statistic;

OR

if the response reports the correct p -value but no calculations or test statistic are shown.

Incorrect (I) if the response fails to meet the criteria for E or P.

Note: The p -value is considered correct if it is consistent with the alternative stated in the response and the calculated test statistic, even if those are incorrect.

Step 4 is scored as follows:

Essentially correct (E) if the response provides a correct conclusion in context, with justification based on linkage between the p -value and the given $\alpha = 0.05$.

Partially correct (P) if the response provides a correct conclusion, with linkage to the p -value, but not in context;

OR

if the response provides a correct conclusion in context, but without justification based on linkage to the p -value.

Incorrect (I) if the response does not meet the criteria for E or P.

Notes:

- The conclusion must be related to the alternative hypothesis.
- If the p -value is incorrect, then step 4 is scored as E if the response includes proper linkage and a conclusion in context consistent with that p -value.
- If the p -value is less than 0.05, wording that states or implies that the alternative hypothesis is *proven* lowers the score one level (that is, from E to P or P to I) in step 4.
- If the p -value is incorrect and greater than 0.05, wording that states or implies that the null hypothesis is *accepted* lowers the score one level (that is, from E to P or P to I) in step 4.

2015 SCORING GUIDELINES**Question 4 (continued)**

Each essentially correct (E) step counts as 1 point. Each partially correct (P) step counts as $\frac{1}{2}$ point.

- 4 Complete Response**
- 3 Substantial Response**
- 2 Developing Response**
- 1 Minimal Response**

If a response is between two scores (for example, $2\frac{1}{2}$ points), use a holistic approach to decide whether to score up or down, depending on the overall strength of the response and communication.

SCORING GUIDELINES

Question

Intent of Question

The primary goals of this question were to assess a student's ability to (1) use the information provided by a scatterplot to describe the relationship between two quantitative variables; (2) interpret and use the information given by lines displayed on a scatterplot; and (3) use a regression equation to estimate a predicted value of y for a given x value.

Solution**Part**

There is a moderately strong, positive, linear relationship between height and arm span so that taller students tend to have longer arm spans.

Part (b):

The line in Graph 2 is the one that is helpful. For each student, the graph illustrates whether arm is equal to height (points on the line), arm span is greater than height (points above the line), or arm span is less than height (points below the line).

Classification	Square	Tall Rectangle	Short Rectangle
Frequency	3	4	5

Part (c):

The predicted arm span is $\hat{y} = 11.74 + 0.8247x = 11.74 + 0.8247(61) = 62.05$ inches.

Scoring

b), and (c) are scored as essentially correct (E), partially correct (P), or incorrect (I).

Part (a) is scored as follows:

Essentially correct (E) if the response includes the following four components:

1. The relationship is approximately linear.
2. The relationship is positive.
3. There is a moderate to strong association (or relationship).
4. The response is given in context.

Partially correct (P) if the response includes only two or three of the four components.

Incorrect (I) if the response includes at most one of the four components.

SCORING GUIDELINES

Question 5 (continued)

Notes:

- Correlation alone is not sufficient to indicate a linear relationship. However, linear correlation sufficient to indicate a linear relationship.
- Reporting that the correlation coefficient is 0.81, the correlation coefficient is close to 1.0, or some other value or range of values for the correlation coefficient is not sufficient to satisfy component 3.

Part (b) is scored as follows:

Essentially correct (E) if the response contains the following four components:

- The line in Graph 2 is selected.
- A reasonable explanation for selecting the line in Graph 2 that links the body shapes to the regions defined by the line is given. The explanation should indicate that square shapes (arm span is equal to height) are represented by points on the line, tall rectangle shapes (arm span less than height) are represented by points below the line, and short rectangle shapes (arm span is greater than height) are represented by points above the line.
- Correct counting of body types demonstrated by reporting correct frequencies (3, 4, and 5) reporting correct proportions $\left(\frac{1}{4} = \frac{3}{12}, \frac{1}{3} = \frac{4}{12}, \text{ and } \frac{5}{12}\right)$ for the square, tall rectangle, and short rectangle body types, respectively.
- The correct frequencies (3, 4, and 5) are reported in the table.

Partially correct (P) if the response includes only two or three of the four components.

Incorrect (I) if the response includes at most one of the four components.

Notes:

- Selecting the regression line on Graph 1 cannot satisfy either component 1 or component 2.
- To satisfy component 1, it is sufficient to refer to the $y = x$ line without explicitly mentioning Graph 2.
- Use of incorrect labels, such as regression line or least-squares regression line, in referring to the $y = x$ line in Graph 2 is an incorrect use of terminology that should be strongly discouraged, but it is ignored in this scoring rubric.
- The explanation for selecting Graph 2 is acceptable if it explicitly includes at least two of the following: (i) square body shapes (arm span equal to height) are represented by points on the line, (ii) tall rectangle body shapes (arm span less than height) are represented by points *below* the line, or (iii) short rectangle body shapes (arm span greater than height) are represented by points *above* the line.
- Incorrectly counting the points on the $y = x$ line in Graph 2 is considered a minor error. Frequencies (2, 4, and 5) are accepted for both component 3 and component 4. Reporting corresponding proportions $\left(\frac{2}{11}, \frac{4}{11}, \frac{5}{11}\right)$ or $\left(\frac{2}{12}, \frac{4}{12}, \frac{5}{12}\right)$ in the table satisfies component 3 but does not satisfy component 4.
- Frequencies reported in the table as (2, 5, 4) or (3, 5, 4), interchanging the counts for tall and short rectangle body shapes, satisfies component 3, but does not satisfy component 4. Reporting corresponding proportions, $\left(\frac{2}{11}, \frac{5}{11}, \frac{4}{11}\right)$ or $\left(\frac{3}{12}, \frac{5}{12}, \frac{4}{12}\right)$, in the table does not satisfy either component 3 or component 4.

SCORING GUIDELINES

Question 5 (continued)

Part (c) is scored as follows:

Essentially correct (E) if the response contains the following three components:

1. Correct formula for predicting arm span with 61 inserted for x .
2. Correct value for the predicted arm span.
3. Units for the predicted arm span given as inches.

Partially correct (P) if the response includes only two of the three components.

Incorrect (I) if the response includes at most one of the three components.

Notes:

- Any value for the predicted arm span between 61.5 and 62.5 is acceptable; values outside this interval do not satisfy component 2. This allows for inaccuracy in obtaining a value for the predicted arm span directly from the line displayed on Graph 1 and for rounding in applying the prediction formula; but it excludes predictions based on the $y = x$ line and other unreasonable predictions.
- Reporting a prediction equation that is similar to the equation given in the stem, with 61 inserted for x , satisfies components 1 and 2 if the value of the predicted arm span is between 61.5 and 62.5 inches. Otherwise, neither of those two components is satisfied. This could occur if a student enters the data from the graph into a calculator to compute the least squares regression line, instead of using the formula provided in the stem.

4 Complete Response

All three parts essentially correct

3 Substantial Response

Two parts essentially correct and one part partially correct

2 Developing Response

Two parts essentially correct and one part incorrect

OR

One part essentially correct and two parts partially correct

OR

One part essentially correct, one part partially correct, and one part incorrect

OR

Three parts partially correct

1 Minimal Response

One part essentially correct and two parts incorrect

OR

Two parts partially correct and one part incorrect

2015 SCORING GUIDELINES

Question 6

Intent of Question

The primary goals of this question were to assess a student's ability to (1) describe how sample data would differ using two different sampling methods; (2) describe the sampling distribution of the sample mean for two different sampling methods; and (3) choose the sampling method that will result in the best estimate of the population mean.

Solution**Part (a):**

No, a sample obtained using Method 2 will not be representative of all tortillas made that day. The sample obtained using Method 2 will only represent the tortillas from one production line, not from the entire population because the distributions of diameters for the two production lines are different.

Part (b):

Method 1 was most likely used to select this sample. The bimodal shape in the histogram of sample data indicates that tortillas were selected from both production lines, which is what would happen using Method 1. Method 2 would be likely to produce a unimodal distribution of diameters centered at either 5.9 inches or 6.1 inches.

Part (c):

Method 2 would result in less variability in the sample of 200 tortillas on a given day because the sample comes from only one production line. Because the distributions of diameters are not the same for the two production lines, selecting tortillas from both lines as in Method 1 would result in more variable sample data.

Part (d):

The sampling distribution of the sample mean diameter for samples obtained using Method 1 would be approximately normal with mean 6 inches and standard deviation $\frac{0.11}{\sqrt{200}} \approx 0.0078$ inch.

Part (e):

Method 1 would result in less variability in the sample means over the 365 days, because with Method 2, roughly half of the sample means will be around 5.9 inches and the other half will be around 6.1 inches. With Method 1, however, the sample means will all be very close to 6 inches, as indicated by the standard deviation in part (d).

Part (f):

Method 1 is more likely to produce a sample mean close to 6 inches. Because the sample mean is an unbiased estimator for both methods, the manager should pick the method that would result in less variability in the distribution of the sample mean. Based on the answer to part (e), Method 1 results in less variability in the distribution of the sample mean.

2015 SCORING GUIDELINES

Question 6 (continued)

Scoring

This question is scored in three sections. Section 1 consists of parts (a), (b), and (c), section 2 consists of part (d), and section 3 consists of parts (e) and (f). Sections 1, 2, and 3 are scored as essentially correct (E), partially correct (P), or incorrect (I).

Section 1 is scored as follows:

Essentially correct (E) if the response includes the following three components:

1. In part (a) the response says no *AND* either argues that the sample will only be selected from one production line and not the entire population (that is, the sample will only represent one production line) *OR* argues that the tortillas from the two productions lines are different.
2. In part (b) the response chooses Method 1 *AND* refers to a relevant characteristic of the histogram (shape, center, or variability) that matches what would be expected when using Method 1 (or that does not match what would be expected when using Method 2).
3. In part (c) the response chooses Method 2 *AND* either justifies by stating that the sample comes from only one production line (does not come from both production lines) *OR* justifies by comparing the possible range of diameters for the two methods.

Partially correct (P) if the response includes only two of the three components.

Incorrect (I) if the response includes at most one of the three components.

Note: If a response includes more than one justification in an individual part, score the weaker of the two justifications. For example, a response for part (a) that says “No, because only one line was selected and because the sample size was too small” does not satisfy the first component because the sample size argument is incorrect.

Section 2 is scored as follows:

Essentially correct (E) if the response includes the following three components:

1. The shape is approximately normal.
2. The mean is 6 inches.
3. The standard deviation is $\frac{0.11}{\sqrt{200}}$ inch.

Partially correct (P) if the response includes only two of the three components.

Incorrect (I) if the response includes at most one of the three components.

Note: It is not necessary to include units (inches) for the mean or standard deviation.

Section 3 is scored as follows:

Essentially correct (E) if the response includes the following two components:

1. In part (e) the response chooses Method 1 *AND* describes the sampling distribution of the sample mean for Method 2 as having some *sample means* close to the mean of production line A (5.9 inches) and the other *sample means* close to the mean of production line B (6.1 inches).

2015 SCORING GUIDELINES

Question 6 (continued)

2. In part (f) the response chooses Method 1 *AND*

- refers to a correct answer in part (e);

OR

- describes the sampling distribution of the sample mean for Method 2 as having some *sample means* close to the mean of production line A (5.9, less than 6) and the other *sample means* close to the mean of production line B (6.1, greater than 6);

OR

- argues that on a single day, it would be preferable to get a sample with a mean around 6 rather than getting a sample that would have a mean around 5.9 (less than 6) or a mean around 6.1 (greater than 6).

Partially correct (P) if the response includes one of the two components.

Incorrect (I) if the response does not meet the criteria for E or P.

Notes:

- In part (e), the response must be clear that there is *more than one* mean being described (365 sample means).
 - Correct: The sample means will be around 5.9 or 6.1.
 - Not correct: The sample mean will be around 5.9 or 6.1.
- Parts (e) and (f) are not satisfied if the response does not imply the sample means *vary* from the population means in both methods (or does not imply that the sample mean varies from the population mean when making the single day argument in part (f)). However, if the mistake is made in part (e), do not penalize the response in part (f) for the same mistake.
 - Correct: The sample means will be around 5.9 or 6.1 instead of close to 6.
 - Not correct: The sample means will be 5.9 or 6.1 instead of close to 6.
 - Not correct: The sample means will be 6 instead of around 5.9 or 6.1.

4 Complete Response

All three sections essentially correct

3 Substantial Response

Two sections essentially correct and one section partially correct

2 Developing Response

Two sections essentially correct and one section incorrect

OR

One section essentially correct and either one or two sections partially correct

OR

All three sections partially correct

1 Minimal Response

One section essentially correct and two sections incorrect

OR

Two sections partially correct and one section incorrect