

Begin your response to **QUESTION 6** on this page.

SECTION II, Part B

Suggested Time—25 minutes


1 Question

Directions: Show all your work. Indicate clearly the methods you use, because you will be scored on the correctness of your methods as well as on the accuracy and completeness of your results and explanations.

6. A company sells a certain type of whistle. The price of the whistle varies from store to store. Julio, a statistician at the company, wants to estimate the mean price, in dollars (\$), of this type of whistle at all stores that sell the whistle.

- (a) (i) Identify the appropriate inference procedure for Julio to use.
- (ii) Describe the parameter for the inference procedure you identified in part (a-i) in context.

Julio called the managers of 20 randomly selected stores that sell the whistle and recorded the price of the whistle at each store. Following is a dotplot of Julio’s data.



GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 6** on this page.

The summary statistics for Julio’s data are shown in the following table.

Summary Statistics for Julio’s Data

Sample Size	Mean	Standard Deviation	Minimum	Q_1	Median	Q_3	Maximum
20	5.12	0.743	4.25	4.51	4.885	5.475	6.58

- (b) Julio wants to examine some characteristics of the distribution of the sample of whistle prices.
- (i) Describe the shape of the distribution of the sample of whistle prices. Justify your response using appropriate values from the summary statistics table.
- (ii) Using the $1.5 \times \text{IQR}$ rule, determine whether there are any outliers in the sample of whistle prices. Justify your response.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 6** on this page.

It can often be difficult to determine whether the distribution of sample data is skewed by looking at a graph of the data and the summary statistics, particularly when the sample size is small. Thus, statisticians sometimes measure how skewed a data set is. One such measure is Pearson's coefficient of skewness, which is calculated using the following formula.


$$\text{Pearson's Coefficient of Skewness} = \frac{3(\bar{x} - m)}{s}$$

In the formula, $\bar{x}$ is the sample mean, m is the sample median, and s is the sample standard deviation.

(c) (i) Calculate Pearson's coefficient of skewness for Julio's sample of 20 whistle prices. Show your work.

The following graph shows conclusions that can be made about the shape of the distribution of sample data based on Pearson's coefficient of skewness and sample size.

Conclusion from Pearson's Coefficient of Skewness



(ii) Indicate the value of the Pearson's coefficient of skewness you calculated in part (c-i) for the appropriate sample size by marking it with an "X" on the preceding graph.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 6** on this page.

(d) Consider your work in part (c).

(i) What should you conclude about the shape of the distribution of the sample of whistle prices? Justify your response.

Julio's inference procedure in part (a-i) needs one of the following requirements to be satisfied to verify the normality condition.

- The sample size is greater than or equal to 30.
- If the sample size is less than 30, the distribution of the sample data is not strongly skewed and does not have outliers.

(ii) Using your response to (d-i) and the preceding requirements, is the normality condition satisfied for Julio's data? Explain your response.

GO ON TO THE NEXT PAGE.

STOP

END OF EXAM




Table entry for z is the probability lying below z .

Table A Standard normal probabilities

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
−3.4	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0002
−3.3	.0005	.0005	.0005	.0004	.0004	.0004	.0004	.0004	.0004	.0003
−3.2	.0007	.0007	.0006	.0006	.0006	.0006	.0006	.0005	.0005	.0005
−3.1	.0010	.0009	.0009	.0009	.0008	.0008	.0008	.0008	.0007	.0007
−3.0	.0013	.0013	.0013	.0012	.0012	.0011	.0011	.0011	.0010	.0010
−2.9	.0019	.0018	.0018	.0017	.0016	.0016	.0015	.0015	.0014	.0014
−2.8	.0026	.0025	.0024	.0023	.0023	.0022	.0021	.0021	.0020	.0019
−2.7	.0035	.0034	.0033	.0032	.0031	.0030	.0029	.0028	.0027	.0026
−2.6	.0047	.0045	.0044	.0043	.0041	.0040	.0039	.0038	.0037	.0036
−2.5	.0062	.0060	.0059	.0057	.0055	.0054	.0052	.0051	.0049	.0048
−2.4	.0082	.0080	.0078	.0075	.0073	.0071	.0069	.0068	.0066	.0064
−2.3	.0107	.0104	.0102	.0099	.0096	.0094	.0091	.0089	.0087	.0084
−2.2	.0139	.0136	.0132	.0129	.0125	.0122	.0119	.0116	.0113	.0110
−2.1	.0179	.0174	.0170	.0166	.0162	.0158	.0154	.0150	.0146	.0143
−2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183
−1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
−1.8	.0359	.0351	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
−1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
−1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
−1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
−1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0721	.0708	.0694	.0681
−1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
−1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
−1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
−1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
−0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
−0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
−0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
−0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
−0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
−0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
−0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
−0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
−0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
−0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641




Table entry for z is the probability lying below z .

Table A (Continued)

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
0.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
0.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
0.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
0.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
0.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
0.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
0.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
0.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
0.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.9997
3.4	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9998




Table entry for p and C is the point t^* with probability p lying above it and probability C lying between $-t^*$ and t^* .

Table B t distribution critical values

df	Tail probability p									
	.25	.20	.15	.10	.05	.025	.02	.01	.005	.0025
1	1.000	1.376	1.963	3.078	6.314	12.71	15.89	31.82	63.66	127.3
2	.816	1.061	1.386	1.886	2.920	4.303	4.849	6.965	9.925	14.09
3	.765	.978	1.250	1.638	2.353	3.182	3.482	4.541	5.841	7.453
4	.741	.941	1.190	1.533	2.132	2.776	2.999	3.747	4.604	5.598
5	.727	.920	1.156	1.476	2.015	2.571	2.757	3.365	4.032	4.773
6	.718	.906	1.134	1.440	1.943	2.447	2.612	3.143	3.707	4.317
7	.711	.896	1.119	1.415	1.895	2.365	2.517	2.998	3.499	4.029
8	.706	.889	1.108	1.397	1.860	2.306	2.449	2.896	3.355	3.833
9	.703	.883	1.100	1.383	1.833	2.262	2.398	2.821	3.250	3.690
10	.700	.879	1.093	1.372	1.812	2.228	2.359	2.764	3.169	3.581
11	.697	.876	1.088	1.363	1.796	2.201	2.328	2.718	3.106	3.497
12	.695	.873	1.083	1.356	1.782	2.179	2.303	2.681	3.055	3.428
13	.694	.870	1.079	1.350	1.771	2.160	2.282	2.650	3.012	3.372
14	.692	.868	1.076	1.345	1.761	2.145	2.264	2.624	2.977	3.326
15	.691	.866	1.074	1.341	1.753	2.131	2.249	2.602	2.947	3.286
16	.690	.865	1.071	1.337	1.746	2.120	2.235	2.583	2.921	3.252
17	.689	.863	1.069	1.333	1.740	2.110	2.224	2.567	2.898	3.222
18	.688	.862	1.067	1.330	1.734	2.101	2.214	2.552	2.878	3.197
19	.688	.861	1.066	1.328	1.729	2.093	2.205	2.539	2.861	3.174
20	.687	.860	1.064	1.325	1.725	2.086	2.197	2.528	2.845	3.153
21	.686	.859	1.063	1.323	1.721	2.080	2.189	2.518	2.831	3.135
22	.686	.858	1.061	1.321	1.717	2.074	2.183	2.508	2.819	3.119
23	.685	.858	1.060	1.319	1.714	2.069	2.177	2.500	2.807	3.104
24	.685	.857	1.059	1.318	1.711	2.064	2.172	2.492	2.797	3.091
25	.684	.856	1.058	1.316	1.708	2.060	2.167	2.485	2.787	3.078
26	.684	.856	1.058	1.315	1.706	2.056	2.162	2.479	2.779	3.067
27	.684	.855	1.057	1.314	1.703	2.052	2.158	2.473	2.771	3.057
28	.683	.855	1.056	1.313	1.701	2.048	2.154	2.467	2.763	3.047
29	.683	.854	1.055	1.311	1.699	2.045	2.150	2.462	2.756	3.038
30	.683	.854	1.055	1.310	1.697	2.042	2.147	2.457	2.750	3.030
40	.681	.851	1.050	1.303	1.684	2.021	2.123	2.423	2.704	2.971
50	.679	.849	1.047	1.299	1.676	2.009	2.109	2.403	2.678	2.937
60	.679	.848	1.045	1.296	1.671	2.000	2.099	2.390	2.660	2.915
80	.678	.846	1.043	1.292	1.664	1.990	2.088	2.374	2.639	2.887
100	.677	.845	1.042	1.290	1.660	1.984	2.081	2.364	2.626	2.871
1000	.675	.842	1.037	1.282	1.646	1.962	2.056	2.330	2.581	2.813
∞	.674	.841	1.036	1.282	1.645	1.960	2.054	2.326	2.576	2.807
Confidence level C										
50% 60% 70% 80% 90% 95% 96% 98% 99% 99.5% 99.8% 99.9%										

Table entry for p is the point (χ^2) with probability p lying above it.




Table C χ^2 critical values

df	Tail probability p											
	.25	.20	.15	.10	.05	.025	.02	.01	.005	.0025	.001	.0005
1	1.32	1.64	2.07	2.71	3.84	5.02	5.41	6.63	7.88	9.14	10.83	12.12
2	2.77	3.22	3.79	4.61	5.99	7.38	7.82	9.21	10.60	11.98	13.82	15.20
3	4.11	4.64	5.32	6.25	7.81	9.35	9.84	11.34	12.84	14.32	16.27	17.73
4	5.39	5.99	6.74	7.78	9.49	11.14	11.67	13.28	14.86	16.42	18.47	20.00
5	6.63	7.29	8.12	9.24	11.07	12.83	13.39	15.09	16.75	18.39	20.51	22.11
6	7.84	8.56	9.45	10.64	12.59	14.45	15.03	16.81	18.55	20.25	22.46	24.10
7	9.04	9.80	10.75	12.02	14.07	16.01	16.62	18.48	20.28	22.04	24.32	26.02
8	10.22	11.03	12.03	13.36	15.51	17.53	18.17	20.09	21.95	23.77	26.12	27.87
9	11.39	12.24	13.29	14.68	16.92	19.02	19.68	21.67	23.59	25.46	27.88	29.67
10	12.55	13.44	14.53	15.99	18.31	20.48	21.16	23.21	25.19	27.11	29.59	31.42
11	13.70	14.63	15.77	17.28	19.68	21.92	22.62	24.72	26.76	28.73	31.26	33.14
12	14.85	15.81	16.99	18.55	21.03	23.34	24.05	26.22	28.30	30.32	32.91	34.82
13	15.98	16.98	18.20	19.81	22.36	24.74	25.47	27.69	29.82	31.88	34.53	36.48
14	17.12	18.15	19.41	21.06	23.68	26.12	26.87	29.14	31.32	33.43	36.12	38.11
15	18.25	19.31	20.60	22.31	25.00	27.49	28.26	30.58	32.80	34.95	37.70	39.72
16	19.37	20.47	21.79	23.54	26.30	28.85	29.63	32.00	34.27	36.46	39.25	41.31
17	20.49	21.61	22.98	24.77	27.59	30.19	31.00	33.41	35.72	37.95	40.79	42.88
18	21.60	22.76	24.16	25.99	28.87	31.53	32.35	34.81	37.16	39.42	42.31	44.43
19	22.72	23.90	25.33	27.20	30.14	32.85	33.69	36.19	38.58	40.88	43.82	45.97
20	23.83	25.04	26.50	28.41	31.41	34.17	35.02	37.57	40.00	42.34	45.31	47.50
21	24.93	26.17	27.66	29.62	32.67	35.48	36.34	38.93	41.40	43.78	46.80	49.01
22	26.04	27.30	28.82	30.81	33.92	36.78	37.66	40.29	42.80	45.20	48.27	50.51
23	27.14	28.43	29.98	32.01	35.17	38.08	38.97	41.64	44.18	46.62	49.73	52.00
24	28.24	29.55	31.13	33.20	36.42	39.36	40.27	42.98	45.56	48.03	51.18	53.48
25	29.34	30.68	32.28	34.38	37.65	40.65	41.57	44.31	46.93	49.44	52.62	54.95
26	30.43	31.79	33.43	35.56	38.89	41.92	42.86	45.64	48.29	50.83	54.05	56.41
27	31.53	32.91	34.57	36.74	40.11	43.19	44.14	46.96	49.64	52.22	55.48	57.86
28	32.62	34.03	35.71	37.92	41.34	44.46	45.42	48.28	50.99	53.59	56.89	59.30
29	33.71	35.14	36.85	39.09	42.56	45.72	46.69	49.59	52.34	54.97	58.30	60.73
30	34.80	36.25	37.99	40.26	43.77	46.98	47.96	50.89	53.67	56.33	59.70	62.16
40	45.62	47.27	49.24	51.81	55.76	59.34	60.44	63.69	66.77	69.70	73.40	76.09
50	56.33	58.16	60.35	63.17	67.50	71.42	72.61	76.15	79.49	82.66	86.66	89.56
60	66.98	68.97	71.34	74.40	79.08	83.30	84.58	88.38	91.95	95.34	99.61	102.7
80	88.13	90.41	93.11	96.58	101.9	106.6	108.1	112.3	116.3	120.1	124.8	128.3
100	109.1	111.7	114.7	118.5	124.3	129.6	131.1	135.8	140.2	144.3	149.4	153.2

Descriptive Statistics

$$\bar{x} = \frac{1}{n} \sum x_i = \frac{\sum x_i}{n}$$

$$\hat{y} = a + bx$$

$$r = \frac{1}{n-1} \sum \left(\frac{x_i - \bar{x}}{s_x} \right) \left(\frac{y_i - \bar{y}}{s_y} \right)$$

$$s_x = \sqrt{\frac{1}{n-1} \sum (x_i - \bar{x})^2} = \sqrt{\frac{\sum (x_i - \bar{x})^2}{n-1}}$$

$$\bar{y} = a + b\bar{x}$$

$$b = r \frac{s_y}{s_x}$$

Descriptive Statistics

$$\bar{x} = \frac{1}{n} \sum x_i = \frac{\sum x_i}{n}$$

$$\hat{y} = a + bx$$

$$r = \frac{1}{n-1} \sum \left(\frac{x_i - \bar{x}}{s_x} \right) \left(\frac{y_i - \bar{y}}{s_y} \right)$$

$$s_x = \sqrt{\frac{1}{n-1} \sum (x_i - \bar{x})^2} = \sqrt{\frac{\sum (x_i - \bar{x})^2}{n-1}}$$


$$\bar{y} = a + b\bar{x}$$

$$b = r \frac{s_y}{s_x}$$

**STATISTICS
SECTION II****Part A****Questions 1-5****Spend about 1 hour and 5 minutes on this part of the exam.****Percent of Section II score—75**

Directions: Show all your work. Indicate clearly the methods you use, because you will be scored on the correctness of your methods as well as on the accuracy and completeness of your results and explanations.


1. The sizes, in square feet, of the 20 rooms in a student residence hall at a certain university are summarized in the following histogram.



- (a) Based on the histogram, write a few sentences describing the distribution of room size in the residence hall.
- (b) Summary statistics for the sizes are given in the following table.

Mean	Standard Deviation	Min	Q1	Median	Q3	Max
231.4	68.12	134	174	253.5	292	315

Determine whether there are potential outliers in the data. Then use the following grid to sketch a boxplot of room size.




- (c) What characteristic of the shape of the distribution of room size is apparent from the histogram but not from the boxplot?

STATISTICS**SECTION II****Part A****Questions 1-5****Spend about 65 minutes on this part of the exam.****Percent of Section II score—75**

Directions: Show all your work. Indicate clearly the methods you use, because you will be scored on the correctness of your methods as well as on the accuracy and completeness of your results and explanations.

1. Robin works as a server in a small restaurant, where she can earn a tip (extra money) from each customer she serves. The histogram below shows the distribution of her 60 tip amounts for one day of work.



- (a) Write a few sentences to describe the distribution of tip amounts for the day shown.
- (b) One of the tip amounts was \$8. If the \$8 tip had been \$18, what effect would the increase have had on the following statistics? Justify your answers.

The mean:

The median: