


The following information applies to parts A and B.

6. To win a game, a baseball team needs to score runs. To score runs, players need to get on base. The primary way players get on base is by hitting the ball. Figure 1 shows the relationship between the number of hits and the number of runs for 30 randomly selected professional baseball teams.

Figure 1. Scatterplot of Number of Runs by Number of Hits



Part A

Consider the scatterplot in Figure 1.

- Describe the relationship between the number of hits and the number of runs for these baseball teams, in context.
- Based on the data from the sample of 30 teams, an equation of the least-squares regression line model for predicting the number of runs from the number of hits is as follows.


$$\text{Predicted number of runs} = -372.2 + 0.823(\text{number of hits})$$

Using the given regression equation, calculate the point estimate for the predicted number of runs a team would score if they were to achieve their goal of 1,250 hits. Show your work.

Part B

Each team has a total salary equal to the sum of the salaries of all players on the team. The median total salary for the sample of 30 teams is \$160 million. In Figure 2, points with dots represent teams with a total salary greater than the median, and points with squares represent teams with a total salary less than the median.

Figure 2. Scatterplot of Number of Runs by Number of Hits and Salary Classification



Consider the scatterplot in Figure 2.

- Compare the team represented by the point that is circled and labeled A with the other teams that have the same total salary classification.
- For each salary classification, consider the linear relationship between the number of hits and the number of runs. For teams with a total salary greater than the median, is the strength of the linear relationship stronger than, weaker than, or similar to the strength of the linear relationship for teams with a total salary less than the median? Explain.

Part C

Different types of intervals can be used when working with linear regression models. One type of interval is a confidence interval for the slope of the least-squares regression line. Two other types of intervals in the context of this problem are as follows:

- A confidence interval that estimates the mean number of runs for all teams with a specific number of hits.
- A prediction interval that predicts the number of runs for a single team with a specific number of hits.

All three intervals use the same basic formula:

$$\text{Point Estimate} \pm t^* (\text{standard error}),$$

where critical value t^* comes from a t -distribution that has $n - 2$ degrees of freedom. The same critical value is used when finding each of these intervals with the same level of confidence.

Consider the point estimate for the predicted number of runs found in part A (ii) for the team whose goal is to achieve 1,250 hits next year. Recall that this value was calculated using the linear regression model based on the data from the sample of 30 teams.

- What is the critical value for a 95% confidence level that would be used for the confidence interval for the mean number of runs as well as for the prediction interval for the number of runs? Indicate the answer to two decimal places.
- The standard error for the confidence interval for the mean number of runs is 17.48. Assuming the conditions for inference are met, calculate the 95% confidence interval for the mean number of runs for all teams with 1,250 hits. Show your work.
- The standard error for the prediction interval for the number of runs is 56.78. Assuming the conditions for inference are met, calculate the 95% prediction interval for the number of runs for a single team with 1,250 hits. Show your work.

Part D

- Would a distribution of sample means be expected to have more variability or less variability than a distribution of individual observations? Explain.
- The standard error used in the confidence interval that estimates the mean number of runs for all teams with x hits can be found using the following formula.

$$\sqrt{s^2 \left(\frac{1}{n} + \frac{(x - \bar{x})^2}{\sum (x_i - \bar{x})^2} \right)}$$


The standard error used in the prediction interval that predicts the number of runs for a single team with x hits can be found using the following formula.

$$\sqrt{s^2 + s^2 \left(\frac{1}{n} + \frac{(x - \bar{x})^2}{\sum (x_i - \bar{x})^2} \right)}$$

In both standard error formulas, s is the standard deviation of the residuals, n is the sample size, and $\bar{x}$ is the sample mean number of hits. Based on the answer from part D (i) and the standard error formulas for the confidence interval and the prediction interval, explain why the prediction interval calculated in part C (iii) is wider than the confidence interval calculated in part C (ii).

STOP
END OF EXAM

Wildlife biologists are interested in the health of tule elk, a species of deer found in California. An important measurement of tule elk health is their weight. The weight of a tule elk is difficult to measure in the wild. However, chest circumference, which is believed to be related to the weight of a tule elk, can easily be measured from a safe distance using a harmless laser. A study was done to investigate whether chest circumference, in centimeters (cm), could be used to accurately estimate the weight, in kilograms (kg), of male tule elk. For the study, wildlife biologists captured 30 male tule elk, measured their chest circumference and weight, and then released the elk. The data for the 30 male tule elk are shown in the scatterplot.



(a) Describe the relationship between chest circumference and weight of male tule elk in context.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 5** on this page.

Following is the equation of the least-squares regression line relating chest circumference and weight for male tule elk.

$$\text{predicted weight} = -350.3 + 3.7455(\text{chest circumference})$$

(b) The weight of one male tule elk with a chest circumference of 145.9 cm is 204.3 kg.

(i) Using the equation of the least-squares regression line, calculate the predicted weight for this male tule elk. Show your work.

(ii) Calculate the residual for this male tule elk. Show your work.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 4** on this page.

The equation of the least-squares regression line relating chest circumference and weight for male tule elk is repeated here.

$$\text{predicted weight} = -350.3 + 3.7455(\text{chest circumference})$$

(c) Interpret the slope of the least-squares regression line in context.

(d) The sambar, another species of deer, is similar in size to the tule elk. The slope of the population regression line relating chest circumference and weight for all male sambars is 4.5 kilograms per centimeter. A wildlife biologist wants to determine whether the slope of the population regression line for male tule elk is different than that for male sambars. Let β represent the slope of the population regression line for male tule elk. The wildlife biologist conducted a test of the following hypotheses using the sample of 30 tule elk.

$$H_0: \beta = 4.5$$

$$H_a: \beta \neq 4.5$$

The test statistic was calculated to be 3.408. Assume all conditions for inference were met.

(i) Determine the p -value of the test.

GO ON TO THE NEXT PAGE.Continue your response to **QUESTION 5** on this page.

(ii) At a significance level of $\alpha = 0.05$, what conclusion should the wildlife biologist make regarding the slope of the population regression line for male tule elk? Justify your response.

GO ON TO THE NEXT PAGE.


Part B

Question 6

Spend about 25 minutes on this part of the exam.

Directions: Show all your work. Indicate clearly the methods you use, because you will be scored on the correctness of your methods as well as on the accuracy and completeness of your results and explanations.

5. Attendance at games for a certain baseball team is being investigated by the team owner. The following boxplots summarize the attendance, measured as average number of attendees per game, for 47 years of the team's existence. The boxplots include the 30 years of games played in the old stadium and the 17 years played in the new stadium.




- (a) Compare the distributions of average attendance between the old and new stadiums.

GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

The following scatterplot shows average attendance versus year.




- (b) Compare the trends in average attendance over time between the old and new stadium.

GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

c) Consider the following scatterplots.



(i) Graph I shows the average attendance versus number of games won for each year. Describe the relationship between the variables.

(ii) Graph II shows the same information as Graph I, but also indicates the old and new stadiums. Does Graph II suggest that the rate at which attendance changes as number of games won increases is different in the new stadium compared to the old stadium? Explain your reasoning.

GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

(d) Consider the three variables: number of games won, year, and stadium. Based on the graphs, explain how one of those variables could be a confounding variable in the relationship between average attendance and the other variables.


GO ON TO THE NEXT PAGE.

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

STOP**END OF EXAM**

Use a pencil or pen with black or dark blue ink only. Do NOT write your name. Do NOT write outside the box.

Table entry for z is the probability lying below z .

**Table A** Standard normal probabilities

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
-3.4	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0002
-3.3	.0005	.0005	.0005	.0004	.0004	.0004	.0004	.0004	.0004	.0003
-3.2	.0007	.0007	.0006	.0006	.0006	.0006	.0006	.0005	.0005	.0005
-3.1	.0010	.0009	.0009	.0009	.0008	.0008	.0008	.0008	.0007	.0007
-3.0	.0013	.0013	.0013	.0012	.0012	.0011	.0011	.0011	.0010	.0010
-2.9	.0019	.0018	.0018	.0017	.0016	.0016	.0015	.0015	.0014	.0014
-2.8	.0026	.0025	.0024	.0023	.0023	.0022	.0021	.0021	.0020	.0019
-2.7	.0035	.0034	.0033	.0032	.0031	.0030	.0029	.0028	.0027	.0026
-2.6	.0047	.0045	.0044	.0043	.0041	.0040	.0039	.0038	.0037	.0036
-2.5	.0062	.0060	.0059	.0057	.0055	.0054	.0052	.0051	.0049	.0048
-2.4	.0082	.0080	.0078	.0075	.0073	.0071	.0069	.0068	.0066	.0064
-2.3	.0107	.0104	.0102	.0099	.0096	.0094	.0091	.0089	.0087	.0084
-2.2	.0139	.0136	.0132	.0129	.0125	.0122	.0119	.0116	.0113	.0110
-2.1	.0179	.0174	.0170	.0166	.0162	.0158	.0154	.0150	.0146	.0143
-2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183
-1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
-1.8	.0359	.0351	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
-1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
-1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
-1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
-1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0721	.0708	.0694	.0681
-1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
-1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
-1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
-1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
-0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
-0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
-0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
-0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
-0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
-0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
-0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
-0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
-0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
-0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641




Table entry for z is the probability lying below z .

Table A (Continued)

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
0.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
0.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
0.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
0.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
0.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
0.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
0.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
0.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
0.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.9997
3.4	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9998





Table entry for p and C is the point t^* with probability p lying above it and probability C lying between $-t^*$ and t^* .

Table B t distribution critical values

df	Tail probability p										
	.25	.20	.15	.10	.05	.025	.02	.01	.005	.0025	.001
1	1.000	1.376	1.963	3.078	6.314	12.71	15.89	31.82	63.66	127.3	318.3
2	.816	1.061	1.386	1.886	2.920	4.303	4.849	6.965	9.925	14.09	22.33
3	.765	.978	1.250	1.638	2.353	3.182	3.482	4.541	5.841	7.453	10.21
4	.741	.941	1.190	1.533	2.132	2.776	2.999	3.747	4.604	5.598	7.173
5	.727	.920	1.156	1.476	2.015	2.571	2.757	3.365	4.032	4.773	5.893
6	.718	.906	1.134	1.440	1.943	2.447	2.612	3.143	3.707	4.317	5.208
7	.711	.896	1.119	1.415	1.895	2.365	2.517	2.998	3.499	4.029	4.785
8	.706	.889	1.108	1.397	1.860	2.306	2.449	2.896	3.355	3.833	4.501
9	.703	.883	1.100	1.383	1.833	2.262	2.398	2.821	3.250	3.690	4.297
10	.700	.879	1.093	1.372	1.812	2.228	2.359	2.764	3.169	3.581	4.144
11	.697	.876	1.088	1.363	1.796	2.201	2.328	2.718	3.106	3.497	4.025
12	.695	.873	1.083	1.356	1.782	2.179	2.303	2.681	3.055	3.428	3.930
13	.694	.870	1.079	1.350	1.771	2.160	2.282	2.650	3.012	3.372	3.852
14	.692	.868	1.076	1.345	1.761	2.145	2.264	2.624	2.977	3.326	3.787
15	.691	.866	1.074	1.341	1.753	2.131	2.249	2.602	2.947	3.286	3.733
16	.690	.865	1.071	1.337	1.746	2.120	2.235	2.583	2.921	3.252	3.686
17	.689	.863	1.069	1.333	1.740	2.110	2.224	2.567	2.898	3.222	3.646
18	.688	.862	1.067	1.330	1.734	2.101	2.214	2.552	2.878	3.197	3.611
19	.688	.861	1.066	1.328	1.729	2.093	2.205	2.539	2.861	3.174	3.579
20	.687	.860	1.064	1.325	1.725	2.086	2.197	2.528	2.845	3.153	3.552
21	.686	.859	1.063	1.323	1.721	2.080	2.189	2.518	2.831	3.135	3.527
22	.686	.858	1.061	1.321	1.717	2.074	2.183	2.508	2.819	3.119	3.505
23	.685	.858	1.060	1.319	1.714	2.069	2.177	2.500	2.807	3.104	3.485
24	.685	.857	1.059	1.318	1.711	2.064	2.172	2.492	2.797	3.091	3.467
25	.684	.856	1.058	1.316	1.708	2.060	2.167	2.485	2.787	3.078	3.450
26	.684	.856	1.058	1.315	1.706	2.056	2.162	2.479	2.779	3.067	3.435
27	.684	.855	1.057	1.314	1.703	2.052	2.158	2.473	2.771	3.057	3.421
28	.683	.855	1.056	1.313	1.701	2.048	2.154	2.467	2.763	3.047	3.408
29	.683	.854	1.055	1.311	1.699	2.045	2.150	2.462	2.756	3.038	3.396
30	.683	.854	1.055	1.310	1.697	2.042	2.147	2.457	2.750	3.030	3.385
40	.681	.851	1.050	1.303	1.684	2.021	2.123	2.423	2.704	2.971	3.307
50	.679	.849	1.047	1.299	1.676	2.009	2.109	2.403	2.678	2.937	3.261
60	.679	.848	1.045	1.296	1.671	2.000	2.099	2.390	2.660	2.915	3.232
80	.678	.846	1.043	1.292	1.664	1.990	2.088	2.374	2.639	2.887	3.195
100	.677	.845	1.042	1.290	1.660	1.984	2.081	2.364	2.626	2.871	3.174
1000	.675	.842	1.037	1.282	1.646	1.962	2.056	2.330	2.581	2.813	3.098
∞	.674	.841	1.036	1.282	1.645	1.960	2.054	2.326	2.576	2.807	3.091
Confidence level C											
50% 60% 70% 80% 90% 95% 96% 98% 99% 99.5% 99.8% 99.9%											

Table entry for p is the point (χ^2) with probability p lying above it.

Table C χ^2 critical values


df	Tail probability p										
	.25	.20	.15	.10	.05	.025	.02	.01	.005	.0025	.001
1	1.32	1.64	2.07	2.71	3.84	5.02	5.41	6.63	7.88	9.14	10.83
2	2.77	3.22	3.79	4.61	5.99	7.38	7.82	9.21	10.60	11.98	13.82
3	4.11	4.64	5.32	6.25	7.81	9.35	9.84	11.34	12.84	14.32	16.27
4	5.39	5.99	6.74	7.78	9.49	11.14	11.67	13.28	14.86	16.42	18.47
5	6.63	7.29	8.12	9.24	11.07	12.83	13.39	15.09	16.75	18.39	20.51
6	7.84	8.56	9.45	10.64	12.59	14.45	15.03	16.81	18.55	20.25	22.46
7	9.04	9.80	10.75	12.02	14.07	16.01	16.62	18.48	20.28	22.04	24.32
8	10.22	11.03	12.03	13.36	15.51	17.53	18.17	20.09	21.95	23.77	26.12
9	11.39	12.24	13.29	14.68	16.92	19.02	19.68	21.67	23.59	25.46	27.88
10	12.55	13.44	14.53	15.99	18.31	20.48	21.16	23.21	25.19	27.11	29.59
11	13.70	14.63	15.77	17.28	19.68	21.92	22.62	24.72	26.76	28.73	31.26
12	14.85	15.81	16.99	18.55	21.03	23.34	24.05	26.22	28.30	30.32	32.91
13	15.98	16.98	18.20	19.81	22.36	24.74	25.47	27.69	29.82	31.88	34.53
14	17.12	18.15	19.41	21.06	23.68	26.12	26.87	29.14	31.32	33.43	36.12
15	18.25	19.31	20.60	22.31	25.00	27.49	28.26	30.58	32.80	34.95	37.70
16	19.37	20.47	21.79	23.54	26.30	28.85	29.63	32.00	34.27	36.46	39.25
17	20.49	21.61	22.98	24.77	27.59	30.19	31.00	33.41	35.72	37.95	40.79
18	21.60	22.76	24.16	25.99	28.87	31.53	32.35	34.81	37.16	39.42	42.31
19	22.72	23.90	25.33	27.20	30.14	32.85	33.69	36.19	38.58	40.88	43.82
20	23.83	25.04	26.50	28.41	31.41	34.17	35.02	37.57	40.00	42.34	45.31
21	24.93	26.17	27.66	29.62	32.67	35.48	36.34	38.93	41.40	43.78	46.80
22	26.04	27.30	28.82	30.81	33.92	36.78	37.66	40.29	42.80	45.20	48.27
23	27.14	28.43	29.98	32.01	35.17	38.08	38.97	41.64	44.18	46.62	49.73
24	28.24	29.55	31.13	33.20	36.42	39.36	40.27	42.98	45.56	48.03	51.18
25	29.34	30.68	32.28	34.38	37.65	40.65	41.57	44.31	46.93	49.44	52.62
26	30.43	31.79	33.43	35.56	38.89	41.92	42.86	45.64	48.29	50.83	54.05
27	31.53	32.91	34.57	36.74	40.11	43.19	44.14	46.96	49.64	52.22	55.48
28	32.62	34.03	35.71	37.92	41.34	44.46	45.42	48.28	50.99	53.59	56.89
29	33.71	35.14	36.85	39.09	42.56	45.72	46.69	49.59	52.34	54.97	58.30
30	34.80	36.25	37.99	40.26	43.77	46.98	47.96	50.89	53.67	56.33	59.70
40	45.62	47.27	49.24	51.81	55.76	59.34	60.44	63.69	66.77	69.70	73.40
50	56.33	58.16	60.35	63.17	67.50	71.42	72.61	76.15	79.49	82.66	86.66
60	66.98	68.97	71.34	74.40	79.08	83.30	84.58	88.38	91.95	95.34	99.61
80	88.13	90.41	93.11	96.58	101.9	106.6	108.1	112.3	116.3	120.1	124.8
100	109.1	111.7	114.7	118.5	124.3	129.6	131.1	135.8	140.2	144.3	149.4

STATISTICS
SECTION II
Part B
Question 6

Spend about 25 minutes on this part of the exam.
Percent of Section II score—25

Directions: Show all your work. Indicate clearly the methods you use, because you will be scored on the correctness of your methods as well as on the accuracy and completeness of your results and explanations.

6. A newspaper in Germany reported that the more semesters needed to complete an academic program at the university, the greater the starting salary in the first year of a job. The report was based on a study that used a random sample of 24 people who had recently completed an academic program. Information was collected on the number of semesters each person in the sample needed to complete the program and the starting salary, in thousands of euros, for the first year of a job. The data are shown in the scatterplot below.



- (a) Does the scatterplot support the newspaper report about number of semesters and starting salary? Justify your answer.


The table below shows computer output from a linear regression analysis on the data.

Predictor	Coef	SE Coef	T	P
Constant	34.018	4.455	7.64	0.000
Semesters	1.1594	0.3482	3.33	0.003

$S = 7.37702$ $R\text{-Sq} = 33.5\%$ $R\text{-Sq(adj)} = 30.5\%$

- (b) Identify the slope of the least-squares regression line, and interpret the slope in context.

An independent researcher received the data from the newspaper and conducted a new analysis by separating the data into three groups based on the major of each person. A revised scatterplot identifying the major of each person is shown below.



- (c) Based on the people in the sample, describe the association between starting salary and number of semesters for the business majors.
- (d) Based on the people in the sample, compare the median starting salaries for the three majors.
- (e) Based on the analysis conducted by the independent researcher, how could the newspaper report be modified to give a better description of the relationship between the number of semesters and the starting salary for the people in the sample?

STOP

END OF EXAM




Table entry for z is the probability lying below z .

Table A Standard normal probabilities

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
-3.4	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0002
-3.3	.0005	.0005	.0005	.0004	.0004	.0004	.0004	.0004	.0004	.0003
-3.2	.0007	.0007	.0006	.0006	.0006	.0006	.0006	.0005	.0005	.0005
-3.1	.0010	.0009	.0009	.0009	.0008	.0008	.0008	.0008	.0007	.0007
-3.0	.0013	.0013	.0013	.0012	.0012	.0011	.0011	.0011	.0010	.0010
-2.9	.0019	.0018	.0018	.0017	.0016	.0016	.0015	.0015	.0014	.0014
-2.8	.0026	.0025	.0024	.0023	.0023	.0022	.0021	.0021	.0020	.0019
-2.7	.0035	.0034	.0033	.0032	.0031	.0030	.0029	.0028	.0027	.0026
-2.6	.0047	.0045	.0044	.0043	.0041	.0040	.0039	.0038	.0037	.0036
-2.5	.0062	.0060	.0059	.0057	.0055	.0054	.0052	.0051	.0049	.0048
-2.4	.0082	.0080	.0078	.0075	.0073	.0071	.0069	.0068	.0066	.0064
-2.3	.0107	.0104	.0102	.0099	.0096	.0094	.0091	.0089	.0087	.0084
-2.2	.0139	.0136	.0132	.0129	.0125	.0122	.0119	.0116	.0113	.0110
-2.1	.0179	.0174	.0170	.0166	.0162	.0158	.0154	.0150	.0146	.0143
-2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183
-1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
-1.8	.0359	.0351	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
-1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
-1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
-1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
-1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0721	.0708	.0694	.0681
-1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
-1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
-1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
-1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
-0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
-0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
-0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
-0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
-0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
-0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
-0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
-0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
-0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
-0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641




Table entry for z is the probability lying below z .

Table A (Continued)

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
0.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
0.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
0.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
0.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
0.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
0.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
0.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
0.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
0.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9997	.9997
3.4	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9998




Table entry for p and C is the point t^* with probability p lying above it and probability C lying between $-t^*$ and t^* .

Table B t distribution critical values

df	Tail probability p									
	.25	.20	.15	.10	.05	.025	.02	.01	.005	.0025
1	1.000	1.376	1.963	3.078	6.314	12.71	15.89	31.82	63.66	127.3
2	.816	1.061	1.386	1.886	2.920	4.303	4.849	6.965	9.925	14.09
3	.765	.978	1.250	1.638	2.353	3.182	3.482	4.541	5.841	7.453
4	.741	.941	1.190	1.533	2.132	2.776	2.999	3.747	4.604	5.598
5	.727	.920	1.156	1.476	2.015	2.571	2.757	3.365	4.032	4.773
6	.718	.906	1.134	1.440	1.943	2.447	2.612	3.143	3.707	4.317
7	.711	.896	1.119	1.415	1.895	2.365	2.517	2.998	3.499	4.029
8	.706	.889	1.108	1.397	1.860	2.306	2.449	2.896	3.355	3.833
9	.703	.883	1.100	1.383	1.833	2.262	2.398	2.821	3.250	3.690
10	.700	.879	1.093	1.372	1.812	2.228	2.359	2.764	3.169	3.581
11	.697	.876	1.088	1.363	1.796	2.201	2.328	2.718	3.106	3.497
12	.695	.873	1.083	1.356	1.782	2.179	2.303	2.681	3.055	3.428
13	.694	.870	1.079	1.350	1.771	2.160	2.282	2.650	3.012	3.372
14	.692	.868	1.076	1.345	1.761	2.145	2.264	2.624	2.977	3.326
15	.691	.866	1.074	1.341	1.753	2.131	2.249	2.602	2.947	3.286
16	.690	.865	1.071	1.337	1.746	2.120	2.235	2.583	2.921	3.252
17	.689	.863	1.069	1.333	1.740	2.110	2.224	2.567	2.898	3.222
18	.688	.862	1.067	1.330	1.734	2.101	2.214	2.552	2.878	3.197
19	.688	.861	1.066	1.328	1.729	2.093	2.205	2.539	2.861	3.174
20	.687	.860	1.064	1.325	1.725	2.086	2.197	2.528	2.845	3.153
21	.686	.859	1.063	1.323	1.721	2.080	2.189	2.518	2.831	3.135
22	.686	.858	1.061	1.321	1.717	2.074	2.183	2.508	2.819	3.119
23	.685	.858	1.060	1.319	1.714	2.069	2.177	2.500	2.807	3.104
24	.685	.857	1.059	1.318	1.711	2.064	2.172	2.492	2.797	3.091
25	.684	.856	1.058	1.316	1.708	2.060	2.167	2.485	2.787	3.078
26	.684	.856	1.058	1.315	1.706	2.056	2.162	2.479	2.779	3.067
27	.684	.855	1.057	1.314	1.703	2.052	2.158	2.473	2.771	3.057
28	.683	.855	1.056	1.313	1.701	2.048	2.154	2.467	2.763	3.047
29	.683	.854	1.055	1.311	1.699	2.045	2.150	2.462	2.756	3.038
30	.683	.854	1.055	1.310	1.697	2.042	2.147	2.457	2.750	3.030
40	.681	.851	1.050	1.303	1.684	2.021	2.123	2.423	2.704	2.971
50	.679	.849	1.047	1.299	1.676	2.009	2.109	2.403	2.678	2.937
60	.679	.848	1.045	1.296	1.671	2.000	2.099	2.390	2.660	2.915
80	.678	.846	1.043	1.292	1.664	1.990	2.088	2.374	2.639	2.887
100	.677	.845	1.042	1.290	1.660	1.984	2.081	2.364	2.626	2.871
1000	.675	.842	1.037	1.282	1.646	1.962	2.056	2.330	2.581	2.813
∞	.674	.841	1.036	1.282	1.645	1.960	2.054	2.326	2.576	2.807
Confidence level C										
	50%	60%	70%	80%	90%	95%	96%	98%	99%	99.5%
										99.8%
										99.9%

Table entry for p is the point (χ^2) with probability p lying above it.




Table C χ^2 critical values

df	Tail probability p											
	.25	.20	.15	.10	.05	.025	.02	.01	.005	.0025	.001	.0005
1	1.32	1.64	2.07	2.71	3.84	5.02	5.41	6.63	7.88	9.14	10.83	12.12
2	2.77	3.22	3.79	4.61	5.99	7.38	7.82	9.21	10.60	11.98	13.82	15.20
3	4.11	4.64	5.32	6.25	7.81	9.35	9.84	11.34	12.84	14.32	16.27	17.73
4	5.39	5.99	6.74	7.78	9.49	11.14	11.67	13.28	14.86	16.42	18.47	20.00
5	6.63	7.29	8.12	9.24	11.07	12.83	13.39	15.09	16.75	18.39	20.51	22.11
6	7.84	8.56	9.45	10.64	12.59	14.45	15.03	16.81	18.55	20.25	22.46	24.10
7	9.04	9.80	10.75	12.02	14.07	16.01	16.62	18.48	20.28	22.04	24.32	26.02
8	10.22	11.03	12.03	13.36	15.51	17.53	18.17	20.09	21.95	23.77	26.12	27.87
9	11.39	12.24	13.29	14.68	16.92	19.02	19.68	21.67	23.59	25.46	27.88	29.67
10	12.55	13.44	14.53	15.99	18.31	20.48	21.16	23.21	25.19	27.11	29.59	31.42
11	13.70	14.63	15.77	17.28	19.68	21.92	22.62	24.72	26.76	28.73	31.26	33.14
12	14.85	15.81	16.99	18.55	21.03	23.34	24.05	26.22	28.30	30.32	32.91	34.82
13	15.98	16.98	18.20	19.81	22.36	24.74	25.47	27.69	29.82	31.88	34.53	36.48
14	17.12	18.15	19.41	21.06	23.68	26.12	26.87	29.14	31.32	33.43	36.12	38.11
15	18.25	19.31	20.60	22.31	25.00	27.49	28.26	30.58	32.80	34.95	37.70	39.72
16	19.37	20.47	21.79	23.54	26.30	28.85	29.63	32.00	34.27	36.46	39.25	41.31
17	20.49	21.61	22.98	24.77	27.59	30.19	31.00	33.41	35.72	37.95	40.79	42.88
18	21.60	22.76	24.16	25.99	28.87	31.53	32.35	34.81	37.16	39.42	42.31	44.43
19	22.72	23.90	25.33	27.20	30.14	32.85	33.69	36.19	38.58	40.88	43.82	45.97
20	23.83	25.04	26.50	28.41	31.41	34.17	35.02	37.57	40.00	42.34	45.31	47.50
21	24.93	26.17	27.66	29.62	32.67	35.48	36.34	38.93	41.40	43.78	46.80	49.01
22	26.04	27.30	28.82	30.81	33.92	36.78	37.66	40.29	42.80	45.20	48.27	50.51
23	27.14	28.43	29.98	32.01	35.17	38.08	38.97	41.64	44.18	46.62	49.73	52.00
24	28.24	29.55	31.13	33.20	36.42	39.36	40.27	42.98	45.56	48.03	51.18	53.48
25	29.34	30.68	32.28	34.38	37.65	40.65	41.57	44.31	46.93	49.44	52.62	54.95
26	30.43	31.79	33.43	35.56	38.89	41.92	42.86	45.64	48.29	50.83	54.05	56.41
27	31.53	32.91	34.57	36.74	40.11	43.19	44.14	46.96	49.64	52.22	55.48	57.86
28	32.62	34.03	35.71	37.92	41.34	44.46	45.42	48.28	50.99	53.59	56.89	59.30
29	33.71	35.14	36.85	39.09	42.56	45.72	46.69	49.59	52.34	54.97	58.30	60.73
30	34.80	36.25	37.99	40.26	43.77	46.98	47.96	50.89	53.67	56.33	59.70	62.16
40	45.62	47.27	49.24	51.81	55.76	59.34	60.44	63.69	66.77	69.70	73.40	76.09
50	56.33	58.16	60.35	63.17	67.50	71.42	72.61	76.15	79.49	82.66	86.66	89.56
60	66.98	68.97	71.34	74.40	79.08	83.30	84.58	88.38	91.95	95.34	99.61	102.7
80	88.13	90.41	93.11	96.58	101.9	106.6	108.1	112.3	116.3	120.1	124.8	128.3
100	109.1	111.7	114.7	118.5	124.3	129.6	131.1	135.8	140.2	144.3	149.4	153.2