AP Statistics

Revision Sheet · 考前速记

Learn these by heart before the exam. 考试前把这些内容背熟。

Exploring One-Variable Data

$\bar{x} = \dfrac{\sum x_i}{n}, \quad s_x = \sqrt{\dfrac{\sum (x_i - \bar{x})^2}{n-1}}$ 样本均值与标准差

Describe a distribution with SOCS: Shape, Outliers, Centre, Spread. In context, always.

Skewed right pulls the MEAN above the median. Use median and IQR for skewed data, mean and s for roughly symmetric.

outlier $\iff x < Q_1 - 1.5\,\text{IQR}$ or $x > Q_3 + 1.5\,\text{IQR}$ 离群值判定

$z = \dfrac{x - \mu}{\sigma}$ how many standard deviations from the mean; unitless 标准分数

Linear transform $y = a + bx$: centre scales by b and shifts by a; spread scales by $|b|$ only.

Exploring Two-Variable Data

$\hat{y} = a + bx, \quad b = r\dfrac{s_y}{s_x}, \quad a = \bar{y} - b\bar{x}$ the line always passes through $(\bar{x}, \bar{y})$ 最小二乘回归线

$r = \dfrac{1}{n-1}\sum\left(\dfrac{x_i - \bar{x}}{s_x}\right)\left(\dfrac{y_i - \bar{y}}{s_y}\right)$ $-1 \le r \le 1$, unitless, and NOT resistant to outliers 相关系数

r^2 is the proportion of variation in y explained by the linear model with x. Say it that way.

residual $= y - \hat{y}$ a curved residual plot means a linear model is wrong 残差

Correlation is not causation. Only a randomised EXPERIMENT supports a causal claim.

Influential point: far in the x direction, moves the line a lot. An outlier in y mainly inflates residuals.

Collecting Data

SRS, stratified (homogeneous groups, sample within each), cluster (sample whole groups), systematic. Say which and why.

Bias sources: voluntary response, undercoverage, nonresponse, question wording. Name the specific one.

Experiment needs: comparison, random assignment, replication, control. Random ASSIGNMENT gives causation; random SELECTION gives generalisability.

Blocking is to experiments what stratifying is to samples: group similar units, then randomise WITHIN each block.

Blind means subjects do not know the treatment. Double-blind means the assessors do not either.

Probability and Random Variables

$P(A \cup B) = P(A) + P(B) - P(A \cap B)$ 加法法则

$P(A \mid B) = \dfrac{P(A \cap B)}{P(B)}$ 条件概率

Independent $\iff P(A \mid B) = P(A) \iff P(A \cap B) = P(A)P(B)$. *Mutually exclusive is NOT the same as independent.*

$\mu_X = \sum x_i p_i, \quad \sigma_X^2 = \sum (x_i - \mu_X)^2 p_i$ 离散型随机变量的期望与方差

$\mu_{X \pm Y} = \mu_X \pm \mu_Y, \quad \sigma_{X \pm Y}^2 = \sigma_X^2 + \sigma_Y^2$ variances ADD even when subtracting, and only if X and Y are independent 随机变量的和与差

$P(X = k) = \binom{n}{k} p^k (1-p)^{n-k}, \quad \mu = np, \quad \sigma = \sqrt{np(1-p)}$ BINS: Binary, Independent, fixed Number, Same p 二项分布

$P(X = k) = (1-p)^{k-1} p, \quad \mu = \dfrac{1}{p}$ 几何分布

Sampling Distributions

$\mu_{\bar{x}} = \mu, \quad \sigma_{\bar{x}} = \dfrac{\sigma}{\sqrt{n}}$ 样本均值的抽样分布

$\mu_{\hat{p}} = p, \quad \sigma_{\hat{p}} = \sqrt{\dfrac{p(1-p)}{n}}$ 样本比例的抽样分布

CLT: for $n \ge 30$ the sampling distribution of $\bar{x}$ is approximately normal WHATEVER the population shape.

10% condition: sampling without replacement needs $n \le 0.10N$ for independence.

Large counts for proportions: $np \ge 10$ AND $n(1-p) \ge 10$.

Inference - the common structure

estimate $\pm$ (critical value) $\times$ (standard error) 置信区间的通用形式

$$\text{test statistic} = \frac{\text{statistic} - \text{parameter}}{\text{standard error}}$$ 检验统计量的通用形式

Four steps every time: State (hypotheses + α), Plan (name the test + check conditions), Do (statistic + p-value), Conclude (in context).

Interpret a CI: 'We are 95% confident the true [parameter in context] is between A and B.' Never 'the probability is 95%'.

Interpret a p-value: 'the probability of a result at least this extreme IF H_0 were true'.

$p < \alpha$: reject H_0, evidence for H_a. $p \ge \alpha$: fail to reject. NEVER 'accept H_0'.

Type I: reject a true H_0 (probability α). Type II: fail to reject a false H_0. Power $= 1 - P(\text{Type II})$; it rises with larger n, larger α, or a bigger true effect.

Which test to use

$\hat{p} \pm z^* \sqrt{\dfrac{\hat{p}(1-\hat{p})}{n}}, \quad z = \dfrac{\hat{p} - p_0}{\sqrt{p_0(1-p_0)/n}}$ the TEST uses p_0 in the standard error; the INTERVAL uses $\hat{p}$ 单比例

$\bar{x} \pm t^* \dfrac{s_x}{\sqrt{n}}, \quad t = \dfrac{\bar{x} - \mu_0}{s_x/\sqrt{n}}, \quad df = n - 1$ 单均值

$(\hat{p}_1 - \hat{p}_2) \pm z^* \sqrt{\dfrac{\hat{p}_1(1-\hat{p}_1)}{n_1} + \dfrac{\hat{p}_2(1-\hat{p}_2)}{n_2}}$ the two-proportion TEST pools: $\hat{p}_c = \frac{X_1 + X_2}{n_1 + n_2}$ 两比例之差

$\chi^2 = \sum \dfrac{(O-E)^2}{E}$ goodness-of-fit $df = k - 1$; independence/homogeneity $df = (r-1)(c-1)$; need every $E \ge 5$ 卡方统计量

$b \pm t^* SE_b, \quad t = \dfrac{b}{SE_b}, \quad df = n - 2$ 回归斜率

Paired data (before/after on the SAME unit) is a ONE-sample t on the differences, not a two-sample test.

Exam technique

Always in context. Name the variable and the population in every conclusion; a generic sentence scores E for essentially correct only if contextual.

Check and STATE conditions: random, 10%, and normal/large counts. Listing them without checking earns nothing.

Show the setup: write the formula with numbers substituted, then the answer. Calculator output alone is not enough.