

Probability and Random Variables

AP Statistics • Unit 4


AP Statistics



Probability and Random Variables

What we will cover

- 1 Randomness and probability
- 2 Probability rules
- 3 Conditional and independence
- 4 Random variables
- 5 Combining variables
- 6 Binomial and geometric



1 Randomness

Probability and Random Variables

Randomness

Randomness and probability

Probability 概率 is the long-run **relative frequency** 相对频率 of an outcome.

- short runs vary by **chance** 偶然
- **simulation** 模拟 estimates a hard probability

Law of large numbers 大数定律: more trials $\Rightarrow \hat{p}$ settles toward the true p .



Playing cards

2 Rules

Probability and Random Variables

Rules


Probability rules

From the **sample space** 样本空间, every **event** 事件 obeys $0 \leq P \leq 1$, summing to 1.

$$P(A^c) = 1 - P(A)$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

Mutually exclusive 互斥 events share nothing, so $P(A \cap B) = 0$ and the last term drops.



3 Conditional

Probability and Random Variables
└ Conditional

Conditional probability and independence

A **conditional probability** 条件概率 restricts to the times B happens:

$$P(A | B) = \frac{P(A \cap B)}{P(B)}$$

The **multiplication rule** 乘法法则:
 $P(A \cap B) = P(A) P(B | A)$.

Independent 独立: $P(A | B) = P(A)$, so $P(A \cap B) = P(A) P(B)$. Not the same as mutually exclusive!

	girl	boy
sport	24	36
none	16	24

$P(\text{girl} | \text{sport}) = \frac{24}{60}$

4 Random variables

Probability and Random Variables
└ Random variables


Random variables

A **random variable** 随机变量 X takes a number from a chance process
(**discrete** 离散 or **continuous** 连续).

- its **distribution** 概率分布 lists each x with $P(x)$, summing to 1

$$\mu_X = \sum x_i P(x_i) \quad \sigma_X^2 = \sum (x_i - \mu_X)^2 P(x_i)$$

μ_X is the **expected value** 期望值 – the long-run average outcome.



5 Combining

Probability and Random Variables
└ Combining

Combining random variables

A **linear transformation** 线性变换:

$$\mu_{aX+b} = a\mu_X + b \quad \sigma_{aX+b} = |a| \sigma_X$$

For a **sum or difference** 和或差 of **independent** 独立 variables, means combine and **variances add** – for both:

$$\mu_{X \pm Y} = \mu_X \pm \mu_Y \quad \sigma_{X \pm Y}^2 = \sigma_X^2 + \sigma_Y^2$$

Never subtract variances for a difference – variances always **add**.

Two packages

$\sigma_X = 0.3$, $\sigma_Y = 0.4$, independent.

$$\sigma_{X+Y} = \sqrt{0.3^2 + 0.4^2} = 0.5$$

The difference $X - Y$ has the same $\sigma = 0.5$.

6 Binomial


The binomial distribution

The **binomial setting** 二项分布情境 counts successes – remember **BINS**: **B**inary, **I**ndependent, fixed **N**umber, **S**ame p .

$$P(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}$$

$$\mu_X = np \quad \sigma_X = \sqrt{np(1-p)}$$

Shape: symmetric at $p = 0.5$, skewed otherwise; more bell-shaped as n grows.



Worked example – binomial by hand

A free-throw shooter makes 80%. In $n = 10$ attempts, find $P(X = 8)$ and describe the distribution's centre and spread.

$$\begin{aligned} P(X = 8) &= \binom{10}{8} (0.8)^8 (0.2)^2 \\ &= 45(0.1678)(0.04) \approx 0.302 \\ \mu &= np = 10(0.8) = 8 \text{ makes} \\ \sigma &= \sqrt{10(0.8)(0.2)} \approx 1.26 \end{aligned}$$

$\binom{10}{8} = 45$ counts which 8 shots dropped – forget it and you find one single ordering (0.0067), not the event. Check **BINS** first: without independence it is not binomial at all.


7 Geometric

The geometric distribution

The **geometric setting** 几何分布情境 keeps going *until* the first success (no fixed n).

$$P(X = k) = (1 - p)^{k-1} p \quad \mu_X = \frac{1}{p}$$

X counts *trials until* the first success; a binomial counts successes in a *fixed* number of trials.



8 Recap

Unit 4 – key takeaways

1 Probability

long-run relative frequency; law of large numbers

3 Random vars

$\mu = \sum xP(x)$; variances add for independent

2 Rules

addition for or; multiplication for and; conditional

4 Distributions

binomial $\mu = np$; geometric $\mu = 1/p$

Check yourself

- 1 Write the complement rule for $P(A)$.
- 2 For independent X, Y , how do you combine their variances?
- 3 What does BINS stand for?
- 4 A binomial has $n = 20$, $p = 0.5$: what is μ_X ?



Answers 1. $P(A^c) = 1 - P(A)$ 2. add them: $\sigma_X^2 + \sigma_Y^2$ 3. Binary, Independent, Number, Same p 4. $np = 10$