

1. Find the hypotenuse when the legs are 3 and 4.
- _____
2. For $r=4$ and $\theta=0.5$ radians, find arc length.
- _____
3. Using the small-angle formula, approximate $\sin(0.2 \text{ radians})$.
- _____
4. Find $\sec(2\pi/3)$.
- _____
5. If $\sec\theta=2$ and θ is acute, find $\tan^2\theta$.
- _____
6. Cosine of a sum equals the sum of the two cosine values.
- ☐ True

☐ False
7. The amplitude of a $\cos\theta+b \sin\theta$ is $|a|+|b|$.
- ☐ True

☐ False
8. Solving $\sin 2x$ on one full turn of $2x$ finds every x solution on $[0,2\pi)$.
- ☐ True

☐ False
9. The sine rule pairs each side with any angle in the triangle.
- ☐ True

☐ False
10. A transformed tangent graph has a finite amplitude and can be joined across its vertical asymptotes.
- ☐ True

☐ False

1. **5.0**
- Pythagoras gives $\sqrt{(3^2+4^2)}=5$.
2. **2.0**
- Arc length= $r\theta=4\times 0.5=2$.
3. **0.2**
- $\sin\theta\approx\theta$, with $\theta=0.2$ radians.
4. **-2.0**
- $\cos(2\pi/3)=-1/2$, so its reciprocal is -2 .
5. **3.0**
- $\tan^2\theta=\sec^2\theta-1=4-1=3$.
6. **False**
- $\cos(A+B)=\cos A \cos B-\sin A \sin B$, which is not $\cos A+\cos B$.
7. **False**
- The amplitude is $R=\sqrt{(a^2+b^2)}$: for $3\cos\theta+4\sin\theta$ it is 5, not 7.
8. **False**
- If x runs over $[0,2\pi)$, then $2x$ runs over $[0,4\pi)$, so solve for $2x$ over two full turns.
9. **False**
- Each side is paired with the angle opposite it: $a/\sin A=b/\sin B$.
10. **False**
- A tangent graph has no amplitude, and its branches must not be joined across asymptotes.