

P · Quantities and units in mechanics · supported focus

AQA · A-Level · Mathematics

Scope and prerequisites

AQA A-level Mathematics 7357, section P (Quantities and units in mechanics), version 1.3 (first assessment 2018). Original teaching notes; objective-level mapping lives in content/teaching/aqa_a_level_mathematics/objective-review.yaml.

Prerequisites: scientific notation and unit conversions from GCSE. Each lesson develops its own method before independent transfer.

These are original teaching examples, not official AQA material.

Quantities and units in mechanics

■ What would explain this observation?

- A report says a cyclist did “300 N of work pedalling 50 metres.” A marker crosses it out without checking any numbers. What did the marker see?
- Start with a prediction. State what is wrong with that sentence dimensionally, and what units work, force and power each need.

■ Build the model

- Every mechanics quantity is built from the base units kilogram, metre and second: velocity is m/s, acceleration is m/s², force is the newton (kg m/s²), work is the joule (N m) and power is the watt (J/s).
- **Work** is energy transferred by a force through a displacement: $W = Fs$ in the direction of the force. **Power** is the rate of doing work: $P = W/t$. Units are checked on every line – a correct equation gives the same unit on both sides, so a dimensional mismatch proves a slip before any number is trusted.
- work: Energy transferred by a force through a displacement, $W = Fs$, in joules; power: The rate of energy transfer, $P = W/t$, in watts.

Every mechanics quantity is built from kg, m and s



Check units on every line: $W = Fs$ gives $N \cdot m = J$,
 $F = ma$ gives $kg \times m/s^2 = N$

■ Choose evidence that can test it

- Convert before combining: speeds in km/h must become m/s (divide by 3.6) before meeting forces or energies. A value’s unit is part of its meaning; a bare number is not an answer in mechanics.
- Check a formula dimensionally on the DATA line, not just the result: if $F = ma$ is to give newtons, m must be in kilograms and a in m/s².

■ Work from known quantities

- State the known values and their units. Choose the relation because its assumptions fit this case, then rearrange before substitution.
- Known: a cyclist applies 80 N over 50 m in 20 s. Work = $Fs = 80 \times 50 = 4,000$ J. Power = $W/t = 4,000/20 = 200$ W. The “300 N of work” sentence fails dimensionally: newtons are force, and work is measured in joules.

Example:

A 60 kg student climbs 4 m of stairs in 10 s. Using $g = 9.8$ m/s², find the work against gravity and the average power, using the same sequence: known $\rightarrow W = mgh \rightarrow P = W/t$. (Answer: $W = 60 \times 9.8 \times 4 = 2,352$ J; $P = 2,352/10 \approx 235$ W.)

■ Check the conclusion and its limits

- A dimensional check catches wrong formulas but cannot catch a wrong NUMBER inside a right-shaped formula; units pass, then arithmetic must still be verified. “Powerful” is not a technical term – compare watts, not adjectives.
- Return to the original observation. Explain what the result supports, which conditions it assumes, and one way to test a competing explanation.

Warn:

Work, force and power can be quoted in the same units because they all involve force. This claim is false: force is in newtons, work in joules (N · m) and power in watts (J/s)

– different dimensions, and confusing them produces nonsense like “300 N of work.”

Key:

Quantities and units in mechanics: everything builds from kg, m, s; work is $W = Fs$ in joules, power is W/t in watts, and a dimensional check on every line catches formula slips before the arithmetic is trusted.