

Exercise walk-through

2.1.3

Manipulation

AQA · GCSE · Mathematics

141. Graphical gradients and area estimates · Higher

Graphical gradients and area estimates · Higher

For a distance curve $d=t^2$, the chord from (1,1) to (3,9) has gradient $(9-1)/(3-1)=4$. A tangent at (2,4) passes through (1,0) and (3,8), giving gradient 4 without calculus. A velocity-time trapezium with endpoint velocities 2 and 6 m/s over 3 s has area $(2+6) \times 3/2 = 12$ m. Smaller strips can improve a curved-area estimate; curvature affects over/underestimation. A cost-versus-quantity tangent through (2,12) and (6,28) gives a local rate $(28-12)/(6-2)=4$ yuan per extra item, linking graph slope to a financial interpretation.

183. Reasoning and check

Reasoning and check

For $F(x) = (x^2 - 9)/(x - 3)$, factor $x^2 - 9 = (x - 3)(x + 3)$. Thus $F(x) = x + 3$ for $x \neq 3$; $F(5) = 8$, but $F(3)$ is undefined, not 6. For addition, $1/(x - 1) + 2/(x + 1) = [(x + 1) + 2(x - 1)]/[(x - 1)(x + 1)] = (3x - 1)/(x^2 - 1)$, with $x \neq \pm 1$. At $x = 3$, both forms give 1. For subtraction, the numerator is $(x + 1) - 2(x - 1) = 3 - x$, so the minus sign acts on both terms. Multiplication $[(x - 1)/(x + 2)] \times [(x + 2)/(x + 1)]$ gives $(x - 1)/(x + 1)$, but the original excludes $x = -2$ and $x = -1$. Division $[(x^2 - 1)/(x^2 + 3x + 2)] \div [(x - 1)/(x + 2)]$ gives 1, yet $x = -2, -1, 1$ are excluded: $x = 1$ makes the divisor zero. Finally $F(x) = 6$ reduces to $x + 3 = 6$.

186. Reasoning and check

Reasoning and check

AQA A4 Higher includes algebraic fractions and valid cancellation. The equation example checks whether a candidate is inside the original domain. Explain each factor cancellation, retain original restrictions and check every equation candidate. These operations extend the existing manipulation example into a complete worked arithmetic sequence.

Exercises 2.1·2.3

2.1 (Foundation) Find the coefficient of x in $3x+4x$.

2.2 (Foundation) Find the constant term after expanding $2(x+3)$.

2.3 (Foundation) Find the coefficient of x in $(x+2)(x+3)$.

Exercises 2.1·2.3

2.1 (Foundation) Find the coefficient of x in $3x+4x$.

2.1 Add coefficients $3+4$.

2.2 (Foundation) Find the constant term after expanding $2(x+3)$.

2.2 Distribute 2 to both terms; $2 \times 3 = 6$.

2.3 (Foundation) Find the coefficient of x in $(x+2)(x+3)$.

2.3 The cross terms are $3x+2x=5x$.

Exercise 3.1

3.1 Decide whether this claim is true. Justify with a concrete calculation or counterexample: The terms $3x$ and $4y$ can be collected to give $7xy$.

Exercise 3.1

3.1 Decide whether this claim is true. Justify with a concrete calculation or counterexample: The terms $3x$ and $4y$ can be collected to give $7xy$.

3.1 False. At $x=1, y=2$, $3x+4y=11$, while $7xy=14$. The terms have different variable parts and cannot be collected as like terms.

Local review guidance: show the chosen method; give the checked result; explain the condition or failed shortcut. This is authored classroom guidance.