

1.1.3 Counting

Name: _____ Class: _____ Date: _____

KEY WORDS systematic list

<!-- Session 2 native-skill companion -->

Objective

Practise counting within the labelled native lessons. Public original tasks; locally authored answer guidance.

1 Key knowledge and worked examples

■ Systematic lists and possibility grids · Foundation

Fix one first choice and list every allowed second choice before moving to the next first choice. Use a table to check that no outcome is missing or duplicated. If combinations are forbidden, remove those entries from the list.

$$\{TA, TB, TM, JA, JB, JM\}$$

List tea-apple, tea-banana, tea-melon, juice-apple, juice-banana, juice-melon: six choices. Removing tea-melon leaves five. The codes using 1,2,3 once each are 123,132,213,231,312,321: six. These answers follow from complete lists.

State whether order matters and whether repetition is allowed. Choosing A then B can be different from B then A for a code, but not for an unordered two-person team. A restriction can make a simple product invalid.

Foundation should justify answers with a complete ordered list or grid. Higher may compress a verified structure using the product rule, but must still check restrictions.

■ Systematic lists and the product rule · Higher

Fix one first choice and list every permitted second choice before moving to the next first choice. A table gives the same structure. When every one of m first choices allows n second choices, the Higher product rule gives mn outcomes. If restrictions change the options, count the allowed rows separately.

$$\{TA, TB, TM, JA, JB, JM\}$$

The list is tea-apple, tea-banana, tea-melon, juice-apple, juice-banana, juice-melon: six outcomes. If tea-melon is unavailable, five remain. For a three-digit code with digits 1,2,3

State whether order matters and whether repetition is allowed. Choosing A then B can be different from B then A for a code, but not for an unordered two-person team. A restriction can make a simple product invalid.

Foundation should justify answers with a complete ordered list or grid. Higher may compress a verified structure using the product rule, but must still check restrictions.

A complete grid lists six choices once each

Juice + Apple	Juice + Banana	Juice + Melon
Tea + Apple	Tea + Banana	Tea + Melon

Original native-lesson illustration; labels belong to its worked example.

2 Practice

2.1 (Foundation) How many cafe choices are in the complete list?

2.2 (Foundation) How many remain when tea-melon is removed?

3 Exam-style questions

3.1 Decide whether this claim is true. Justify with a concrete calculation or counterexample: Three available first choices and four second choices always give twelve outcomes even if some combinations are forbidden.
