

2.1.99 Independent transfer: Foundation then Higher

Name: _____ Class: _____ Date: _____

KEY WORDS identity • root

<!-- Session 2 independent transfer -->

Objective


Apply this topic to unfamiliar Foundation and Higher problems; keep the two scopes distinct. Original questions and local marking guidance.

1 Key knowledge and worked examples


Translate fixed and variable costs into an equation with defined units. Elimination combines whole equations. In a rational equation, exclude zero denominators before cancelling factors. For a quadratic inequality, locate roots and test the intervals between them; completing the square locates its vertex.

$20+3x=44+x$ gives $2x=24$ and $x=12$. Both plans then cost 56. In $A=\pi r^2$, divide by π and take the positive square root to obtain $r=\sqrt{A/\pi}$, because r is a length.

$x^2-10x+21=(x-3)(x-7)$. Hence the equation $x^2-10x+21=0$ has roots 3 and 7. Check each root by substitution and mark both intercepts on the graph.



Prerequisite example; use the transfer task's own data.



Prerequisite example; use the transfer task's own data.

2 Practice

2.1 Foundation: A taxi charges a fixed 8 CNY plus 3 CNY per kilometre. Form a cost equation. A trip costs 35 CNY; find its distance and check. Then solve the simultaneous equations $2x + y = 11$, $x - y = 1$.

3 Exam-style questions

3.1 Higher: Solve $(x^2 - 9)/(x - 3) = x^2 - 5$ on its original real domain. Then solve $x^2 - 4x - 5 \leq 0$ completely. State the turning point and roots of $y = x^2 - 4x - 5$.
