

Exercise walk-through

3.1

Planning investigations (Unit 3 spec, planning section)

Pearson Edexcel · International A-Level · Biology

1

The method

Method — Name the variable, then give its reason

The **independent variable** is what you change; the **dependent variable** is what you measure; every other quantity that could affect the dependent variable is a **control variable** and must be held constant *with a reason*: temperature is controlled because it changes membrane fluidity and enzyme action; solution volume because it dilutes the measured pigment differently. A mark needs both halves – the variable and the effect it would have.

Method — A control treatment is not a control variable

A **control treatment** leaves the independent variable out (solvent without trifluralin; water instead of methanol) so any measured difference can be attributed to the independent variable alone. It allows a valid comparison; it is not “the control variable”.

Exercises 2.1·2.3

2.1 In a seedling investigation, state the independent variable when some seedlings sit in solvent containing trifluralin and others sit in solvent alone. [1]

2.2 Explain why the solvent-only seedlings are needed, in terms of what can be concluded. [2]

2.3 A student measures pigment loss from beetroot at 20 °C and at 60 °C. Give two control variables with the reason each must be fixed. [4]

Exercises 2.1·2.3

2.1 In a seedling investigation, state the independent variable when some seedlings sit in solvent containing trifluralin and others sit in solvent alone. [1]

2.1 The presence (or concentration) of trifluralin.

2.2 Explain why the solvent-only seedlings are needed, in terms of what can be concluded. [2]

2.2 It is a control treatment, so any difference in results is due to trifluralin and not the solvent; it allows a valid comparison between like and like.

2.3 A student measures pigment loss from beetroot at 20 °C and at 60 °C. Give two control variables with the reason each must be fixed. [4]

2.3 Any two, each with reason: beetroot piece size/surface area – larger pieces leak more pigment; volume of liquid – dilutes the pigment differently; time of immersion – leakage continues with time; same beetroot/age – pigment content varies; light – could affect membrane state.

Exercises 2.4·3.1

2.4 Explain why the seedlings were germinated on moist paper before treatment began, rather than placed onto treated discs as dry grains. [2]

3.1 Brine shrimp hatch rates are measured at 15 °C, 20 °C, 25 °C and 30 °C. Plan this investigation: identify the variables, describe how the dependent variable is measured, state two controls with reasons, and give one ethical consideration in the specification's terms. [6]

Exercises 2.4·3.1

2.4 Explain why the seedlings were germinated on moist paper before treatment began, rather than placed onto treated discs as dry grains. [2]

2.4 To ensure all grains used were viable and had begun growth, so every treatment starts from seedlings at the same stage of development.

3.1 Brine shrimp hatch rates are measured at 15 °C, 20 °C, 25 °C and 30 °C. Plan this investigation: identify the variables, describe how the dependent variable is measured, state two controls with reasons, and give one ethical consideration in the specification's terms. [6]

3.1 Independent: temperature. Dependent: hatching rate – number hatched per number of eggs per time interval, counted under a hand lens. Two controls with reasons, e.g. salinity of water – affects survival; oxygen/aeration – needed for respiration; number of eggs per dish; volume of water. Measurement detail: replicate dishes per temperature, water baths to hold temperature. Ethical: the specification requires the ethical use of organisms – minimal numbers, no deliberate harm beyond the treatment, appropriate disposal.

Exercise 3.2

3.2 Yeast is grown in glucose solution and its respiration rate followed with a redox indicator. Write a plan that states: the independent and dependent variables; two control variables with reasons; how the data will be presented and used; and one safety point with its control.

[6]

Exercise 3.2

3.2 Yeast is grown in glucose solution and its respiration rate followed with a redox indicator. Write a plan that states: the independent and dependent variables; two control variables with reasons; how the data will be presented and used; and one safety point with its control. [6]

3.2 Independent: substrate (glucose) concentration or temperature as planned. Dependent: rate of colour change of the redox indicator, timed to a fixed endpoint, $1/\text{time}$ as rate proxy. Controls with reasons: temperature – affects enzyme-controlled respiration rate; pH – affects the indicator colour; volume of yeast suspension – more cells respire faster; indicator volume. Data: table of concentration against mean time with units; line graph of rate against the independent variable; compare the initial gradient or rate proxy. Safety: none of the standard human pathogens cultured; wash hands, disinfect benches; keep the water bath electrical parts dry.

Exercise 3.3

3.3 An investigation germinates 40 wheat grains on filter paper soaked in 7 mg cm^{-3} trifluralin solution, incubated at 25°C , with root tips sampled after 12 hours. Justify the 25°C incubation temperature, and explain what the data would show if trifluralin arrested mitosis at metaphase. [5]

Exercise 3.3

3.3 An investigation germinates 40 wheat grains on filter paper soaked in 7 mg cm^{-3} trifluralin solution, incubated at 25°C , with root tips sampled after 12 hours. Justify the 25°C incubation temperature, and explain what the data would show if trifluralin arrested mitosis at metaphase. [5]

3.3 25°C is near the optimum for the enzymes of germination/mitosis, so rates are measurable and enzymes do not denature or slow excessively. If mitosis arrested at metaphase, cells in the treated roots would accumulate in metaphase – a raised percentage of cells at that stage, with prophase still seen but anaphase and telophase absent, because the spindle does not form or cannot separate the chromatids.