

1.4 Numerical solution of equations

Name: _____ Class: _____ Date: _____ **Total: 24 marks**

KEY WORDS interval bisection • linear interpolation
• Newton–Raphson method • continuity

Numerical solution of equations

Objective

Locate roots by interval bisection, linear interpolation and the Newton–Raphson process, showing all stages of working (FP1 3.1). Functions stay within P1/P2 content.

1 Key knowledge and worked examples

Three locators. A sign change of continuous f on $[a, b]$ brackets a root. **Bisection** halves the bracket each time; **linear interpolation** uses the chord: $\beta \approx a - \frac{(b-a)f(a)}{f(b)-f(a)}$; **Newton–Raphson** (booklet) iterates $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$. Calculator-only answers score zero – every substitution is shown.

Worked example – bisection (WFM01 June 2025 Q3i). $f(x) = x^2 + 5 - 8^{5x}$ has a single root α in $[0, 1]$: $f(0) = 4 > 0$; $f(0.5) = 5.25 - 8^{2.5} = -175.77 < 0$ so $\alpha \in [0, 0.5]$; midpoint 0.25: $f(0.25) = 5.0625 - 8^{1.25} = -8.392 < 0$, so the root lies in $[0, 0.25]$ – an interval of width 0.25.

Worked example – interpolation on a transcendental (WFM01 June 2025 Q3ii). $g(x) = 3^{\sin x} - 3 \cos x$: $g(4) = 2.396$, $g(5) = -0.502$. Both continuous and sign change, so a root $\beta \in [4, 5]$. One interpolation: $\beta \approx 4 + \frac{(5-4) \times 2.396}{2.396 - (-0.502)} = 4 + \frac{2.396}{2.898} = 4.827$ (4 s.f.).

Worked example – Newton–Raphson (original). $x^3 - 2x - 5 = 0$ from $x_0 = 2$: $f'(x) = 3x^2 - 2$; $x_1 = 2 - \frac{f(2)}{f'(2)} = 2 - \frac{-1}{10} = 2.1$; $x_2 = 2.1 - \frac{0.061}{11.23} = 2.0946$ (true root 2.09455...).

2 Practice

2.1 Show that $x^3 + x - 3 = 0$ has a root between 1 and 2, then perform two bisections to locate it in an interval of width 0.25. [4]

2.2 Use one linear interpolation on $[2, 3]$ for $h(x) = x + \frac{1}{x} - 3$ (take $h(2) = -0.5$, $h(3) = 0.3\bar{3}$) to estimate the root to 3 significant figures. [3]

2.3 Apply two Newton–Raphson iterations to $x^3 - 3x - 1 = 0$ starting from $x_0 = 2$, giving answers to 4 decimal places. [4]

2.4 (Original.) A student’s bisection table shows $f(1.6) = -0.3$ and $f(1.8) = 0.2$, and concludes “root = 1.7”. Give two criticisms. [2]

3 Exam-style questions

3.1 (*Original.*) $p(x) = x^4 - 3x - 1$. (a) Show $p(x) = 0$ has a root γ in $[1, 2]$. (b) Use interval bisection twice to find an interval of width 0.25 containing γ . (c) Use one linear interpolation on your interval from (b) to estimate γ to 3 significant figures. [7]

3.2 (*Adapted from Edexcel IAL WFM01 June 2025 Q3ii.*) $g(x) = x \cos \frac{x}{2} + 1$, x in radians. (a) Show that $g(x) = 0$ has a root β in the interval $[3, 4]$, making the continuity argument explicit. (b) Use linear interpolation once on $[3, 4]$ to estimate β to 4 significant figures ($g(3) = 1.2122\dots$, $g(4) = -0.6646\dots$; use these values). [4]
