

Exercise walk-through

8.2

Continuous random variables

Pearson Edexcel · International A-Level · Further Mathematics

Check yourself

Check yourself

State distribution assumptions in context; solve moment pairs; scale Poisson rates; normalise piecewise densities; medians, modes and skewness; uniform facts both directions; census versus sample; statistics; small sampling distributions; one- and two-tailed tests with critical regions and actual levels; continuity-corrected normal thresholds.

Exercises 2.1·2.3

2.1 $f(x) = kx^2$ on $[0, 2]$. Find k , $E(X)$ and $\text{Var}(X)$. [4]

2.2 For the same variable, find the median. [2]

2.3 $f(x) = \frac{2x}{9}$ on $[0, 3]$. Find $P(X > 1)$ and $E(X)$. [3]

Exercises 2.1·2.3

2.1 $f(x) = kx^2$ on $[0, 2]$. Find k , $E(X)$ and $\text{Var}(X)$. [4]

$$\mathbf{2.1} \int_0^2 kx^2 dx = \frac{8k}{3} = 1 \text{ gives } k = \frac{3}{8}. E(X) = \frac{3}{8} \int_0^2 x^3 dx = \frac{3}{8}(4) = \frac{3}{2}; E(X^2) = \frac{3}{8} \int_0^2 x^4 dx = \frac{3}{8} \cdot \frac{32}{5} = \frac{12}{5}; \text{Var}(X) = \frac{12}{5} - \frac{9}{4} = \frac{3}{20}.$$

2.2 For the same variable, find the median. [2]

$$\mathbf{2.2} F(m) = \frac{3}{8} \cdot \frac{m^3}{3} = \frac{m^3}{8} = \frac{1}{2}: m^3 = 4, \text{ so } m = 1.59 \text{ (3 s.f.)}.$$

2.3 $f(x) = \frac{2x}{9}$ on $[0, 3]$. Find $P(X > 1)$ and $E(X)$. [3]

$$\mathbf{2.3} F(x) = \frac{x^2}{9}: P(X > 1) = 1 - \frac{1}{9} = \frac{8}{9}. E(X) = \frac{2}{9} \int_0^3 x^2 dx = \frac{2}{9}(9) = 2.$$

Exercise 3.1

3.1 (*Edexcel IAL WST02 June 2025 Q1, condensed.*) The continuous random variable X has cumulative distribution function $F(x) = k(6x^2 - x^4)$ for $0 \leq x \leq 1$ (0 below, 1 above). Given also that $E(X) = 0.64$: (a) show that $k = \frac{1}{5}$; (b) find $P(X > 0.6)$; (c) show that the median m satisfies $2m^4 - 12m^2 + 5 = 0$ and hence find m ; (d) find, using calculus, the mode of X ; (e) state, with a reason, the skewness of X . [11]

Exercise 3.1

3.1 (*Edexcel IAL WST02 June 2025 Q1, condensed.*) The continuous random variable X has cumulative distribution function $F(x) = k(6x^2 - x^4)$ for $0 \leq x \leq 1$ (0 below, 1 above). Given also that $E(X) = 0.64$: (a) show that $k = \frac{1}{5}$; (b) find $P(X > 0.6)$; (c) show that the median m satisfies $2m^4 - 12m^2 + 5 = 0$ and hence find m ; (d) find, using calculus, the mode of X ; (e) state, with a reason, the skewness of X . [11]

3.1 (a) $F(1) = k(6 - 1) = 5k = 1$, so $k = \frac{1}{5}$ ✓. (b) $F(0.6) = \frac{6(0.36) - 0.6^4}{5} = \frac{2.0304}{5} = 0.40608$, so $P(X > 0.6) = 0.59392$ (awrt 0.594). (c) $F(m) = \frac{1}{2}$: $6m^2 - m^4 = \frac{5}{2}$, i.e. $2m^4 - 12m^2 + 5 = 0$ ✓. Solving, $m^2 = \frac{12 - \sqrt{144 - 40}}{4} = 3 - \frac{\sqrt{26}}{2} = 0.4505$, so $m = 0.671$ (the root inside $[0, 1]$). (d) $f(x) = F'(x) = \frac{12x - 4x^3}{5}$; $f'(x) = \frac{12 - 12x^2}{5} = 0$ gives $x = \pm 1$ – only $x = 1$ lies in the support (where $f(1) = \frac{8}{5} > 0$), so the mode is 1. (e) mean $0.64 <$ median $0.671 <$ mode 1: negative skew.

Exercise 3.2

3.2 (*Original.*) $f(x) = kx^2(1 - x)$ on $[0, 1]$. (a) Show that $k = 12$. (b) Find $E(X)$ and $\text{Var}(X)$. (c) Find the mode of X . [6]

Exercise 3.2

3.2 (*Original.*) $f(x) = kx^2(1 - x)$ on $[0, 1]$. (a) Show that $k = 12$. (b) Find $E(X)$ and $\text{Var}(X)$. (c) Find the mode of X . [6]

3.2 (a) $\int_0^1 x^2(1 - x)dx = \frac{1}{3} - \frac{1}{4} = \frac{1}{12}$, so $k = 12$ ✓. (b) $E(X) = 12 \int_0^1 (x^3 - x^4)dx = 12(\frac{1}{4} - \frac{1}{5}) = \frac{3}{5}$; $E(X^2) = 12(\frac{1}{5} - \frac{1}{6}) = \frac{2}{5}$; $\text{Var}(X) = \frac{2}{5} - \frac{9}{25} = \frac{1}{25}$. (c) $f(x) = 12(x^2 - x^3)$; $f'(x) = 12(2x - 3x^2) = 0$ gives $x = \frac{2}{3}$ (and $x = 0$, an endpoint): the mode is $\frac{2}{3}$.

Local review guidance: normalisation over the correct limits; quartic medians solved for the root INSIDE the support; modes rejected outside the support. This is authored classroom guidance.