

Exercise walk-through

9.1

Sampling methods

Pearson Edexcel · International A-Level · Further Mathematics

1. Unit S3 at a glance

Unit S3 at a glance

WST03/01: 1 hour 30 minutes, 75 marks; the booklet carries the Normal and chi-squared tables plus Spearman critical values. Must-know:

$$aX \pm bY \sim N(a\mu_x \pm b\mu_y, a^2\sigma_x^2 + b^2\sigma_y^2); \bar{X} \sim N(\mu, \sigma^2/n); s^2 = \frac{\sum x^2 - n\bar{x}^2}{n-1};$$

$$\chi^2 = \sum \frac{(O-E)^2}{E}; r_s = 1 - \frac{6 \sum d^2}{n(n^2-1)}.$$

Check yourself

Check yourself

Combine independent Normals (differences and absolute differences); four sampling methods with mechanical reasons; bias algebra and paired unbiased-estimator equations; recovering (μ, σ) from two sample-mean probabilities; z intervals and one/two-sample tests with the variance assumption; chi-squared for fit and contingency with df bookkeeping; Spearman's r_s and its table test.

Exercises 2.1·2.3

2.1 A factory has 540 items listed serially; a systematic sample of 18 is needed and item 7 is chosen first. Write down x and the first four items. [2]

2.2 A school of 800 pupils has 480 girls and 320 boys. A stratified sample of 40 is needed: how many of each? [2]

2.3 State one advantage and one disadvantage of quota sampling. [2]

Exercises 2.1·2.3

2.1 A factory has 540 items listed serially; a systematic sample of 18 is needed and item 7 is chosen first. Write down x and the first four items. [2]

2.1 $x = 540/18 = 30$: items 7, 37, 67, 97.

2.2 A school of 800 pupils has 480 girls and 320 boys. A stratified sample of 40 is needed: how many of each? [2]

2.2 Girls $40 \times \frac{480}{800} = 24$; boys $40 \times \frac{320}{800} = 16$.

2.3 State one advantage and one disadvantage of quota sampling. [2]

2.3 Advantage: no sampling frame needed (cheap, quick). Disadvantage: interviewer bias – the choice of who to approach is not random.

Exercise 3.1

3.1 (*Edexcel IAL WST03 June 2025 Q1.*) A golf club has 360 members. The captain needs to select 20 players for a charity tournament. The members are listed by randomly allocated membership numbers and the captain uses systematic sampling. The members are numbered 001 to 360; the 5th member is randomly selected first and then every x th member thereafter. (a) Give a reason for using systematic sampling in this case. (b) Write down the value of x . (c) Find the probability that the sample includes both the 5th member and the 6th member. [3]

Exercise 3.1

3.1 (*Edexcel IAL WST03 June 2025 Q1.*) A golf club has 360 members. The captain needs to select 20 players for a charity tournament. The members are listed by randomly allocated membership numbers and the captain uses systematic sampling. The members are numbered 001 to 360; the 5th member is randomly selected first and then every x th member thereafter. (a) Give a reason for using systematic sampling in this case. (b) Write down the value of x . (c) Find the probability that the sample includes both the 5th member and the 6th member. [3]

3.1 (a) It removes bias/favouritism: with the random start, every member has an equal chance of selection. (b) $x = \frac{360}{20} = 18$. (c) The sample is 5, 23, 41, ..., 347 – members 18 apart; 5 and 6 are consecutive, so both cannot be in the sample: probability 0.

Exercise 3.2

3.2 (*Original.*) A market researcher interviews shoppers outside a store, required to speak to 25 men and 25 women whatever order they arrive. (a) Name the sampling method. (b) State why the sample may still be biased even though the quotas are met. (c) The researcher instead decides to select from a list of 2000 store members a sample of 50, where every member is equally likely: name this method and state one requirement for implementing it. [6]

Exercise 3.2

3.2 (*Original.*) A market researcher interviews shoppers outside a store, required to speak to 25 men and 25 women whatever order they arrive. (a) Name the sampling method. (b) State why the sample may still be biased even though the quotas are met. (c) The researcher instead decides to select from a list of 2000 store members a sample of 50, where every member is equally likely: name this method and state one requirement for implementing it. [6]

3.2 (a) Quota sampling. (b) The interviewer chooses who to approach – the selection within each quota is non-random, so the sample may not represent all shoppers (interviewer bias). (c) Simple random sampling; it requires a sampling frame (the full list of members) and random-number selection from it.

Local review guidance: reasons tied to the method's mechanics, not generic speed claims; spacing as an exact division; impossibility arguments expressed through the spacing. This is authored classroom guidance.