

Solutions

2.1 The other root is $2 - i$: sum $4 = -p$, product $5 = q$. So $p = -4$, $q = 5$.

2.2 $f(1) = 1 - 3 + 7 - 5 = 0$ ✓. Divide: $x^3 - 3x^2 + 7x - 5 = (x - 1)(x^2 - 2x + 5)$.
 $x^2 - 2x + 5 = 0 \Rightarrow x = \frac{2 \pm \sqrt{4 - 20}}{2} = 1 \pm 2i$. Roots: 1, $1 \pm 2i$. (Product: $1 \cdot 5 = 5$ ✓.)

2.3 $1 - 2i$ is also a root, giving the factor $x^2 - 2x + 5$. Dividing: $\frac{x^4 - 2x^3 + 6x^2 - 2x + 5}{x^2 - 2x + 5} = x^2 + 1$, since $(x^2 - 2x + 5)(x^2 + 1) = x^4 - 2x^3 + 6x^2 - 2x + 5$ ✓. The other roots are $\pm i$.

2.4 Product of roots $= (1 + 3i)(1 - 3i) \cdot k^2 = 10k^2$. For a monic quartic the product equals the constant term (sign + for degree 4): $10k^2 = 50 \Rightarrow k = \pm\sqrt{5}$.

3.1 (a) Sum 1, product $\frac{1}{4} + \frac{9}{4} = \frac{5}{2}$: factor $z^2 - z + \frac{5}{2}$, or $2z^2 - 2z + 5$ scaled. (b) Comparing $(2z^2 - 2z + 5)(z^2 + mz + n) = 2z^4 + 2z^3 + 7z^2 + 4z + 15$: constant $5n = 15 \Rightarrow n = 3$; z^3 : $2m - 2 = 2 \Rightarrow m = 2$. Check the rest: $(2z^2 - 2z + 5)(z^2 + 2z + 3) = 2z^4 + 2z^3 + 7z^2 + 4z + 15$ ✓. (c) $z^2 + 2z + 3 = 0$ gives $z = -1 \pm i\sqrt{2}$, so the roots are $\frac{1}{2} \pm \frac{3}{2}i$, $-1 \pm i\sqrt{2}$.

3.2 Factor $z^2 - 6z + 34$; comparing coefficients: $c = 2$, $b = 6$, $P = 7$, $Q = 204$; other roots from $7z^2 + 6z + 2 = 0$: $z = \frac{-6 \pm \sqrt{-20}}{14} = \frac{-3 \pm i\sqrt{5}}{7}$ (and the conjugate $3 - 5i$).

Local review guidance: conjugate stated and the quadratic factor formed from sum and product; coefficient comparison or division carried through exactly; complex roots simplified to $a + bi$ form. This is authored classroom guidance.