

9.3 Confidence intervals and tests for means

Name: _____ Class: _____ Date: _____

Total: 26 marks

KEY WORDS confidence interval • critical value

Confidence intervals and tests for means

Objective

Build and interpret z-based confidence intervals, run one- and two-sample tests for a mean (including a non-zero null gap), and state the variance/CLT assumptions (objectives 3.3–3.8).

1 Key knowledge and worked examples

Formulas. CI: $\bar{x} \pm z\sigma/\sqrt{n}$. One-sample test: $z = \frac{\bar{x} - \mu}{\sigma/\sqrt{n}}$. Two samples with a gap:
 $z = \frac{(\bar{x} - \bar{y}) - (\mu_x - \mu_y)}{\sqrt{\sigma_x^2/n_x + \sigma_y^2/n_y}}$. Large samples: substitute s^2 for σ^2 (CLT for non-Normal populations).

Worked example (WST03 June 2025 Q4). Grey squirrels ($n = 80$, $\bar{x} = 500$, $s_x^2 = 1156$); red squirrels ($n = 80$, $\sum y = 23200$, $\sum y^2 = 6741351$): $\bar{y} = 290$, $s_y^2 = \frac{6741351 - 80(290)^2}{79} = 169$. Testing $H_0: \mu_x - \mu_y = 200$ vs $H_1: > 200$: $z = \frac{10}{\sqrt{265/16}} = 2.457 > 1.6449$ – reject H_0 . Assumption: the sample variances equal the population variances.

2 Practice

2.1 A sample of 36 from a Normal population with $\sigma = 9$ gives $\bar{x} = 50$. Construct a 95% confidence interval for μ . [3]

3.2 (*Original.*) A machine fills bottles; the fill volumes are Normally distributed with standard deviation 4 ml. A sample of 49 bottles gives $\bar{x} = 503$ ml. (a) Construct a 99% confidence interval for the mean fill. (b) The label claims 500 ml: test at the 1% level whether the mean fill exceeds 500 ml, and state the conclusion in context. [7]
