

7.2 Centres of mass

Name: _____ Class: _____ Date: _____

Total: 23 marks

KEY WORDS centre of mass • wire framework

Centres of mass

Objective

Find centres of mass of particle systems, uniform laminas, composite shapes (negative areas) and wire frameworks (length weighting with signed distances), and use suspension geometry (objectives 2.1–2.3).

1 Key knowledge and worked examples

Table method. Weight each piece by mass – for a uniform lamina by AREA, for a wire by LENGTH – and take moments about a convenient line: $\sum(\text{weight})(\bar{x}_{\text{piece}}) = (\text{total})(\bar{x})$. Holes and opposite-side pieces enter NEGATIVE. Booklet results: semi-circular wire $\frac{2r}{\pi}$, semicircular lamina $\frac{4r}{3\pi}$ from the centre. Suspended from a point, the body hangs with G vertically below it; the asked angle usually splits a triangle angle at the suspension point.

Worked example – an S-shaped framework (WME02 June 2025 Q3). Arcs AB ($r = 2a$, length $2\pi a$, COM $\frac{4a}{\pi}$ one side) and CD ($r = a$, length πa , COM $\frac{2a}{\pi}$ the OTHER side) joined by straights $AC = DB = a$ on the line: moments about the line give $2\pi a \cdot \frac{4a}{\pi} - \pi a \cdot \frac{2a}{\pi} = 6a^2$; total length $(3\pi + 2)a$; so $\bar{x} = \frac{6a}{3\pi+2}$. Suspended from A with E on arc AB at $\angle AOE = 90^\circ$: G lies on OE, $\tan \angle OAG = \frac{3}{3\pi+2}$, and $\theta = 45^\circ - \angle OAG$ (since $OA = OE$), giving $\tan \theta = \frac{3\pi-1}{3\pi+5}$.

2 Practice

2.1 Particles of mass 2 kg, 3 kg and 5 kg sit at $x = 0, 4$ and 6 on the x-axis. Find their centre of mass. [2]

2.2 A uniform rectangular lamina 6 m by 4 m has a 2 m by 2 m square hole cut from one corner. Find the distance of the centre of mass from each outer edge adjacent to that corner. [4]

2.3 Three uniform rods form a uniform wire triangle with sides 5, 5 and 6. Find the distance of the centre of mass from the 6-side. [3]

3.1 (*Edexcel IAL WME02 June 2025 Q3, condensed.*) A uniform rigid wire framework consists of two semicircular arcs AB and CD, both of centre O, joined by two straight pieces AC and DB. The arc AB has radius $2a$, the arc CD has radius a , $AC = DB = a$, and ACDB is a straight line. The arcs lie on opposite sides of ACDB. The centre of mass of a semicircular arc of radius r is $\frac{2r}{\pi}$ from its centre. (a) Show that the centre of mass of the framework is $\frac{6a}{3\pi+2}$ from AB. (b) The point E lies on the arc AB with $\angle AOE = 90^\circ$. The framework is freely suspended from A and hangs in equilibrium with AE at angle θ to the downward vertical. Show that $\tan \theta = \frac{3\pi - 1}{3\pi + 5}$. [10]

3.2 (Original.) A uniform semicircular lamina of radius 6 cm is freely suspended from one end P of its straight edge. Find the angle between the straight edge and the vertical, giving an exact tangent. [4]